Calculus II
Homework
Building block integrals

1. Let $x \in (0,1)$. Express the following using x and $\sqrt{1-x^2}$.

   (a) $\sin(\arcsin(x))$.
   (b) $\sin(2\arcsin(x))$.
   (c) $\sin(3\arcsin(x))$.
   (d) $\sin(\arccos(x))$.
   (e) $\sin(2\arccos(x))$.
   (f) $\sin(3\arccos(x))$.
   (g) $\cos(2\arcsin(x))$.
   (h) $\cos(3\arccos(x))$.

2. Express as the following as an algebraic expression of x. In other words, "get rid" of the trigonometric and inverse trigonometric expressions.

   (a) $\cos^2(\arctan x)$.
   (b) $-\sin^2(\operatorname{arccot} x)$.
   (c) $\dfrac{1}{\cos(\arcsin x)}$.
   (d) $-\dfrac{1}{\sin(\arccos x)}$.

3. Rewrite as a rational function of t. This problem will be later used to derive the Euler substitutions (an important technique for integrating).

   (a) $\cos(2\arctan t)$.
   (b) $\sin(2\arctan t)$.
   (c) $\tan(2\arctan t)$.
   (d) $\cot(2\arctan t)$.
   (e) $\csc(2\arctan t)$.
   (f) $\sec(2\arctan t)$.
   (g) $\cos(2\operatorname{arccot} t)$.
   (h) $\sin(2\operatorname{arccot} t)$.
   (i) $\tan(2\operatorname{arccot} t)$.
   (j) $\cot(2\operatorname{arccot} t)$.
   (k) $\csc(2\operatorname{arccot} t)$.
   (l) $\sec(2\operatorname{arccot} t)$.

4. Compute the derivative (derive the formula).

   (a) $(\arctan x)'$.
   (b) $(\operatorname{arccot} x)'$.
   (c) $(\arcsin x)'$.
   (d) $(\arccos x)'$.
   (e) Let arcsec denote the inverse of the secant function. Compute $(\operatorname{arcsec} x)'$.

5. (a) Let $a+b \neq k\pi$, $a \neq k\pi + \frac{\pi}{2}$ and $b \neq k\pi + \frac{\pi}{2}$ for any $k \in \mathbb{Z}$ (integers). Prove that

   $$\frac{\tan a + \tan b}{1 - \tan a \tan b} = \tan(a+b) \quad .$$

   (b) Let x and y be real. Prove that, for $xy \neq 1$, we have

   $$\arctan x + \arctan y = \arctan\left(\frac{x+y}{1-xy}\right)$$

   if the left hand side lies between $\left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.

6. Evaluate the indefinite integral. Illustrate the steps of your solutions.

(a) $\int x \sin x \mathrm{d}x$.

(b) $\int x e^{-x} \mathrm{d}x$.

(c) $\int x^2 e^{x} \mathrm{d}x$.

(d) $\int x \sin(-2x) \mathrm{d}x$.

(e) $\int x^2 \cos(3x) \mathrm{d}x$.

(f) $\int x^2 e^{-2x} \mathrm{d}x$.

(g) $\int x \sin(2x) \mathrm{d}x$.

(h) $\int x \cos(3x) \mathrm{d}x$.

(i) $\int x^2 e^{2x} \mathrm{d}x$.

(j) $\int x^3 e^{x} \mathrm{d}x$.

7. Evaluate the indefinite integral. Illustrate the steps of your solutions.

(a) $\int x^2 \cos(2x) \mathrm{d}x$.

(b) $\int x^2 e^{ax} \mathrm{d}x$, where a is a constant.

(c) $\int x^2 e^{-ax} \mathrm{d}x$, where a is a constant.

(d) $\int x^2 \dfrac{(e^{ax}+e^{-ax})^2}{4} \mathrm{d}x$, where a is a constant.

(e) $\int \dfrac{1}{\cos^2 x} \mathrm{d}x$. (Hint: This problem does not require integration by parts. What is the derivative of $\tan x$?)

(f) $\int (\tan^2 x) \mathrm{d}x$. (Hint: This problem does not require integration by parts. We can use $\tan^2 x = \frac{1}{\cos^2 x} - 1$ and the previous problem.)

(g) $\int x \tan^2 x \mathrm{d}x$. (Hint: $\tan^2 x \mathrm{d}x = \mathrm{d}(F(x))$, where $F(x)$ is the answer from the preceding problem).

(h) $\int e^{-\sqrt{x}} \mathrm{d}x$.

(i) $\int \cos^2 x \ \mathrm{d}x$.

(j) $\int \dfrac{x}{1+x^2} \mathrm{d}x$ (Hint: use substitution rule, don't use integration by parts)

(k) $\int (\arctan x) \mathrm{d}x$.

(l) $\int (\arcsin x) \mathrm{d}x$.

(m) $\int (\arcsin x)^2 \mathrm{d}x$. (Hint: Try substituting $x = \sin y$.)

(n) $\int \arctan\left(\dfrac{1}{x}\right) \mathrm{d}x$.

(o) $\int \sin x e^{x} \mathrm{d}x$

(p) $\int \cos x e^{x} \mathrm{d}x$

(q) $\int \sin(\ln(x)) \mathrm{d}x$.

(r) $\int \cos(\ln(x)) \mathrm{d}x$.

(s) $\int \ln x \mathrm{d}x$

(t) $\int x \ln x \ \mathrm{d}x$.

(u) $\int \dfrac{\ln x}{\sqrt{x}} \mathrm{d}x$.

(v) $\int (\ln x)^2 \mathrm{d}x$.

(w) $\int (\ln x)^3 \mathrm{d}x$.

(x) $\int x^2 \cos^2 x \mathrm{d}x$. (This problem is related to Problem 7.d as $\cos x = \frac{e^{ix}+e^{-ix}}{2}$).

8. Compute $\int x^n e^x \mathrm{d}x$, where n is a non-negative integer.

9. Integrate. Illustrate the steps of your solution.

(a) $\int \dfrac{1}{x+1} \mathrm{d}x$

(b) $\int \dfrac{x-1}{x+1} \mathrm{d}x$

(c) $\int \dfrac{1}{(x+1)^2} \mathrm{d}x$

(d) $\int \dfrac{x}{(x+1)^2} \mathrm{d}x$

(e) $\int \dfrac{1}{(2x+3)^2} \mathrm{d}x$

(f) $\int \dfrac{x}{2x^2+3} \mathrm{d}x$

(g) $\int \dfrac{1}{2x^2+3} \mathrm{d}x$

(h) $\int \dfrac{x}{2x^2+x+1} \mathrm{d}x$.

(i) $\int \dfrac{x}{2x^2+x+3} \mathrm{d}x$

(j) $\int \dfrac{x}{x^2-x+3} \mathrm{d}x$

(k) $\displaystyle\int \frac{1}{\left(x^2+1\right)^2}\mathrm{d}x$

(l) $\displaystyle\int \frac{1}{\left(x^2+x+1\right)^2}\mathrm{d}x$

(m) $\displaystyle\int \frac{1}{\left(x^2+1\right)^3}\mathrm{d}x$

10. Let a, b, c, A, B be real numbers. Suppose in addition $a \neq 0$ and $b^2 - 4ac < 0$. Integrate

$$\int \frac{Ax+B}{ax^2+bx+c}\mathrm{d}x \quad .$$

The purpose of this exercise is to produce a formula in form ready for implementation in a computer algebra system.

11. Let a, b, c, A, B be real numbers and let $n > 1$ be an integer. Suppose in addition $a \neq 0$ and $b^2 - 4ac < 0$. Let

$$J(n) = \int \frac{1}{\left(x^2+\frac{b}{a}x+\frac{c}{a}\right)^n}\mathrm{d}x \quad .$$

(a) Express the integral

$$\int \frac{Ax+B}{\left(ax^2+bx+c\right)^n}\mathrm{d}x$$

via $J(n)$.

(b) Express $J(n)$ recursively via $J(n-1)$

The purpose of this exercise is to produce a formula in form ready for implementation in a computer algebra system.