

## Calculus II

# Curve length miscellaneous problem, part 2

Todor Milev

2019

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] \quad .$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] \quad .$$

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = ?$  and  $y'(t) = ?$

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left( ? \right)^2 + \left( ? \right)^2} dt$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = ?$

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + ( ? )^2} dt$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = ?$

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + ( ? )^2} dt$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] \quad .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}}$

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(\frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}}\right)^2} dt$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] \quad .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4} t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + (\text{?})^2} dt$$



## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] \quad .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\text{?} + \text{?}} dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + ?} dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + ?} dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4] .$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left| \frac{1}{2\sqrt{t}} + 2 \right| dt \end{aligned}$$



## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[ \frac{1}{\sqrt{t}} + 2t \right]_1^4 \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[ \sqrt{t} + 2t \right]_1^4 \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[\sqrt{t} + 2t\right]_1^4 \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[\sqrt{t} + 2t\right]_1^4 \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma : \begin{cases} x(t) = \sqrt{t} - 2 \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[\sqrt{t} + 2t\right]_1^4 = \sqrt{4} + 2 \cdot 4 - (\sqrt{1} + 2 \cdot 1) \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[ \sqrt{t} + 2t \right]_1^4 = \sqrt{4} + 2 \cdot 4 - (\sqrt{1} + 2 \cdot 1) \end{aligned}$$

## Example

Find the length of the curve  $\gamma$ .

$$\gamma: \begin{cases} x(t) = \sqrt{t} - 2t \\ y(t) = \frac{8}{3}t^{\frac{3}{4}} \end{cases}, t \in [1, 4]$$

We have that  $x'(t) = \frac{1}{2\sqrt{t}} - 2$  and  $y'(t) = \frac{8}{3} \cdot \frac{3}{4}t^{-\frac{1}{4}} = 2t^{-\frac{1}{4}}$ .

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} - 2\right)^2 + \left(2t^{-\frac{1}{4}}\right)^2} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} - \frac{2}{\sqrt{t}} + 4 + \frac{4}{\sqrt{t}}} dt \\ &= \int_1^4 \sqrt{\frac{1}{4t} + \frac{2}{\sqrt{t}} + 4} dt = \int_1^4 \sqrt{\left(\frac{1}{2\sqrt{t}} + 2\right)^2} dt \\ &= \int_1^4 \left(\frac{1}{2\sqrt{t}} + 2\right) dt = \left[\sqrt{t} + 2t\right]_1^4 = \sqrt{4} + 2 \cdot 4 - (\sqrt{1} + 2 \cdot 1) = 7. \end{aligned}$$