

Calculus II

Homework

Integrals of involving radicals of quadratics

1. Let $x \in (0, 1)$. Express the following using x and $\sqrt{1 - x^2}$.

- | | |
|----------------------------|----------------------------|
| (a) $\sin(\arcsin(x))$. | (e) $\sin(2 \arccos(x))$. |
| (b) $\sin(2 \arcsin(x))$. | (f) $\sin(3 \arccos(x))$. |
| (c) $\sin(3 \arcsin(x))$. | (g) $\cos(2 \arcsin(x))$. |
| (d) $\sin(\arccos(x))$. | (h) $\cos(3 \arccos(x))$. |

2. Express as the following as an algebraic expression of x . In other words, “get rid” of the trigonometric and inverse trigonometric expressions.

- | | |
|--|------------------------------------|
| (a) $\cos^2(\arctan x)$. | (c) $\frac{1}{\cos(\arcsin x)}$. |
| (b) $-\sin^2(\operatorname{arccot} x)$. | (d) $-\frac{1}{\sin(\arccos x)}$. |

3. Rewrite as a rational function of t . This problem will be later used to derive the Euler substitutions (an important technique for integrating).

- | | |
|---------------------------|---|
| (a) $\cos(2 \arctan t)$. | (g) $\cos(2 \operatorname{arccot} t)$. |
| (b) $\sin(2 \arctan t)$. | (h) $\sin(2 \operatorname{arccot} t)$. |
| (c) $\tan(2 \arctan t)$. | (i) $\tan(2 \operatorname{arccot} t)$. |
| (d) $\cot(2 \arctan t)$. | (j) $\cot(2 \operatorname{arccot} t)$. |
| (e) $\csc(2 \arctan t)$. | (k) $\csc(2 \operatorname{arccot} t)$. |
| (f) $\sec(2 \arctan t)$. | (l) $\sec(2 \operatorname{arccot} t)$. |

4. Compute the derivative (derive the formula).

- | | |
|------------------------------------|---|
| (a) $(\arctan x)'$. | (d) $(\arccos x)'$. |
| (b) $(\operatorname{arccot} x)'$. | (e) Let arcsec denote the inverse of the secant function. Compute $(\operatorname{arcsec} x)'$. |
| (c) $(\arcsin x)'$. | |

5. (a) Let $a + b \neq k\pi$, $a \neq k\pi + \frac{\pi}{2}$ and $b \neq k\pi + \frac{\pi}{2}$ for any $k \in \mathbb{Z}$ (integers). Prove that

$$\frac{\tan a + \tan b}{1 - \tan a \tan b} = \tan(a + b) \quad .$$

(b) Let x and y be real. Prove that, for $xy \neq 1$, we have

$$\arctan x + \arctan y = \arctan \left(\frac{x + y}{1 - xy} \right)$$

if the left hand side lies between $(-\frac{\pi}{2}, \frac{\pi}{2})$.

6. Evaluate the indefinite integral. Illustrate the steps of your solutions.

$$(a) \int x \sin x dx.$$

$$(b) \int x e^{-x} dx.$$

$$(c) \int x^2 e^x dx.$$

$$(d) \int x \sin(-2x) dx.$$

$$(e) \int x^2 \cos(3x) dx.$$

$$(f) \int x^2 e^{-2x} dx.$$

$$(g) \int x \sin(2x) dx.$$

$$(h) \int x \cos(3x) dx.$$

$$(i) \int x^2 e^{2x} dx.$$

$$(j) \int x^3 e^x dx.$$

7. Evaluate the indefinite integral. Illustrate the steps of your solutions.

$$(a) \int x^2 \cos(2x) dx.$$

$$(b) \int x^2 e^{ax} dx, \text{ where } a \text{ is a constant.}$$

$$(c) \int x^2 e^{-ax} dx, \text{ where } a \text{ is a constant.}$$

$$(d) \int x^2 \frac{(e^{ax} + e^{-ax})^2}{4} dx, \text{ where } a \text{ is a constant.}$$

$$(e) \int \frac{1}{\cos^2 x} dx. \quad (\text{Hint: This problem does not require integration by parts. What is the derivative of } \tan x?)$$

$$(f) \int (\tan^2 x) dx. \quad (\text{Hint: This problem does not require integration by parts. We can use } \tan^2 x = \frac{1}{\cos^2 x} - 1 \text{ and the previous problem.})$$

$$(g) \int x \tan^2 x dx. \quad (\text{Hint: } \tan^2 x dx = d(F(x)), \text{ where } F(x) \text{ is the answer from the preceding problem}).$$

$$(h) \int e^{-\sqrt{x}} dx.$$

$$(i) \int \cos^2 x dx.$$

$$(j) \int \frac{x}{1+x^2} dx \quad (\text{Hint: use substitution rule, don't use integration by parts})$$

$$(k) \int (\arctan x) dx.$$

$$(l) \int (\arcsin x) dx.$$

$$(m) \int (\arcsin x)^2 dx. \quad (\text{Hint: Try substituting } x = \sin y.)$$

$$(n) \int \arctan\left(\frac{1}{x}\right) dx.$$

$$(o) \int \sin x e^x dx$$

$$(p) \int \cos x e^x dx$$

$$(q) \int \sin(\ln(x)) dx.$$

$$(r) \int \cos(\ln(x)) dx.$$

$$(s) \int \ln x dx$$

$$(t) \int x \ln x dx.$$

$$(u) \int \frac{\ln x}{\sqrt{x}} dx.$$

$$(v) \int (\ln x)^2 dx.$$

$$(w) \int (\ln x)^3 dx.$$

$$(x) \int x^2 \cos^2 x dx. \quad (\text{This problem is related to Problem 7.d as } \cos x = \frac{e^{ix} + e^{-ix}}{2}).$$

8. Compute $\int x^n e^x dx$, where n is a non-negative integer.

9. Integrate. Illustrate the steps of your solution.

$$(a) \int \frac{1}{x+1} dx$$

$$(b) \int \frac{x-1}{x+1} dx$$

$$(c) \int \frac{1}{(x+1)^2} dx$$

$$(d) \int \frac{x}{(x+1)^2} dx$$

$$(e) \int \frac{1}{(2x+3)^2} dx$$

$$(f) \int \frac{x}{2x^2+3} dx$$

$$(g) \int \frac{1}{2x^2+3} dx$$

$$(h) \int \frac{x}{2x^2+x+1} dx.$$

$$(i) \int \frac{x}{2x^2+x+3} dx$$

$$(j) \int \frac{x}{x^2-x+3} dx$$

$$(k) \int \frac{1}{(x^2 + 1)^2} dx$$

$$(m) \int \frac{1}{(x^2 + 1)^3} dx$$

$$(l) \int \frac{1}{(x^2 + x + 1)^2} dx$$

10. Let a, b, c, A, B be real numbers. Suppose in addition $a \neq 0$ and $b^2 - 4ac < 0$. Integrate

$$\int \frac{Ax + B}{ax^2 + bx + c} dx \quad .$$

The purpose of this exercise is to produce a formula in form ready for implementation in a computer algebra system.

11. Let a, b, c, A, B be real numbers and let $n > 1$ be an integer. Suppose in addition $a \neq 0$ and $b^2 - 4ac < 0$. Let

$$J(n) = \int \frac{1}{(x^2 + \frac{b}{a}x + \frac{c}{a})^n} dx \quad .$$

(a) Express the integral

$$\int \frac{Ax + B}{(ax^2 + bx + c)^n} dx$$

via $J(n)$.

(b) Express $J(n)$ recursively via $J(n - 1)$

The purpose of this exercise is to produce a formula in form ready for implementation in a computer algebra system.

12. Integrate. Some of the examples require partial fraction decomposition and some do not. Illustrate the steps of your solution.

$$(a) \int \frac{1}{4x^2 + 4x + 1} dx$$

$$(h) \int \frac{x}{3x^2 + x - 2} dx$$

$$(b) \int \frac{1}{1 - x^2} dx$$

$$(i) \int \frac{x}{3x^2 + x + 2} dx$$

$$(c) \int \frac{1}{5 - x^2} dx$$

$$(j) \int \frac{x}{2x^2 + x + 1} dx$$

$$(d) \int \frac{x}{4x^2 + x + \frac{1}{16}} dx$$

$$(k) \int \frac{x}{2x^2 + x - 1} dx$$

$$(e) \int \frac{x + 1}{2x^2 + x} dx$$

$$(l) \int \frac{1}{x^2 + x + 1} dx$$

$$(f) \int \frac{x}{4x^2 + x + 5} dx$$

$$(m) \int \frac{1}{2x^2 + 5x + 1} dx$$

$$(g) \int \frac{x}{4x^2 + x - 5} dx$$

13. Evaluate the indefinite integral. Illustrate all steps of your solution.

$$(a) \int \frac{x^3 + 4}{x^2 + 4} dx$$

$$(h) \int \frac{15x^2 - 4x - 81}{(x - 3)(x + 4)(x - 1)} dx$$

$$(b) \int \frac{4x^2}{2x^2 - 1} dx$$

$$(i) \int \frac{x^4 + 10x^3 + 18x^2 + 2x - 13}{x^4 + 4x^3 + 3x^2 - 4x - 4} dx$$

$$(c) \int \frac{x^3}{x^2 + 2x - 3} dx$$

Check first that $(x - 1)(x + 2)^2(x + 1) = x^4 + 4x^3 + 3x^2 - 4x - 4$.

$$(d) \int \frac{x^3}{x^2 + 3x - 4} dx$$

$$(j) \int \frac{x^4}{(x^2 + 2)(x + 2)} dx$$

$$(e) \int \frac{x^3}{2x^2 + 3x - 5} dx$$

$$(k) \int \frac{x^5}{x^3 - 1} dx$$

$$(f) \int \frac{x^2 + 1}{(x - 3)(x - 2)^2} dx$$

$$(l) \int \frac{x^4}{(x^2 + 2)(x + 1)^2} dx$$

$$(g) \int \frac{x^4}{(x + 1)^2(x + 2)} dx$$

$$(m) \int \frac{3x^2 + 2x - 1}{(x-1)(x^2+1)} dx$$

$$(n) \int \frac{x^2 - 1}{x(x^2 + 1)^2} dx$$

14. Integrate

$$\int \frac{x^6 - x^5 + \frac{9}{2}x^4 - 4x^3 + \frac{13}{2}x^2 - \frac{7}{2}x + \frac{11}{4}}{x^5 - x^4 + 3x^3 - 3x^2 + \frac{9}{4}x - \frac{9}{4}} dx \quad .$$

15. Integrate.

$$(a) \int \frac{1}{3 + \cos x} dx.$$

$$(d) \int \frac{1}{2 + \tan x} dx. \text{ (Hint: this integral can be done simply with the substitution } x = \arctan t.)$$

$$(b) \int \frac{1}{4 + \cos x} dx.$$

$$(c) \int \frac{1}{3 + \sin x} dx.$$

$$(e) \int \frac{dx}{2 \sin x - \cos x + 5}.$$

16. Integrate. The answer key has not been proofread, use with caution.

$$(a) \int \sin(3x) \cos(2x) dx.$$

$$(b) \int \sin x \cos(5x) dx.$$

$$(c) \int \cos(3x) \sin(2x) dx.$$

$$(d) \int \sin(5x) \sin(3x) dx.$$

$$(e) \int \cos(x) \cos(3x) dx.$$

17. Integrate.

$$(a) \int \sin^2 x \cos x dx.$$

$$(c) \int \cos^3 x dx.$$

$$(b) \int \sin^2 x dx.$$

$$(d) \int \sin^3 x \cos^4 x dx.$$

18. Integrate.

$$(a) \int \sec x dx.$$

$$(b) \int \sec^3 x dx.$$

$$(c) \int \tan^3 x dx.$$

$$(d) \int \sec^2 x \tan^2 x dx.$$

19. Find a linear substitution (via completing the square) to transform the radical to a multiple of an expression of the form $\sqrt{u^2 + 1}$, $\sqrt{u^2 - 1}$ or $\sqrt{1 - u^2}$.

$$(a) \sqrt{x^2 + x + 1}.$$

$$(b) \sqrt{-2x^2 + x + 1}.$$

20. Compute the integral.

$$(a) \int \frac{\sqrt{1 + x^2}}{x^2} dx.$$

21. Compute the integral using a trigonometric substitution.

$$(a) \int \frac{\sqrt{9-x^2}}{x^2} dx \quad .$$

22. Compute the integral.

$$(a) \int \sqrt{x^2+1} dx$$

$$(b) \int \sqrt{x^2+2} dx$$

$$(c) \int \sqrt{x^2+x+1} dx$$

$$(d) \int \sqrt{(2x^2+2x+1)} dx$$

$$(e) \int \sqrt{(3x^2+2x+1)} dx$$

$$(f) \int \frac{\sqrt{x^2+1}}{x+1} dx$$

23. Let $b^2 - 4ac < 0$ and $a > 0$ be (real) numbers. Show that

$$\int \sqrt{(ax^2+bx+c)} dx = \frac{\sqrt{a}D}{2} \left(\ln \left(\sqrt{\left(\frac{2xa+b}{2\sqrt{Da}} \right)^2 + 1} + \frac{2xa+b}{2\sqrt{Da}} \right) + \frac{2xa+b}{2\sqrt{Da}} \sqrt{\left(\frac{2xa+b}{2\sqrt{Da}} \right)^2 + 1} \right) + C,$$

$$\text{where } D = \frac{4ac - b^2}{4a^2}.$$

24. Integrate

$$(a) \int \sqrt{1-x^2} dx$$

$$(b) \int \sqrt{2-x^2} dx$$

$$(c) \int \sqrt{-x^2+x+1} dx$$

$$(d) \int \sqrt{2-x-x^2} dx$$

$$(e) \int \frac{\sqrt{1-x^2}}{1+x} dx$$

$$(f) \int \frac{\sqrt{1-x^2}}{2+x} dx$$

25. Integrate

$$(a) \int \sqrt{x^2-1} dx$$

$$(b) \int \sqrt{x^2-2} dx$$

$$(c) \int \sqrt{2x^2+x-1} dx$$

$$(d) \int \sqrt{x^2+x-1} dx$$

26. (a) Express x , dx and $\sqrt{x^2+1}$ via θ and $d\theta$ for the trigonometric substitution $x = \cot \theta$, $\theta \in (0, \pi)$.

(b) Express x , dx and $\sqrt{x^2+1}$ via t and dt for the Euler substitution $x = \cot(2 \arctan t)$, $t > 0$. Express t via x .

27. Let the variables x and t be related via $\sqrt{x^2+1} = x+t$.

(a) Express x via t .

(b) Express $\sqrt{x^2+1}$ via t alone.

(c) Express dx via t and dt .

28. (a) Express x , dx and $\sqrt{x^2 + 1}$ via θ and $d\theta$ for the trigonometric substitution $x = \tan \theta$, $\theta \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$.
 (b) Express x , dx and $\sqrt{x^2 + 1}$ via t and dt for the Euler substitution $x = \tan(2 \arctan t)$, $t \in (-1, 1)$. Express t via x .
29. Let the variables x and t be related via $\sqrt{x^2 + 1} = \frac{x}{t} - 1$.
 (a) Express x via t .
 (b) Express $\sqrt{x^2 + 1}$ via t alone.
 (c) Express dx via t and dt .
30. (a) Express x , dx and $\sqrt{1 - x^2}$ via θ and $d\theta$ for the trigonometric substitution $x = \cos \theta$, $\theta \in [0, \pi]$.
 (b) Express x , dx and $\sqrt{1 - x^2}$ via t and dt for the Euler substitution $x = \cos(2 \arctan t)$, $t \geq 0$. Express t via x .
31. Let the variables x and t be related via $\sqrt{-x^2 + 1} = (1 - x)t$.
 (a) Express x via t .
 (b) Express $\sqrt{-x^2 + 1}$ via t alone.
 (c) Express dx via t and dt .
32. (a) Express x , dx and $\sqrt{1 - x^2}$ via θ and $d\theta$ for the trigonometric substitution $x = \sin \theta$, $\theta \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$.
 (b) Express x , dx and $\sqrt{1 - x^2}$ via t and dt for the Euler substitution $x = \sin(2 \arctan t)$, $t \in [-1, 1]$. Express t via x .
33. Let the variables x and t be related via $\sqrt{-x^2 + 1} = 1 - xt$.
 (a) Express x via t .
 (b) Express $\sqrt{-x^2 + 1}$ via t alone.
 (c) Express dx via t and dt .
34. (a) Express x , dx and $\sqrt{x^2 - 1}$ via θ and $d\theta$ for the trigonometric substitution $x = \csc \theta$, $\theta \in \left[0, \frac{\pi}{2}\right] \cup \left[\pi, \frac{3\pi}{2}\right]$.
 (b) Express x , dx and $\sqrt{x^2 - 1}$ via t and dt for the Euler substitution $x = \sec(2 \arctan t)$, $t \in (-\infty, -1) \cup [1, 0)$. Express t via x .
35. Let the variables x and t be related via $\sqrt{x^2 - 1} = (x + 1)t$.
 (a) Express x via t .
 (b) Express $\sqrt{x^2 - 1}$ via t alone.
 (c) Express dx via t and dt .
36. (a) Express x , dx and $\sqrt{1 - x^2}$ via θ and $d\theta$ for the trigonometric substitution $x = \csc \theta$, $\theta \in \left[0, \frac{\pi}{2}\right] \cup \left[\pi, \frac{3\pi}{2}\right]$.
 (b) Express x , dx and $\sqrt{1 - x^2}$ via t and dt for the Euler substitution $x = \csc(2 \arctan t)$, $t \in (-\infty, -1) \cup [0, 1)$. Express t via x .
37. Let the variables x and t be related via $\sqrt{x^2 - 1} = \frac{1}{t} - x$.
 (a) Express x via t .
 (b) Express $\sqrt{x^2 - 1}$ via t alone.
 (c) Express dx via t and dt .