

Calculus II

Area swept by one clover leaf $r = \sin(n\theta)$,
 $r = \cos(n\theta)$

Todor Milev

2019

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.
The region enclosed by the right
loop corresponds to points whose θ
polar coordinate lies in the interval
 $\leq \theta \leq ?$.

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right
loop corresponds to points whose θ
polar coordinate lies in the interval

$$-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}.$$

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right loop corresponds to points whose θ polar coordinate lies in the interval

$$-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}.$$

$$A = \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2} r^2 d\theta$$

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right loop corresponds to points whose θ polar coordinate lies in the interval $-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$.

$$\begin{aligned} A &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2} r^2 d\theta \\ &= \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \end{aligned}$$

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right loop corresponds to points whose θ polar coordinate lies in the interval $-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$.

$$\begin{aligned}
 A &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2} r^2 d\theta \\
 &= \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \cos^2(2\theta) d\theta
 \end{aligned}$$

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right loop corresponds to points whose θ polar coordinate lies in the interval $-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$.

$$\begin{aligned}
 A &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2} r^2 d\theta \\
 &= \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \frac{1}{2} (1 + \cos(4\theta)) d\theta
 \end{aligned}$$

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right loop corresponds to points whose θ polar coordinate lies in the interval $-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$.

$$\begin{aligned}
 A &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2} r^2 d\theta \\
 &= \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \frac{1}{2} (1 + \cos(4\theta)) d\theta \\
 &= \frac{1}{2} \left[\theta + \frac{1}{4} \sin(4\theta) \right]_0^{\frac{\pi}{4}}
 \end{aligned}$$

Example

Find the area enclosed by one loop of the four-leaved rose $r = \cos 2\theta$.

The region enclosed by the right loop corresponds to points whose θ polar coordinate lies in the interval $-\frac{\pi}{4} \leq \theta \leq \frac{\pi}{4}$.

$$\begin{aligned}
 A &= \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \frac{1}{2} r^2 d\theta \\
 &= \frac{1}{2} \int_{-\frac{\pi}{4}}^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \cos^2(2\theta) d\theta \\
 &= \int_0^{\frac{\pi}{4}} \frac{1}{2} (1 + \cos(4\theta)) d\theta \\
 &= \frac{1}{2} \left[\theta + \frac{1}{4} \sin(4\theta) \right]_0^{\frac{\pi}{4}} \\
 &= \frac{\pi}{8}
 \end{aligned}$$