

Calculus II

Lecture 21

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1 Complex numbers

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:

<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit.

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1.$$

The number i is called the imaginary unit.

$$\pm \sqrt{-1} = i.$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. **By definition,** $\pm \sqrt{-1} = i$.

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. **By definition,** $\sqrt{-1} = i$.

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bdi^2$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$(a + bi)(c + di) = ac + adi + bci + bd i^2 = ac + adi + bci - bd$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = \textcolor{red}{ac} + adi + bci - \textcolor{red}{bd} \\ &= (\textcolor{red}{ac} - \textcolor{red}{bd}) + i(ad + bc)\end{aligned}$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- Complex multiplication

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = ac + ad\textcolor{red}{i} + bc\textcolor{red}{i} - bd \\ &= (ac - bd) + \textcolor{red}{i}(ad + bc)\end{aligned}$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$

The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- **Complex addition/subtraction**

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$

- **Complex multiplication**

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = ac + \textcolor{red}{ad}i + \textcolor{red}{bc}i - bd \\ &= (ac - bd) + i(\textcolor{red}{ad} + \textcolor{red}{bc})\end{aligned}$$

Definition (Complex numbers)

The set of complex numbers \mathbb{C} is defined as the set

$$\{a + bi \mid a, b - \text{real numbers}\},$$

where the number i is a number for which

$$i^2 = -1 \quad .$$


The number i is called the imaginary unit. By definition, $\sqrt{-1} = i$.

- Complex addition/subtraction

$$(a + bi) \pm (c + di) = (a \pm c) + (b \pm d)i \quad .$$


- Complex multiplication

$$\begin{aligned}(a + bi)(c + di) &= ac + adi + bci + bdi^2 = ac + adi + bci - bd \\ &= (ac - bd) + i(ad + bc)\end{aligned}$$



Definition (Complex numbers)


The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.



Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.


- Real numbers are usually denoted by \mathbb{R} .



Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the **real part** of z ,




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the **imaginary part** of z .




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the **real part** of z , y is called the **imaginary part** of z .
We write $x = \text{Re}z = \text{Re}(x + yi)$, $y = \text{Im}z = \text{Im}(x + yi)$.




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z .
We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z .
We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + yi$ as the point (position vector) (x, y) in a two-dimensional space.




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + iy$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to **vector addition**.




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + yi$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to **vector addition**.




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + iy$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to **vector addition**.




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + iy$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to vector addition.
- **Multiplication by a real number c corresponds to vector scalar multiplication by c (scaling).**




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + iy$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to vector addition.
- **Multiplication by a real number c corresponds to vector scalar multiplication by c (scaling).**




Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + iy$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to vector addition.
- **Multiplication by a real number c corresponds to vector scalar multiplication by c (scaling).**



Definition (Complex numbers)

The complex numbers are the set $\{x + yi | x, y \in \mathbb{R}\}$.

- Real numbers are usually denoted by \mathbb{R} .
- Complex numbers are usually denoted by \mathbb{C} .

Consider $z = x + yi$.

- x is called the real part of z , y is called the imaginary part of z . We write $x = \operatorname{Re} z = \operatorname{Re}(x + yi)$, $y = \operatorname{Im} z = \operatorname{Im}(x + yi)$.
- Real & imaginary part of z can be used as x, y -coords. to depict z .
- In this way we view complex number $x + iy$ as the point (position vector) (x, y) in a two-dimensional space.
- The addition of complex numbers corresponds to vector addition.
- Multiplication by a real number c corresponds to vector scalar multiplication by c (scaling).
- The space the complex numbers is referred to as the **complex plane** (sometimes alternatively called the complex line).

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v =$$

.

Example (Subtraction)

$$u - v =$$

.

Example (Multiplication)

$$u \cdot v =$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = ?$$

Example (Subtraction)

$$u - v =$$

Example (Multiplication)

$$u \cdot v =$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v =$$

Example (Multiplication)

$$u \cdot v =$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = ?$$

Example (Multiplication)

$$u \cdot v =$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$u \cdot v =$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$u \cdot v = (2 + 3i) \cdot (5 - 7i)$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned}u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\&= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\&= 10 - 14i + 15i - 21i^2\end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\ &= 10 - 14i + 15i - 21i^2 \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\ &= 10 - 14i + 15i - 21i^2 \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\ &= 10 - 14i + 15i - 21i^2 \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\ &= 10 - 14i + 15i - 21i^2 \\ &= 10 + i - (-21) \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$

Example (Multiplication)

$$\begin{aligned} u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\ &= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\ &= 10 - 14i + 15i - 21i^2 \\ &= 10 + i - (-21) \end{aligned}$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Addition)

$$u + v = (2 + 3i) + (5 - 7i) = (2 + 5) + (3 - 7)i = 7 - 4i.$$

Example (Subtraction)

$$u - v = (2 + 3i) - (5 - 7i) = (2 - 5) + (3 - (-7))i = -3 + 10i.$$


Example (Multiplication)

$$\begin{aligned}u \cdot v &= (2 + 3i) \cdot (5 - 7i) \\&= 2 \cdot 5 + 2 \cdot (-7)i + 3i \cdot 5 + 3i(-7i) \\&= 10 - 14i + 15i - 21i^2 \\&= 10 + i - (-21) \\&= 31 + i\end{aligned}$$

Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)


We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate of z* .




Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.






Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.



Let $z = x + iy$ be a complex number.


Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.



Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.


Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$$z\bar{z} =$$





Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.


Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$$z\bar{z} = (x + iy)(x - iy) =$$





Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.


Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 =$$





Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.


$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2$ is real and non-negative.



Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.




In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2$ is real and non-negative.
 $z\bar{z} = 0$ implies $x^2 + y^2 = 0$, □



Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.


In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2$ is real and non-negative.
 $z\bar{z} = 0$ implies $x^2 + y^2 = 0$, which implies $x = y = 0$. □



Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.

In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2$ is real and non-negative.
 $z\bar{z} = 0$ implies $x^2 + y^2 = 0$, which implies $x = y = 0$. □


Definition

The quantity $|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}$ is called the *absolute value of z* .

Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.



In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2$ is real and non-negative.
 $z\bar{z} = 0$ implies $x^2 + y^2 = 0$, which implies $x = y = 0$. □


Definition

The quantity $|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}$ is called the *absolute value* of z .

Let $z = x + iy$ be a complex number.

Definition (Complex conjugation)

We say that $\bar{z} = \overline{(x + iy)} = x - iy$ is the *complex conjugate* of z . The transformation that maps z to \bar{z} is called *complex conjugation*.



In the complex plane, complex conj. = reflection across real axis.

Theorem

$z\bar{z}$ is a non-negative real number. $z\bar{z}$ equals 0 if and only if $z = 0$.

Proof.

$z\bar{z} = (x + iy)(x - iy) = x^2 - (iy)^2 = x^2 + y^2$ is real and non-negative.
 $z\bar{z} = 0$ implies $x^2 + y^2 = 0$, which implies $x = y = 0$. □

Definition

The quantity $|z| = \sqrt{z\bar{z}} = \sqrt{x^2 + y^2}$ is called the *absolute value* of z .

Theorem (Conjugation preserves $+$, \cdot)

1 $\overline{z \cdot w} = \overline{z} \cdot \overline{w}$

2 $\overline{z + w} = \overline{z} + \overline{w}$

Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\overline{z \cdot w} =$$

$$\bullet \quad \overline{z \cdot w} =$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\overline{z \cdot w} = \overline{(x + yi) \cdot (u + vi)}$$

$$\bullet \quad \overline{z \cdot w} =$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\overline{z \cdot w} = \overline{(x + yi) \cdot (u + vi)}$$

$$\bullet \quad \overline{z \cdot w} =$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\overline{z \cdot w} = \overline{(x + yi) \cdot (u + vi)} = (x - yi)(u - vi)$$

$$\bullet \quad \overline{z \cdot w} =$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = (\overline{x} - \overline{yi})(\overline{u} - \overline{vi}) \\ &= (\overline{x}u - \overline{y}v) - (\overline{x}v + \overline{y}u)i \\ \overline{z \cdot w} &= \end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

1 $\overline{z \cdot w} = \overline{z} \cdot \overline{w}$

2 $\overline{z + w} = \overline{z} + \overline{w}$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = (x - yi)(u - vi) \\ &= (xu - yv) - (xv + yu)i \\ \overline{z \cdot w} &= \end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = (\overline{x} - \overline{yi})(\overline{u} - \overline{vi}) \\ &= (xu - yv) - (xv + yu)i \\ \overline{z \cdot w} &= \end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = (x - yi)(u - vi) \\ &= (xu - yv) - (xv + yu)i \\ \bullet \quad \overline{z \cdot w} &= \end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i} \\ \bullet \quad \overline{z \cdot w} &= \overline{(x + yi)(u + iv)}\end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i} \\ \bullet \quad \overline{z \cdot w} &= \overline{(x + yi)(u + iv)}\end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\bullet \quad \overline{z \cdot w} = \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\overline{z \cdot w} = \overline{(x + \textcolor{red}{yi})(u + \textcolor{red}{iv})} = \overline{xu - \textcolor{red}{yv} + (xv + yu)i}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\bullet \quad \overline{z \cdot w} = \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\bullet \quad \overline{z \cdot w} = \overline{(x + \textcolor{red}{yi})(\textcolor{red}{u} + \textcolor{red}{iv})} = \overline{xu - yv + (xv + \textcolor{red}{yu})i}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i.\end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= \overline{(xu - yv) - (xv + yu)i}.\end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i} \end{aligned}$$

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i \end{aligned}$$

$$\overline{z} + \overline{w} = \overline{(x + yi)} + \overline{(u + vi)}$$

$$\overline{z} + \overline{w} =$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i} \end{aligned}$$

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i \end{aligned}$$

$$\overline{z} + \overline{w} = \overline{(x + yi)} + \overline{(u + vi)} = (x - yi) + (u - vi)$$

$$\overline{z + w} =$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i} \end{aligned}$$

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i \end{aligned}$$

$$\begin{aligned} \overline{z} + \overline{w} &= \overline{(x + yi)} + \overline{(u + vi)} = (x - yi) + (u - vi) \\ &= (x + u) - (y + v)i \end{aligned}$$

$$\overline{z + w} =$$



Theorem (Conjugation preserves $+$, \cdot)

1 $\overline{z \cdot w} = \overline{z} \cdot \overline{w}$

2 $\overline{z + w} = \overline{z} + \overline{w}$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i.\end{aligned}$$

$$\begin{aligned}\overline{z + w} &= \overline{(x + yi) + (u + vi)} = \overline{(x - yi) + (u - vi)} \\ &= \overline{(x + u) - (y + v)i}\end{aligned}$$

$$\overline{z + w} = \overline{(x + yi) + (u + iv)}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i.\end{aligned}$$

$$\begin{aligned}\overline{z + w} &= \overline{(x + yi) + (u + vi)} = \overline{(x - yi) + (u - vi)} \\ &= \overline{(x + u) - (y + v)i}\end{aligned}$$

$$\overline{z + w} = \overline{(x + yi) + (u + iv)} = \overline{(x + u) + (y + v)i}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i}\end{aligned}$$

$$\begin{aligned}\overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i.\end{aligned}$$

$$\begin{aligned}\overline{z + w} &= \overline{(x + yi) + (u + vi)} = \overline{(x - yi) + (u - vi)} \\ &= \overline{(x + u) - (y + v)i}\end{aligned}$$

$$\begin{aligned}\overline{z + w} &= \overline{(x + yi) + (u + iv)} = \overline{(x + u) + (y + v)i} \\ &= (x + u) - (y + v)i.\end{aligned}$$



Theorem (Conjugation preserves $+$, \cdot)

$$1 \quad \overline{z \cdot w} = \overline{z} \cdot \overline{w}$$

$$2 \quad \overline{z + w} = \overline{z} + \overline{w}$$

Proof.

Let $z = x + yi$, $w = u + vi$.

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi) \cdot (u + vi)} = \overline{(x - yi)(u - vi)} \\ &= \overline{(xu - yv) - (xv + yu)i} \end{aligned}$$

$$\begin{aligned} \overline{z \cdot w} &= \overline{(x + yi)(u + iv)} = \overline{xu - yv + (xv + yu)i} \\ &= (xu - yv) - (xv + yu)i \end{aligned}$$

$$\begin{aligned} \overline{z} + \overline{w} &= \overline{(x + yi)} + \overline{(u + vi)} = (x - yi) + (u - vi) \\ &= (x + u) - (y + v)i \end{aligned}$$

$$\begin{aligned} \overline{z + w} &= \overline{(x + yi) + (u + iv)} = \overline{(x + u) + (y + v)i} \\ &= (x + u) - (y + v)i \end{aligned}$$



In the preceding slide we proved the following.

Theorem (Conjugation preserves \cdot)

$$\overline{z \cdot w} = \overline{z} \cdot \overline{w}.$$

Corollary

$$|zw| = |z||w|.$$

Proof.

.



In the preceding slide we proved the following.

Theorem (Conjugation preserves \cdot)

$$\overline{z \cdot w} = \overline{z} \cdot \overline{w}.$$

Corollary

$$|zw| = |z||w|.$$

Proof.

$$|zw| = \sqrt{zw\overline{zw}}$$

.



In the preceding slide we proved the following.

Theorem (Conjugation preserves \cdot)

$$\overline{z \cdot w} = \overline{z} \cdot \overline{w}.$$

Corollary

$$|zw| = |z||w|.$$

Proof.

$$|zw| = \sqrt{zw\overline{zw}} = \sqrt{zw\overline{z}\overline{w}}.$$



In the preceding slide we proved the following.

Theorem (Conjugation preserves \cdot)

$$\overline{z \cdot w} = \overline{z} \cdot \overline{w}.$$

Corollary

$$|zw| = |z||w|.$$

Proof.

$$|zw| = \sqrt{zw\overline{zw}} = \sqrt{zw\overline{z}\overline{w}} = \sqrt{z\overline{z}}\sqrt{w\overline{w}} \quad .$$



In the preceding slide we proved the following.

Theorem (Conjugation preserves \cdot)

$$\overline{z \cdot w} = \overline{z} \cdot \overline{w}.$$

Corollary

$$|zw| = |z||w|.$$

Proof.

$$|zw| = \sqrt{zw\overline{zw}} = \sqrt{zw\overline{z}\overline{w}} = \sqrt{z\overline{z}}\sqrt{w\overline{w}} = |z||w|. \quad \square$$

In the preceding slide we proved the following.

Theorem (Conjugation preserves \cdot)

$$\overline{z \cdot w} = \overline{z} \cdot \overline{w}.$$

Corollary

$$|zw| = |z||w|.$$

Proof.

$$|zw| = \sqrt{zw\overline{zw}} = \sqrt{zw\overline{z}\overline{w}} = \sqrt{z\overline{z}}\sqrt{w\overline{w}} = |z||w|. \quad \square$$

Corollary

$$\left|\frac{z}{w}\right| = \frac{|z|}{|w|}, \quad w \neq 0.$$

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\frac{u}{v} = \frac{2 + 3i}{5 - 7i}$$

|

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i) ?}{(5 - 7i) ?}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i)(5 + 7i)}{(5 - 7i)(5 + 7i)}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i) (5 + 7i)}{(5 - 7i) (5 + 7i)} \\ &= \frac{(2 + 3i)(5 + 7i)}{\textcolor{red}{?}}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i)(5 + 7i)}{(5 - 7i)(5 + 7i)} \\ &= \frac{(2 + 3i)(5 + 7i)}{5^2 - (7i)^2}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i)(5 + 7i)}{(5 - 7i)(5 + 7i)} \\ &= \frac{(2 + 3i)(5 + 7i)}{5^2 - (7i)^2} \\ &= \frac{10 + 15i + 14i + 21i^2}{5^2 + 7^2}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i)(5 + 7i)}{(5 - 7i)(5 + 7i)} \\ &= \frac{(2 + 3i)(5 + 7i)}{5^2 - (7i)^2} \\ &= \frac{10 + 15i + 14i + 21i^2}{5^2 + 7^2} \\ &= \frac{10 - 21 + 29i}{25 + 49}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}\frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\ &= \frac{(2 + 3i)(5 + 7i)}{(5 - 7i)(5 + 7i)} \\ &= \frac{(2 + 3i)(5 + 7i)}{5^2 - (7i)^2} \\ &= \frac{10 + 15i + 14i + 21i^2}{5^2 + 7^2} \\ &= \frac{10 - 21 + 29i}{25 + 49} \\ &= \frac{-11 + 29i}{74}\end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = 2 + 3i$, $v = 5 - 7i$.

Example (Division)

$$\begin{aligned}
 \frac{u}{v} &= \frac{2 + 3i}{5 - 7i} \\
 &= \frac{(2 + 3i)(5 + 7i)}{(5 - 7i)(5 + 7i)} \\
 &= \frac{(2 + 3i)(5 + 7i)}{5^2 - (7i)^2} \\
 &= \frac{10 + 15i + 14i + 21i^2}{5^2 + 7^2} \\
 &= \frac{10 - 21 + 29i}{25 + 49} \\
 &= \frac{-11 + 29i}{74} \\
 &= -\frac{11}{74} + \frac{29}{74}i
 \end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Let $u = a + bi$, $v = c + di$, $v \neq 0$.

Example (Complex number division)

$$\begin{aligned}
 \frac{u}{v} &= \frac{a + bi}{c + di} \\
 &= \frac{(a + bi)(c - di)}{(c + di)(c - di)} \\
 &= \frac{(a + bi)(c - di)}{c^2 - (di)^2} \\
 &= \frac{ac - adi + cbi - bdi^2}{c^2 + d^2} \\
 &= \frac{ac + bd + (bc - ad)i}{c^2 + d^2} \\
 &= \frac{ac + bd}{c^2 + d^2} + \frac{(bc - ad)}{c^2 + d^2}i
 \end{aligned}$$

Multiply and divide
by complex conjugate
of denominator

Definition (Complex number division)

The quotient $\frac{u}{v}$, $v \neq 0$ is defined via the formula above.

Theorem

$$\text{Let } e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!},$$

Theorem

Let $e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, $z \in \mathbb{C}$.

Power series over \mathbb{C} are defined similarly to power series over \mathbb{R} .

Theorem

Let $e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, $z \in \mathbb{C}$. Then $e(z)e(w) = e(z + w)$.

Power series over \mathbb{C} are defined similarly to power series over \mathbb{R} .

Theorem

Let $e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, $z \in \mathbb{C}$. Then $e(z)e(w) = e(z + w)$.

Power series over \mathbb{C} are defined similarly to power series over \mathbb{R} . The following proof lies outside of scope Calc II. Details are omitted and get filled in a course of Complex Analysis. You will not be tested on it.

Proof.



Theorem

Let $e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, $z \in \mathbb{C}$. Then $e(z)e(w) = e(z + w)$.

Power series over \mathbb{C} are defined similarly to power series over \mathbb{R} . The following proof lies outside of scope Calc II. Details are omitted and get filled in a course of Complex Analysis. You will not be tested on it.

Proof.

$$\begin{aligned} e(z)e(w) &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{m=0}^{\infty} \frac{w^m}{m!} = \sum_{s=0}^{\infty} \sum_{k=0}^s \frac{z^k w^{s-k}}{k!(s-k)!} \\ &= \sum_{s=0}^{\infty} \sum_{k=0}^s \frac{z^k w^{s-k}}{s!} \frac{s!}{k!(s-k)!} = \sum_{s=0}^{\infty} \frac{(z+w)^s}{s!} = e(z+w). \end{aligned}$$



Theorem

Let $e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$, $z \in \mathbb{C}$. Then $e(z)e(w) = e(z+w)$.

Power series over \mathbb{C} are defined similarly to power series over \mathbb{R} . The following proof lies outside of scope Calc II. Details are omitted and get filled in a course of Complex Analysis. You will not be tested on it.

Proof.

$$\begin{aligned} e(z)e(w) &= \sum_{n=0}^{\infty} \frac{z^n}{n!} \sum_{m=0}^{\infty} \frac{w^m}{m!} = \sum_{s=0}^{\infty} \sum_{k=0}^s \frac{z^k w^{s-k}}{k!(s-k)!} \\ &= \sum_{s=0}^{\infty} \sum_{k=0}^s \frac{z^k w^{s-k}}{s!} \frac{s!}{k!(s-k)!} = \sum_{s=0}^{\infty} \frac{(z+w)^s}{s!} = e(z+w). \end{aligned}$$



Lemma (Newton Binomial formula)

$$(z+w)^s = \sum_{k=0}^s z^k w^{s-k} \frac{s!}{k!(s-k)!}.$$

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow z \\ p \text{ is rational}}} e^p$.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

- For real z , e^z may be defined via Definition I.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

- For real z , e^z may be defined via Definition I.
- For complex z , e^z is defined via Definition II.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

- For real z , e^z may be defined via Definition I.
- For complex z , e^z is defined via Definition II.
- Real numbers are complex numbers (with zero imaginary part).
Thus Definition II is also valid when z is a real number, and therefore Definition II is more general.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

- For real z , e^z may be defined via Definition I.
- For complex z , e^z is defined via Definition II.
- Real numbers are complex numbers (with zero imaginary part). Thus Definition II is also valid when z is a real number, and therefore Definition II is more general.
- A calculus course may be built by presenting Definition II first and proving Definition I as a theorem.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

- For real z , e^z may be defined via Definition I.
- For complex z , e^z is defined via Definition II.
- Real numbers are complex numbers (with zero imaginary part). Thus Definition II is also valid when z is a real number, and therefore Definition II is more general.
- A calculus course may be built by presenting Definition II first and proving Definition I as a theorem.
- Alternatively, a calculus course may be built by first presenting Definition I, and then expanding it to Definition II.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

- For real z , e^z may be defined via Definition I.
- For complex z , e^z is defined via Definition II.
- Real numbers are complex numbers (with zero imaginary part). Thus Definition II is also valid when z is a real number, and therefore Definition II is more general.
- A calculus course may be built by presenting Definition II first and proving Definition I as a theorem.
- Alternatively, a calculus course may be built by first presenting Definition I, and then expanding it to Definition II.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow z \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition I implies Definition II over \mathbb{R} .

Under Definition I the Maclaurin series of e^z was computed to be

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}.$$



Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition I implies Definition II over \mathbb{R} .

Under Definition I the Maclaurin series of e^z was computed to be $\sum_{n=0}^{\infty} \frac{z^n}{n!}$. Under Definition I, it can be shown that e^z equals its Maclaurin series, □

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition I implies Definition II over \mathbb{R} .

Under Definition I the Maclaurin series of e^z was computed to be $\sum_{n=0}^{\infty} \frac{z^n}{n!}$. Under Definition I, it can be shown that e^z equals its Maclaurin series, which is the defining expression for Definition II. □

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition II implies Definition I over \mathbb{R} .

We showed that $e(z + w) = e(z)e(w)$.



Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition II implies Definition I over \mathbb{R} .

We showed that $e(z + w) = e(z)e(w)$. Using that statement alone, one can show that the two definitions agree over the rational numbers.



Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow z \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition II implies Definition I over \mathbb{R} .

We showed that $e(z + w) = e(z)e(w)$. Using that statement alone, one can show that the two definitions agree over the rational numbers. Two continuous functions are equal if they are equal over the rationals, □

Definition (Real exponent, Definition I)

Let $z \in \mathbb{R}$. The real exponent e^z is defined as $\lim_{\substack{p \rightarrow x \\ p \text{ is rational}}} e^p$.

Definition (Exponent, Definition II)

Let $z \in \mathbb{C}$. The complex exponent e^z is defined by $e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}$.

Theorem

When $z \in \mathbb{R}$, Definition I and Definition II are equivalent.

Sketch of Proof. Definition II implies Definition I over \mathbb{R} .

We showed that $e(z + w) = e(z)e(w)$. Using that statement alone, one can show that the two definitions agree over the rational numbers. Two continuous functions are equal if they are equal over the rationals, and the theorem follows. \square

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots + \frac{(-1)^n x^{2n+1}}{(2n+1)!} + \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots + \frac{(-1)^n x^{2n}}{(2n)!} + \dots$$

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots + \frac{x^n}{n!} + \dots$$

Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant .

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^z = 1 + z + \frac{z^2}{2!} + \frac{z^3}{3!} + \frac{z^4}{4!} + \frac{z^5}{5!} + \dots$$

Rearrange.



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix + \frac{(ix)^2}{2!} + \frac{(ix)^3}{3!} + \frac{(ix)^4}{4!} + \frac{(ix)^5}{5!} + \dots$$

Rearrange. Plug-in $z = ix$.



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in $z = ix$. Use $i^2 = -1$.



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$i \sin x = ix - i \frac{x^3}{3!} + i \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in $z = ix$. Use $i^2 = -1$. Multiply $\sin x$ by i .



Euler's Formula

Theorem (Euler's Formula)

$$e^{ix} = \cos x + i \sin x,$$

where $e \approx 2.71828$ is Euler's/Napier's constant.

Proof.

Recall $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot (n-1) \cdot n$. Borrow from Calc II the f-las:

$$i \sin x = ix - i \frac{x^3}{3!} + i \frac{x^5}{5!} - \dots$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - i \frac{x^3}{3!} + \frac{x^4}{4!} + i \frac{x^5}{5!} - \dots$$

Rearrange. Plug-in $z = ix$. Use $i^2 = -1$. Multiply $\sin x$ by i . **Add to get**
 $e^{ix} = \cos x + i \sin x.$



Lemma


$$\left| \frac{z}{|z|} \right|$$

Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|}$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.
- Then $0, z, \frac{z}{|z|}$ lie on a ray



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.
- Then $0, z, \frac{z}{|z|}$ lie on a ray



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta$.



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$


- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta$.



Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta$.




Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$


Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$


Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$


Lemma

$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .

- Let $\rho = \ln |z|$



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .

- Let $\rho = \ln |z| = \ln \sqrt{x^2 + y^2}$.



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .

- Let $\rho = \ln |z| = \ln \sqrt{x^2 + y^2} = \frac{1}{2} \ln(x^2 + y^2)$.



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .

- Let $\rho = \ln |z| = \ln \sqrt{x^2 + y^2} = \frac{1}{2} \ln(x^2 + y^2)$.
- Then $z = |z|(\cos \theta + i \sin \theta) = e^{\rho}(\cos \theta + i \sin \theta)$



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)

Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .

- Let $\rho = \ln |z| = \ln \sqrt{x^2 + y^2} = \frac{1}{2} \ln(x^2 + y^2)$.
- Then $z = |z|(\cos \theta + i \sin \theta) = e^{\rho}(\cos \theta + i \sin \theta) = e^{\rho} e^{i\theta}$



Theorem (Euler's formula)

$$e^{i\theta} = \cos \theta + i \sin \theta.$$

Lemma


$$\left| \frac{z}{|z|} \right| = \frac{|z|}{|z|} = 1.$$

- Let $z = x + iy$ be a non-zero complex number.
- Then 0 , z , $\frac{z}{|z|}$ lie on a ray and $\frac{z}{|z|}$ lies on the unit circle.
- Let θ - angle between the real axis and the ray between 0 and z .
- Then $\frac{z}{|z|} = \cos \theta + i \sin \theta = e^{i\theta}$.

Definition (Polar form of complex numbers)


Let $z \neq 0$. $z = |z|(\cos \theta + i \sin \theta) = |z|e^{i\theta}$ is called polar form of z .

- Let $\rho = \ln |z| = \ln \sqrt{x^2 + y^2} = \frac{1}{2} \ln(x^2 + y^2)$.
- Then $z = |z|(\cos \theta + i \sin \theta) = e^{\rho}(\cos \theta + i \sin \theta) = e^{\rho} e^{i\theta} = e^{\rho + i\theta}$.



Definition (Polar form of complex numbers)

Let $z \neq 0$. Then $z = |z|(\cos \theta + i \sin \theta)$ is called polar form of z .




Definition (Polar form of complex numbers)

Let $z \neq 0$. Then $z = |z|(\cos \theta + i \sin \theta)$ is called polar form of z .

- θ is called an argument of z . We write

$$\theta = \arg z.$$




Definition (Polar form of complex numbers)

Let $z \neq 0$. Then $z = |z|(\cos \theta + i \sin \theta)$ is called polar form of z .

- θ is called an argument of z . We write

$$\theta = \arg z.$$

- If θ is an argument of z , so is $\theta + 2k\pi$ for all integers k .




Definition (Polar form of complex numbers)

Let $z \neq 0$. Then $z = |z|(\cos \theta + i \sin \theta)$ is called polar form of z .

- θ is called an argument of z . We write

$$\theta = \arg z.$$

- If θ is an argument of z , so is $\theta + 2k\pi$ for all integers k .
- If $\theta \in (-\pi, \pi]$, we say that θ is the principal argument of z .




Definition (Polar form of complex numbers)

Let $z \neq 0$. Then $z = |z|(\cos \theta + i \sin \theta)$ is called polar form of z .

- θ is called an argument of z . We write

$$\theta = \arg z.$$

- If θ is an argument of z , so is $\theta + 2k\pi$ for all integers k .
- If $\theta \in (-\pi, \pi]$, we say that θ is the principal argument of z .
- If we write $\theta = \arg z$ without clarifying the choice of the argument, it is implied that θ is the principal argument of z , $\theta \in (-\pi, \pi]$.



Definition (Polar form of complex numbers)


Let $z \neq 0$. Then $z = |z|(\cos \theta + i \sin \theta)$ is called polar form of z .

- θ is called an argument of z . We write

$$\theta = \arg z.$$

- If θ is an argument of z , so is $\theta + 2k\pi$ for all integers k .
- If $\theta \in (-\pi, \pi]$, we say that θ is the principal argument of z .
- If we write $\theta = \arg z$ without clarifying the choice of the argument, it is implied that θ is the principal argument of z , $\theta \in (-\pi, \pi]$.
- One should never write $\theta = \arg z$ without clarifying the choice of argument.

Example




Plot the number z . Write z in polar form, using the principal value of the argument of z (polar angle).

Recall that θ is the principal argument $\Rightarrow \theta \in (-\pi, \pi]$.

z	$ z $	θ	$ z (\cos \theta + i \sin \theta)$
1	1	0	$\cos 0 + i \sin 0$
i	1	$\frac{\pi}{2}$	$\cos \left(\frac{\pi}{2}\right) + i \sin \left(\frac{\pi}{2}\right)$
-1	1	π	$\cos \pi + i \sin \pi$
$-i$	1	$-\frac{\pi}{2}$	$\cos \left(-\frac{\pi}{2}\right) + i \sin \left(-\frac{\pi}{2}\right)$

Example



Plot the number z . Write z in polar form, using the principal value of the argument of z (polar angle).

Recall that θ is the principal argument $\Rightarrow \theta \in (-\pi, \pi]$.

z	$ z $	θ	$ z (\cos \theta + i \sin \theta)$
$\frac{1}{2} + \frac{\sqrt{3}}{2}i$	1	$\frac{\pi}{3}$	$\cos\left(\frac{\pi}{3}\right) + i \sin\left(\frac{\pi}{3}\right)$
$1 + i$	2	$\frac{\pi}{4}$	$2\left(\cos\left(\frac{\pi}{4}\right) + i \sin\left(\frac{\pi}{4}\right)\right)$
$1 - i$	2	$-\frac{\pi}{4}$	$2\left(\cos\left(-\frac{\pi}{4}\right) + i \sin\left(-\frac{\pi}{4}\right)\right)$
$-\sqrt{3} - i$	2	$-\frac{2\pi}{3}$	$2\left(\cos\left(\frac{2\pi}{3}\right) + i \sin\left(\frac{2\pi}{3}\right)\right)$
$\frac{3}{5} + \frac{4}{5}i$	5	$\arctan\left(\frac{4}{3}\right)$	$5\left(\cos\left(\arctan\left(\frac{4}{3}\right)\right) + i \sin\left(\arctan\left(\frac{4}{3}\right)\right)\right)$

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

- For the duration of this slide, assume Definition I of real exponent.

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via

$$e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: **Euler's f-la**).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: **Euler's f-la**).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta)$$



Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta) = \cos(\alpha + \beta) + i \sin(\alpha + \beta) \quad . \quad \square$$

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta) = \cos(\alpha + \beta) + i \sin(\alpha + \beta) \quad \square$$

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: **Euler's f-la**).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta) = \cos(\alpha + \beta) + i \sin(\alpha + \beta) = e^{i(\alpha+\beta)}. \quad \square$$

Definition (Real exponent)

Let $\rho \in \mathbb{R}$. The real exponent e^ρ is defined as $\lim_{\substack{p \rightarrow \rho \\ p \text{ is rational}}} e^p$.

Definition (Extension to \mathbb{C})

Let $\rho, \theta \in \mathbb{R}$. Define the complex exponent $e^{\rho+i\theta}$ via $e^{\rho+i\theta} = e^\rho (\cos \theta + i \sin \theta)$

- For the duration of this slide, assume Definition I of real exponent.
- Extend this def. to complex numbers (motivation: Euler's f-la).

Theorem

- (a) Let $\alpha, \beta \in \mathbb{R}$. Then $e^{i\alpha} e^{i\beta} = e^{i\alpha+i\beta} = e^{i(\alpha+\beta)}$.
- (b) Let $z, w \in \mathbb{C}$. Then $e^z e^w = e^{z+w}$.

Proof of (a).

$$e^{i\alpha} e^{i\beta} = (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) = (\cos \alpha \cos \beta - \sin \alpha \sin \beta) + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta) = \cos(\alpha + \beta) + i \sin(\alpha + \beta) = e^{i(\alpha+\beta)}. \quad \square$$

- The trig. f-las used above need separate (relatively long) proof.

Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

- For the duration of this slide, assume Definition II of exponent.

Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.

Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, \quad z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.



Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\cos(\alpha + \beta) + i \sin(\alpha + \beta) = e^{i(\alpha + \beta)}$$



Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\cos(\alpha + \beta) + i \sin(\alpha + \beta) = e^{i(\alpha+\beta)} = e^{i\alpha+i\beta}$$



Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\cos(\alpha + \beta) + i \sin(\alpha + \beta) = e^{i(\alpha+\beta)} = e^{i\alpha+i\beta} = e^{i\alpha} e^{i\beta}$$



Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\begin{aligned} \cos(\alpha + \beta) + i \sin(\alpha + \beta) &= e^{i(\alpha+\beta)} = e^{i\alpha+i\beta} = e^{i\alpha} e^{i\beta} \\ &= (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) \end{aligned}$$



Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, \quad z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\begin{aligned} \cos(\alpha + \beta) + i \sin(\alpha + \beta) &= e^{i(\alpha+\beta)} = e^{i\alpha+i\beta} = e^{i\alpha} e^{i\beta} \\ &= (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) \\ &= (\cos \alpha \cos \beta - \sin \alpha \sin \beta) \\ &\quad + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta). \end{aligned}$$



Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, \quad z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\begin{aligned} \cos(\alpha + \beta) + i \sin(\alpha + \beta) &= e^{i(\alpha+\beta)} = e^{i\alpha+i\beta} = e^{i\alpha} e^{i\beta} \\ &= (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) \\ &= (\cos \alpha \cos \beta - \sin \alpha \sin \beta) \\ &\quad + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta). \end{aligned}$$

Compare **real** and imaginary part to get the desired trig identities. □

Definition (Exponent, Def. II)

$$e^z = e(z) = \sum_{n=0}^{\infty} \frac{z^n}{n!}.$$

Definition (Polar form)

$$|z| = e^{\rho}, \quad z = e^{\rho}(\cos \theta + i \sin \theta)$$

- For the duration of this slide, assume Definition II of exponent.
- In preceding slides/lectures, by algebraic manipulations of series, we showed that $e(z)e(w) = e^z e^w = e^{z+w} = e(z+w)$.

Theorem

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + i \sin \beta \cos \alpha$$


$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta, \quad \text{where } \alpha, \beta \in \mathbb{R}$$

Proof.

$$\begin{aligned} \cos(\alpha + \beta) + i \sin(\alpha + \beta) &= e^{i(\alpha+\beta)} = e^{i\alpha+i\beta} = e^{i\alpha} e^{i\beta} \\ &= (\cos \alpha + i \sin \alpha)(\cos \beta + i \sin \beta) \\ &= (\cos \alpha \cos \beta - \sin \alpha \sin \beta) \\ &\quad + i(\cos \alpha \sin \beta + \sin \alpha \cos \beta). \end{aligned}$$


Compare real and **imaginary** part to get the desired trig identities. □

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
$$|z| = e^\rho, \quad |w| = e^\sigma.$$
- Let α, β be arguments of z, w .
$$z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$$
$$w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
$$|z| = e^\rho, \quad |w| = e^\sigma.$$
- Let α, β be arguments of z, w .
$$z = e^\rho (\cos \alpha + i \sin \alpha) = e^{\rho + i\alpha}$$
$$w = e^\sigma (\cos \beta + i \sin \beta) = e^{\sigma + i\beta}.$$

Geometric interpretation of complex multiplication



- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
$$z = e^\rho (\cos \alpha + i \sin \alpha) = e^{\rho + i\alpha}$$
$$w = e^\sigma (\cos \beta + i \sin \beta) = e^{\sigma + i\beta}.$$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha+\beta) + i \sin(\alpha+\beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication




- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Geometric interpretation of complex multiplication



- Let $z, w \neq 0$ and let
 $|z| = e^\rho, \quad |w| = e^\sigma.$
- Let α, β be arguments of z, w .
 $z = e^\rho(\cos \alpha + i \sin \alpha) = e^{\rho+i\alpha}$
 $w = e^\sigma(\cos \beta + i \sin \beta) = e^{\sigma+i\beta}.$

Theorem (Summary)

$$\begin{aligned}
 zw &= |z|(\cos \alpha + i \sin \alpha)|w|(\cos \beta + i \sin \beta) \\
 &= e^\rho(\cos \alpha + i \sin \alpha)e^\sigma(\cos \beta + i \sin \beta) = e^{\rho+i\alpha}e^{\sigma+i\beta} \\
 &= e^{\rho+\sigma+i(\alpha+\beta)} = |z||w|(\cos(\alpha + \beta) + i \sin(\alpha + \beta)).
 \end{aligned}$$

- An argument (polar angle) of zw is $\alpha + \beta$.
- \Rightarrow Multiplying complex numbers adds arguments (polar angles).
- Multiplying complex numbers multiplies absolute values.

Theorem (de Moivre's formula)

$$(\cos \alpha + i \sin \alpha)^n = \cos(n\alpha) + i \sin(n\alpha).$$

Proof.



The formula is named after the French mathematician A. de Moivre (1667-1754).

Theorem (de Moivre's formula)

$$(\cos \alpha + i \sin \alpha)^n = \cos(n\alpha) + i \sin(n\alpha).$$

Proof.

$$(\cos \alpha + i \sin \alpha)^n = (e^{i\alpha})^n$$



The formula is named after the French mathematician A. de Moivre (1667-1754).

Theorem (de Moivre's formula)

$$(\cos \alpha + i \sin \alpha)^n = \cos(n\alpha) + i \sin(n\alpha).$$

Proof.

$$(\cos \alpha + i \sin \alpha)^n = (e^{i\alpha})^n = e^{in\alpha}$$



The formula is named after the French mathematician A. de Moivre (1667-1754).

Theorem (de Moivre's formula)

$$(\cos \alpha + i \sin \alpha)^n = \cos(n\alpha) + i \sin(n\alpha).$$

Proof.

$$(\cos \alpha + i \sin \alpha)^n = (e^{i\alpha})^n = e^{in\alpha} = \cos(n\alpha) + i \sin(n\alpha). \quad \square$$

The formula is named after the French mathematician A. de Moivre (1667-1754).

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = ? (\cos (?) + i \sin (?))$.

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2(\cos(?) + i \sin(?))$.

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 (\cos (?) + i \sin (?))$.

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right)$.

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$(\sqrt{3} + i)^{2014} = \left(2e^{i\frac{\pi}{6}} \right)^{2014}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned} (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\ &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned} (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\ &= 2^{2014} e^{i \cdot 2014 \cdot \frac{\pi}{6}} \\ &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned} (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\ &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\ &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\ &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned} (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\ &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\ &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\ &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\ &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned} (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\ &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\ &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\ &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\ &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned}
 (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\
 &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\
 &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\
 &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\
 &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \\
 &= -2^{2014} \left(? + i? \right)
 \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned}
 (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\
 &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\
 &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\
 &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\
 &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \\
 &= -2^{2014} \left(-\frac{1}{2} + i? \right)
 \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned}
 (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\
 &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\
 &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\
 &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\
 &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \\
 &= -2^{2014} \left(-\frac{1}{2} + i? \right)
 \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.

Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.

$$\begin{aligned}
 (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\
 &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\
 &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\
 &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\
 &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \\
 &= -2^{2014} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right)
 \end{aligned}$$

Polar form $z = |z|(\cos \theta + i \sin \theta)$.


Example

Compute $(\sqrt{3} + i)^{2014}$ and its polar form.

Write $\sqrt{3} + i$ in polar form: $\sqrt{3} + i = 2 \left(\cos \left(\frac{\pi}{6} \right) + i \sin \left(\frac{\pi}{6} \right) \right) = 2e^{i\frac{\pi}{6}}$.


$$\begin{aligned}
 (\sqrt{3} + i)^{2014} &= \left(2e^{i\frac{\pi}{6}} \right)^{2014} \\
 &= 2^{2014} e^{i2014 \cdot \frac{\pi}{6}} \\
 &= 2^{2014} (e^{i(335 + \frac{2}{3})\pi}) \\
 &= 2^{2014} e^{i335\pi} e^{i\frac{2}{3}\pi} \\
 &= 2^{2014} (-1) \left(\cos \left(\frac{2\pi}{3} \right) + i \sin \left(\frac{2\pi}{3} \right) \right) \\
 &= -2^{2014} \left(-\frac{1}{2} + i \frac{\sqrt{3}}{2} \right) \\
 &= 2^{2013} (1 - \sqrt{3}i).
 \end{aligned}$$

Example



Find all complex solutions of the equation $z^4 = 1$.


Example



Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$.


Example



Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$


Example



Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

Example




Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$.

Example




Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$. This implies $\sin(4\theta) = 0$, $\cos(4\theta) = 1$

Example




Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$. This implies $\sin(4\theta) = 0$, $\cos(4\theta) = 1$ and so $4\theta = 2k\pi$, k -integer.

Example




Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$. This implies $\sin(4\theta) = 0$, $\cos(4\theta) = 1$ and so $4\theta = 2k\pi$, k -integer. Therefore $\theta = k\frac{\pi}{2}$.

Example




Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$. This implies $\sin(4\theta) = 0$, $\cos(4\theta) = 1$ and so $4\theta = 2k\pi$, k -integer. Therefore $\theta = k\frac{\pi}{2}$. Among those values, $\theta = -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi$ belong to $(-\pi, \pi]$.

Example




Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$. This implies $\sin(4\theta) = 0$, $\cos(4\theta) = 1$ and so $4\theta = 2k\pi$, k -integer. Therefore $\theta = k\frac{\pi}{2}$. Among those values, $\theta = -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi$ belong to $(-\pi, \pi]$. We may discard the other values of θ as do not give rise to new points.

Example



Find all complex solutions of the equation $z^4 = 1$.

Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z with $\theta \in (-\pi, \pi]$. Since $|z|^4 = |z^4| = 1$ it follows that $|z| = 1$ and so $z = \cos \theta + i \sin \theta$.

By de Moivre's equality $z^4 = \cos(4\theta) + i \sin(4\theta) = 1$. This implies $\sin(4\theta) = 0$, $\cos(4\theta) = 1$ and so $4\theta = 2k\pi$, k -integer. Therefore $\theta = k\frac{\pi}{2}$. Among those values, $\theta = -\frac{\pi}{2}, 0, \frac{\pi}{2}, \pi$ belong to $(-\pi, \pi]$. We may discard the other values of θ as do not give rise to new points.

Therefore the equation $z^4 = 1$ has 4 roots given by


$$z = \cos\left(-\frac{\pi}{2}\right) + i \sin\left(-\frac{\pi}{2}\right) = -i$$

$$z = \cos 0 + i \sin 0 = 1$$

$$z = \cos\left(\frac{\pi}{2}\right) + i \sin\left(\frac{\pi}{2}\right) = i$$

$$z = \cos \pi + i \sin \pi = -1$$

Example




Find all complex numbers z such that $z^3 = i$.

$$\begin{aligned}
 i &= (\cos(\theta) + i \sin(\theta))^3 \\
 \cos\left(\frac{\pi}{2}\right) + i \sin\left(\frac{\pi}{2}\right) &= \cos(3\theta) + i \sin(3\theta) \\
 3\theta &= \frac{\pi}{2} + 2k\pi \\
 \theta &= \frac{\pi}{6} + \frac{2k}{3}\pi
 \end{aligned}$$

de Moivre
Polar form i
 k any integer
 k any integer

Values of θ that differ by even multiple of π produce the same value for $z \Rightarrow$ restrict our attention to $\theta \in (-\pi, \pi]$, i.e. $k = 0, 1, -1 \Rightarrow \theta = \frac{\pi}{6}, \frac{5\pi}{6}, -\frac{\pi}{2}$. Our final answer is **to be continued**.

Example




Find all complex numbers z such that $z^3 = i$.
 Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z
 for which $\theta \in (-\pi, \pi]$.

	i	$=$	$(\cos(\theta) + i \sin(\theta))^3$	de Moivre Polar form i k any integer k any integer
$\cos\left(\frac{\pi}{2}\right) + i \sin\left(\frac{\pi}{2}\right)$	$=$	$\cos(3\theta) + i \sin(3\theta)$		
3θ	$=$	$\frac{\pi}{2} + 2k\pi$		
θ	$=$	$\frac{\pi}{6} + \frac{2k}{3}\pi$		

Values of θ that differ by even multiple of π produce the same value for $z \Rightarrow$ restrict our attention to $\theta \in (-\pi, \pi]$, i.e. $k = 0, 1, -1 \Rightarrow$
 $\theta = \frac{\pi}{6}, \frac{5\pi}{6}, -\frac{\pi}{2}$. Our final answer is **to be continued**.

Example




Find all complex numbers z such that $z^3 = i$.
 Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z for which $\theta \in (-\pi, \pi]$. We have that
 $1 = |i| = |z^3| = |z|^3$.

i	$= (\cos(\theta) + i \sin(\theta))^3$	de Moivre Polar form i k any integer k any integer
$\cos\left(\frac{\pi}{2}\right) + i \sin\left(\frac{\pi}{2}\right)$	$= \cos(3\theta) + i \sin(3\theta)$	
3θ	$= \frac{\pi}{2} + 2k\pi$	
θ	$= \frac{\pi}{6} + \frac{2k}{3}\pi$	

Values of θ that differ by even multiple of π produce the same value for $z \Rightarrow$ restrict our attention to $\theta \in (-\pi, \pi]$, i.e. $k = 0, 1, -1 \Rightarrow$
 $\theta = \frac{\pi}{6}, \frac{5\pi}{6}, -\frac{\pi}{2}$. Our final answer is **to be continued**.

Example



Find all complex numbers z such that $z^3 = i$.
 Let $z = |z|(\cos \theta + i \sin \theta)$ be the polar form of z for which $\theta \in (-\pi, \pi]$. We have that
 $1 = |i| = |z^3| = |z|^3$. Since $|z|$ is a positive real number, $|z|^3 = 1$ implies $|z| = 1$.

i	$= (\cos(\theta) + i \sin(\theta))^3$	de Moivre Polar form i k any integer k any integer
$\cos\left(\frac{\pi}{2}\right) + i \sin\left(\frac{\pi}{2}\right)$	$= \cos(3\theta) + i \sin(3\theta)$	
3θ	$= \frac{\pi}{2} + 2k\pi$	
θ	$= \frac{\pi}{6} + \frac{2k}{3}\pi$	

Values of θ that differ by even multiple of π produce the same value for $z \Rightarrow$ restrict our attention to $\theta \in (-\pi, \pi]$, i.e. $k = 0, 1, -1 \Rightarrow$
 $\theta = \frac{\pi}{6}, \frac{5\pi}{6}, -\frac{\pi}{2}$. Our final answer is **to be continued**.