

Calculus III

Lecture 13

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1 Double Integrals

- Riemann Sums, Double Integral Definition
- Double integral properties
- Iterated integrals

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
`https://creativecommons.org/licenses/by/3.0/us/`
and the links therein.

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

$$\text{population}(\mathcal{R}) = \sum_k \text{population}(D_k) = \sum_k \text{density}(D_k) \cdot \text{area}(D_k)$$

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

$$\text{population}(\mathcal{R}) = \sum_k \text{population}(D_k) = \sum_k \text{density}(D_k) \cdot \text{area}(D_k)$$

- To find the population density in D_k we need to count everyone (what an actual census does).

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

$$\text{population}(\mathcal{R}) = \sum_k \text{population}(D_k) = \sum_k \text{density}(D_k) \cdot \text{area}(D_k)$$

- To find the population density in D_k we need to count everyone (what an actual census does).
- Instead, we estimate the population density as follows.

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

$$\text{population}(\mathcal{R}) = \sum_k \text{population}(D_k) = \sum_k \text{density}(D_k) \cdot \text{area}(D_k)$$

- To find the population density in D_k we need to count everyone (what an actual census does).
- Instead, we estimate the population density as follows.
 - We pick a sample point P_k in each region D_k .

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

$$\text{population}(\mathcal{R}) = \sum_k \text{population}(D_k) = \sum_k \text{density}(D_k) \cdot \text{area}(D_k)$$

- To find the population density in D_k we need to count everyone (what an actual census does).
- Instead, we estimate the population density as follows.
 - We pick a sample point P_k in each region D_k .
 - We estimate the population density $\text{density}(D_k)$ by counting people in a small region around P_k ($\text{density_near}(P_k)$).

A Cheaper “Census”

Imagine we want cheap procedure to estimate population in region \mathcal{R} .

- Decompose \mathcal{R} into pairwise non-overlapping smaller regions D_k (states, counties, finer division...).

$$\text{population}(\mathcal{R}) = \sum_k \text{population}(D_k) = \sum_k \text{density}(D_k) \cdot \text{area}(D_k)$$

- To find the population density in D_k we need to count everyone (what an actual census does).
- Instead, we estimate the population density as follows.
 - We pick a sample point P_k in each region D_k .
 - We estimate the population density $\text{density}(D_k)$ by counting people in a small region around P_k ($\text{density_near}(P_k)$).
- Our population estimate becomes

$$\text{population}(\mathcal{R}) = \sum \text{pop.}(D_k) \simeq \sum \text{density_near}(P_k) \text{area}(D_k).$$

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} .

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} . Let $\{D_k\}$ be finite set of regions covering \mathcal{R}

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} . Let $\{D_k\}$ be finite set of regions covering \mathcal{R} with the following properties.

- Each D_k is a compact set.

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} . Let $\{D_k\}$ be finite set of regions covering \mathcal{R} with the following properties.

- Each D_k is a compact set.
- The boundary of each D_k is a collection of smooth curves.

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} . Let $\{D_k\}$ be finite set of regions covering \mathcal{R} with the following properties.

- Each D_k is a compact set.
- The boundary of each D_k is a collection of smooth curves.
- Two regions D_i and D_j may overlap only on their boundaries.

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} . Let $\{D_k\}$ be finite set of regions covering \mathcal{R} with the following properties.

- Each D_k is a compact set.
- The boundary of each D_k is a collection of smooth curves.
- Two regions D_i and D_j may overlap only on their boundaries.

Let P_k be a collection of sampling points with $P_k \in D_k$ for all k .

Riemann sum in two variables

Let \mathcal{R} be a compact (closed, bounded) region in the plane, and let $f: \mathcal{R} \rightarrow \mathbb{R}$ be a function on \mathcal{R} . Let $\{D_k\}$ be finite set of regions covering \mathcal{R} with the following properties.

- Each D_k is a compact set.
- The boundary of each D_k is a collection of smooth curves.
- Two regions D_i and D_j may overlap only on their boundaries.

Let P_k be a collection of sampling points with $P_k \in D_k$ for all k .

Definition (Riemann sum)

The *Riemann sum* defined by such data is $\sum_k f(P_k) \text{ area}(D_k)$.

Double Integrals

\mathcal{R} -region covered by D_k , D_k don't overlap except at boundaries.

Definition (Riemann sum)

The *Riemann sum* defined by such data is $\sum_k f(P_k) \text{ area}(D_k)$.

Definition


If the limit

$$\lim_{\max_k (\text{diam } D_k) \rightarrow 0} \sum_k f(P_k) \text{ area}(D_k)$$

exists and is finite, then its value is called the *double integral of f over \mathcal{R} (with respect to area)*, and is denoted by


$$\iint_{\mathcal{R}} f(P) dA \quad .$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.
$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.
$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.
$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.

Midpoint Rule



- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.
$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.


Midpoint Rule



- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at **midpoint**
 $P_{s,t} = ?$


Midpoint Rule



- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.

$$\iint_{\mathcal{R}} f(x, y) dx dy = \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st})$$

Midpoint Rule




- Suppose region of integration \mathcal{R} is rectangle, i.e., $\mathcal{R} = [a, b] \times [c, d]$, integration w.r.t. $dA = dx dy$.

$$\iint_{\mathcal{R}} f(P) dA = \iint_{[a,b] \times [c,d]} f(x, y) dx dy.$$
- If integral exists: approximate by fine enough Riemann sum.
- Simplest way: divide \mathcal{R} into $n \times n$ equal pieces, sides $\Delta x = \frac{b-a}{n}$, $\Delta y = \frac{d-c}{n}$.
- For $(s, t)^{th}$ rectangle D_{st} , sample at midpoint $P_{s,t} = (a + (s - \frac{1}{2}) \Delta x, c + (t - \frac{1}{2}) \Delta y)$.


$$\begin{aligned} \iint_{\mathcal{R}} f(x, y) dx dy &= \lim_{n \rightarrow \infty} \sum_{1 \leq s, t \leq n} f(P_{s,t}) \text{area}(D_{st}) \\ &\approx \sum_{1 \leq i, j \leq n} f(P_{s,t}) \Delta x \Delta y . \end{aligned}$$

Example



Use the Midpoint Rule to approximate
 $\iint_{[0,4] \times [0,2]} x^2 y dx dy$, with each side divided into
 $n = 2$ pieces.

Example




Use the Midpoint Rule to approximate

$$\iint_{[0,4] \times [0,2]} x^2 y dx dy, \text{ with each side divided into } n = 2 \text{ pieces.}$$

The small rectangles have dimensions ?

Example




Use the Midpoint Rule to approximate

$\iint_{[0,4] \times [0,2]} x^2 y dx dy$, with each side divided into
 $n = 2$ pieces.

The small rectangles have dimensions
 $\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2.

Example



Use the Midpoint Rule to approximate


$\iint_{[0,4] \times [0,2]} x^2 y dx dy$, with each side divided into $n = 2$ pieces.

The small rectangles have dimensions

$\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2. The midpoints are

?

Example



Use the Midpoint Rule to approximate


$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy$, with each side divided into $n = 2$ pieces.

The small rectangles have dimensions

$\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2. **The midpoints are**

$$P_{11} = \left(1, \frac{1}{2}\right), \quad P_{12} = \left(1, \frac{3}{2}\right), \quad P_{21} = \left(3, \frac{1}{2}\right), \quad P_{22} = \left(3, \frac{3}{2}\right).$$

Example



Use the Midpoint Rule to approximate

$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy$, with each side divided into $n = 2$ pieces.


The small rectangles have dimensions

$\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2. The midpoints are

$$P_{11} = \left(1, \frac{1}{2}\right), \quad P_{12} = \left(1, \frac{3}{2}\right), \quad P_{21} = \left(3, \frac{1}{2}\right), \quad P_{22} = \left(3, \frac{3}{2}\right).$$

$$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy \approx ?$$

Example



Use the Midpoint Rule to approximate

$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy$, with each side divided into $n = 2$ pieces.


The small rectangles have dimensions

$\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2. The midpoints are

$$P_{11} = \left(1, \frac{1}{2}\right), \quad P_{12} = \left(1, \frac{3}{2}\right), \quad P_{21} = \left(3, \frac{1}{2}\right), \quad P_{22} = \left(3, \frac{3}{2}\right).$$

$$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy \approx 2 \left(f\left(1, \frac{1}{2}\right) + f\left(3, \frac{1}{2}\right) + f\left(1, \frac{3}{2}\right) + f\left(3, \frac{3}{2}\right) \right)$$

Example



Use the Midpoint Rule to approximate

$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy$, with each side divided into $n = 2$ pieces.


The small rectangles have dimensions

$\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2. The midpoints are

$$P_{11} = \left(1, \frac{1}{2}\right), \quad P_{12} = \left(1, \frac{3}{2}\right), \quad P_{21} = \left(3, \frac{1}{2}\right), \quad P_{22} = \left(3, \frac{3}{2}\right).$$

$$\begin{aligned} \iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy &\approx 2 \left(f\left(1, \frac{1}{2}\right) + f\left(3, \frac{1}{2}\right) + f\left(1, \frac{3}{2}\right) + f\left(3, \frac{3}{2}\right) \right) \\ &= 1 \cdot \frac{1}{2} \cdot 2 + 9 \cdot \frac{1}{2} \cdot 2 + 1 \cdot \frac{3}{2} \cdot 2 + 9 \cdot \frac{3}{2} \cdot 2 \end{aligned}$$

Example



Use the Midpoint Rule to approximate

$\iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy$, with each side divided into $n = 2$ pieces.

The small rectangles have dimensions

$\frac{4-0}{2} \cdot \frac{2-0}{2} = 2 \cdot 1$ and area 2. The midpoints are

$$P_{11} = \left(1, \frac{1}{2}\right), \quad P_{12} = \left(1, \frac{3}{2}\right), \quad P_{21} = \left(3, \frac{1}{2}\right), \quad P_{22} = \left(3, \frac{3}{2}\right).$$

$$\begin{aligned} \iint_{[0,4] \times [0,2]} x^2 y \, dx \, dy &\approx 2 \left(f\left(1, \frac{1}{2}\right) + f\left(3, \frac{1}{2}\right) + f\left(1, \frac{3}{2}\right) + f\left(3, \frac{3}{2}\right) \right) \\ &= 1 \cdot \frac{1}{2} \cdot 2 + 9 \cdot \frac{1}{2} \cdot 2 + 1 \cdot \frac{3}{2} \cdot 2 + 9 \cdot \frac{3}{2} \cdot 2 \\ &= 1 + 9 + 3 + 27 = 40. \end{aligned}$$

Theoretical Examples

- The total population over a region \mathcal{R} is:

$$\text{population}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA \simeq \sum_k \text{density}(P_k) \text{area}(D_k) .$$

Theoretical Examples

- The total population over a region \mathcal{R} is:

$$\text{population}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA \simeq \sum_k \text{density}(P_k) \text{area}(D_k) .$$

- Mass is the double integral of density with respect to area:

$$\text{mass}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA .$$

Theoretical Examples

- The total population over a region \mathcal{R} is:

$$\text{population}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA \simeq \sum_k \text{density}(P_k) \text{area}(D_k) .$$

- Mass is the double integral of density with respect to area:

$$\text{mass}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA .$$

- Volume under the graph of $h: \mathcal{R} \rightarrow [0, \infty)$

$$\text{Volume} = \iint_{\mathcal{R}} h(P) \, dA .$$

Theoretical Examples

- The total population over a region \mathcal{R} is:

$$\text{population}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA \simeq \sum_k \text{density}(P_k) \text{area}(D_k) .$$

- Mass is the double integral of density with respect to area:

$$\text{mass}(\mathcal{R}) = \iint_{\mathcal{R}} \text{density}(P) \, dA .$$

- Volume under the graph of $h: \mathcal{R} \rightarrow [0, \infty)$

$$\text{Volume} = \iint_{\mathcal{R}} h(P) \, dA .$$

- Area of a region:

$$\text{Area}(\mathcal{R}) = \iint_{\mathcal{R}} 1 \, dA .$$

Double Integral Properties

$$\iint_{\mathcal{R}} f(P) \, dA = \lim_{\max_k (\text{diam } D_k) \rightarrow 0} \sum_k f(P_k) \, \text{area}(D_k)$$

Double Integral Properties

$$\iint_{\mathcal{R}} f(P) \, dA = \lim_{\max_k (\text{diam } D_k) \rightarrow 0} \sum_k f(P_k) \, \text{area}(D_k)$$

- If f is bounded and continuous, except maybe on a finite number of smooth curves, then the limit exists and is finite.

Double Integral Properties

$$\iint_{\mathcal{R}} f(P) \, dA = \lim_{\max_k (\text{diam } D_k) \rightarrow 0} \sum_k f(P_k) \, \text{area}(D_k)$$

- If f is bounded and continuous, except maybe on a finite number of smooth curves, then the limit exists and is finite.
- Linearity

$$\iint_{\mathcal{R}} [\lambda f(P) + \mu g(P)] \, dA = \lambda \iint_{\mathcal{R}} f(P) \, dA + \mu \iint_{\mathcal{R}} g(P) \, dA .$$

Double Integral Properties

$$\iint_{\mathcal{R}} f(P) \, dA = \lim_{\max_k (\text{diam } D_k) \rightarrow 0} \sum_k f(P_k) \, \text{area}(D_k)$$

- If f is bounded and continuous, except maybe on a finite number of smooth curves, then the limit exists and is finite.
- Linearity

$$\iint_{\mathcal{R}} [\lambda f(P) + \mu g(P)] \, dA = \lambda \iint_{\mathcal{R}} f(P) \, dA + \mu \iint_{\mathcal{R}} g(P) \, dA .$$

- Domain additivity: if \mathcal{R}_1 and \mathcal{R}_2 intersect only along boundaries:

$$\iint_{\mathcal{R}_1 \cup \mathcal{R}_2} f(P) \, dA = \iint_{\mathcal{R}_1} f(P) \, dA + \iint_{\mathcal{R}_2} f(P) \, dA$$

Double Integral Properties

$$\iint_{\mathcal{R}} f(P) \, dA = \lim_{\max_k (\text{diam } D_k) \rightarrow 0} \sum_k f(P_k) \, \text{area}(D_k)$$

- If f is bounded and continuous, except maybe on a finite number of smooth curves, then the limit exists and is finite.
- Linearity

$$\iint_{\mathcal{R}} [\lambda f(P) + \mu g(P)] \, dA = \lambda \iint_{\mathcal{R}} f(P) \, dA + \mu \iint_{\mathcal{R}} g(P) \, dA .$$

- Domain additivity: if \mathcal{R}_1 and \mathcal{R}_2 intersect only along boundaries:

$$\iint_{\mathcal{R}_1 \cup \mathcal{R}_2} f(P) \, dA = \iint_{\mathcal{R}_1} f(P) \, dA + \iint_{\mathcal{R}_2} f(P) \, dA$$

- Monotonicity property: If $m \leq f(P) \leq M$ for all P in \mathcal{R} , then

$$m \, \text{area}(\mathcal{R}) \leq \iint_{\mathcal{R}} f(P) \, dA \leq M \, \text{area}(\mathcal{R}) .$$

Applications

- Average value of f on \mathcal{R} .

$$\begin{aligned}\iint_{\mathcal{R}} f(P) \, dA &= \iint_{\mathcal{R}} (\text{average value of } f \text{ on } \mathcal{R}) \, dA \\ &= (\text{average value of } f \text{ on } \mathcal{R}) \iint_{\mathcal{R}} dA \\ &= (\text{average value of } f \text{ on } \mathcal{R}) \cdot \text{area}(\mathcal{R})\end{aligned}$$

$$\text{average value of } f \text{ on } \mathcal{R} = \frac{1}{\text{area}(\mathcal{R})} \iint_{\mathcal{R}} f(P) \, dA .$$

Theorem (Mean Value Theorem)

If f is continuous on \mathcal{R} , then there exists P_0 in \mathcal{R} such that

$$f(P_0) = \frac{1}{\text{area}(\mathcal{R})} \iint_{\mathcal{R}} f(Q) \, dA$$

Theorem (Analog of Fundamental Theorem of Calculus)

If f is continuous around P , then

$$\lim_{D \rightarrow \{P\}} \frac{1}{\text{area}(D)} \iint_D f(Q) \, dA = f(P)$$

Vectorial Integrals

The double integral definition extends directly to f-ns with vector output.

Definition

$$\iint_{\mathcal{R}} \mathbf{F}(P) \, dA = \lim_{\max \text{diam}(\mathcal{D}) \rightarrow 0} \sum_k \mathbf{F}(P_k) \, \text{area}(D_k)$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?

$$dq = (\text{density of charge})dA$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \varepsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\varepsilon Qc \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \varepsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\epsilon Q c \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \epsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\varepsilon Q c \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \varepsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\varepsilon Qc \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \varepsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq = \varepsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\varepsilon Qc \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \varepsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq = \varepsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA$$

$$\mathbf{F} = \iint_{\mathcal{R}} d\mathbf{F}$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\epsilon Qc \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \epsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq = \epsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA$$

$$\mathbf{F} = \iint_{\mathcal{R}} d\mathbf{F} = \iint_{\mathcal{R}} \epsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA$$

Theoretical example: Electric force on a lamina

- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\varepsilon Qc \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \varepsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq = \varepsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA$$

$$\begin{aligned} \mathbf{F} &= \iint_{\mathcal{R}} d\mathbf{F} = \iint_{\mathcal{R}} \varepsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA \\ &= \varepsilon \frac{Qq}{A(\mathcal{R})} \iint_{\mathcal{R}} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA \end{aligned}$$

Theoretical example: Electric force on a lamina


- Given:
 - a charge Q , located at the origin;
 - charge q , uniformly distributed on a planar lamina \mathcal{R} .
- What is the resulting (total) force \mathbf{F} on Q ?
- Recall that the attraction force exerted on a charge Q located at the origin by a charge c located at a point with position vector \mathbf{r} is $\epsilon Qc \frac{\mathbf{r}}{|\mathbf{r}|^3}$.

$$dq = (\text{density of charge})dA = \frac{q}{A(\mathcal{R})}dA$$

$$d\mathbf{F} = \epsilon Q \frac{\mathbf{r}}{|\mathbf{r}|^3} dq = \epsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA$$


$$\begin{aligned} \mathbf{F} &= \iint_{\mathcal{R}} d\mathbf{F} = \iint_{\mathcal{R}} \epsilon \frac{Qq}{A(\mathcal{R})} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA \\ &= \epsilon \frac{Qq}{A(\mathcal{R})} \iint_{\mathcal{R}} \frac{\mathbf{r}}{|\mathbf{r}|^3} dA \end{aligned}$$

Iterated Integrals




$$\iint_{[a,b] \times [c,d]} f(x,y) dx dy$$

Iterated Integrals




$$\iint_{[a,b] \times [c,d]} f(x,y) dx dy \approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y$$

Iterated Integrals




$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$

Iterated Integrals




$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$

Iterated Integrals




$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$

Iterated Integrals




$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$

Iterated Integrals




$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$

Iterated Integrals




$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$

Iterated Integrals



$$\begin{aligned}\iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.\end{aligned}$$


Iterated Integrals



$$\begin{aligned}
 \iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\
 &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.
 \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

Iterated Integrals




$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x, y) dx dy &\approx \sum_{1 \leq i, j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y. \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

$$\sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y \approx \sum_{j=1}^n g(y_j) \Delta y \approx \int_{y=c}^{y=d} g(y) dy$$

Iterated Integrals




$$\begin{aligned}
 \iint_{[a,b] \times [c,d]} f(x, y) dx dy &\approx \sum_{1 \leq i, j \leq n} f(x_i, y_j) \Delta x \Delta y \\
 &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y.
 \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

$$\sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y \approx \sum_{j=1}^n g(y_j) \Delta y \approx \int_{y=c}^{y=d} g(y) dy$$

Iterated Integrals




$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x, y) dx dy &\approx \sum_{1 \leq i, j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y. \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

$$\begin{aligned} \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y &\approx \sum_{j=1}^n g(y_j) \Delta y \approx \int_{y=c}^{y=d} g(y) dy \\ &= \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x, y) dx \right) dy \end{aligned}$$

Iterated Integrals




$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x, y) dx dy &\approx \sum_{1 \leq i, j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y. \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

$$\begin{aligned} \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y &\approx \sum_{j=1}^n g(y_j) \Delta y \approx \int_{y=c}^{y=d} g(y) dy \\ \iint_{[a,b] \times [c,d]} f(x, y) dx dy &= \int_{y=c}^{y=d} g(y) dy = \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x, y) dx \right) dy \end{aligned}$$

Iterated Integrals




$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y. \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

$$\begin{aligned} \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y &\approx \sum_{\substack{j=1 \\ y=c \\ y=d}}^n g(y_j) \Delta y \approx \int_{y=c}^{y=d} g(y) dy \\ \iint_{[a,b] \times [c,d]} f(x,y) dx dy &= \int_{y=c}^{y=d} g(y) dy = \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x,y) dx \right) dy \end{aligned}$$

Iterated Integrals



$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x,y) dx dy &\approx \sum_{1 \leq i,j \leq n} f(x_i, y_j) \Delta x \Delta y \\ &= \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y. \end{aligned}$$

The j^{th} summand is a Riemann sum for $g(y_j) = \int_{x=a}^{x=b} f(x, y_j) dx$.

$$\begin{aligned} \sum_{j=1}^n \left(\sum_{i=1}^n f(x_i, y_j) \Delta x \right) \Delta y &\approx \sum_{j=1}^n g(y_j) \Delta y \approx \int_{y=c}^{y=d} g(y) dy \\ \iint_{[a,b] \times [c,d]} f(x,y) dx dy &= \int_{y=c}^{y=d} g(y) dy = \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x,y) dx \right) dy \end{aligned}$$

Theorem

If f is continuous the double integral $\iint_{[a,b] \times [c,d]} f(x,y) dx dy$ exists.

Theorem

If f is continuous the double integral $\iint_{[a,b] \times [c,d]} f(x,y) dx dy$ exists.

Theorem (Fubini's Theorem)

Suppose the double integral of f exists. Then, except at a set of measure 0, the iterated integrals exist and

$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x,y) dx dy &= \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x,y) dx \right) dy \\ &= \int_{x=a}^{x=b} \left(\int_{y=c}^{y=d} f(x,y) dy \right) dx . \end{aligned}$$

Theorem

If f is continuous the double integral $\iint_{[a,b] \times [c,d]} f(x,y) dx dy$ exists.

Theorem (Fubini's Theorem)

Suppose the double integral of f exists. Then, except at a set of measure 0, the iterated integrals exist and

$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x,y) dx dy &= \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x,y) dx \right) dy \\ &= \int_{x=a}^{x=b} \left(\int_{y=c}^{y=d} f(x,y) dy \right) dx . \end{aligned}$$

This theorem allows to integrate non-continuous functions.

Theorem

If f is continuous the double integral $\iint_{[a,b] \times [c,d]} f(x,y) dx dy$ exists.


Theorem (Fubini's Theorem)

Suppose the double integral of f exists. Then, except at a **set of measure 0**, the iterated integrals exist and

$$\begin{aligned} \iint_{[a,b] \times [c,d]} f(x,y) dx dy &= \int_{y=c}^{y=d} \left(\int_{x=a}^{x=b} f(x,y) dx \right) dy \\ &= \int_{x=a}^{x=b} \left(\int_{y=c}^{y=d} f(x,y) dy \right) dx . \end{aligned}$$


This theorem allows to integrate non-continuous functions. The term “**set of measure 0**” is too technical to define here; usually studied in the subject(s) “Real Analysis/Measure Theory”.

Example



Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.


Example



Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between
?


Example



Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$.


Example



Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between ?


Example



Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy = \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) \, dy$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy = \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) \, dy$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy = \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) \, dy$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned} \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\ &= \int_{y=2}^{y=3} [\quad ? \quad] \quad ? \quad dy \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned} \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\ &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned} \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\ &= \int_{y=2}^{y=3} [x^2 + 3y^2 x] \stackrel{?}{=} dy \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned} \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\ &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) dx dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned} \iint_{[1,2] \times [2,3]} (2x + 3y^2) dx dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) dx \right) dy \\ &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned} \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\ &= \int_{y=2}^{y=3} [x^2 + 3y^2x]_{x=1}^{x=2} dy \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned}
 \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\
 &= \int_{y=2}^{y=3} [x^2 + 3y^2x]_{x=1}^{x=2} dy \\
 &= \int_{y=2}^{y=3} ((4 + 6y^2) - (1 + 3y^2)) dy
 \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned}
 \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\
 &= \int_{y=2}^{y=3} [x^2 + 3y^2x]_{x=1}^{x=2} dy \\
 &= \int_{y=2}^{y=3} ((4 + 6y^2) - (1 + 3y^2)) dy
 \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned}
 \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\
 &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \\
 &= \int_{y=2}^{y=3} ((4 + 6y^2) - (1 + 3y^2)) dy \\
 &= \int_{y=2}^{y=3} (3 + 3y^2) dy
 \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned}
 \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\
 &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \\
 &= \int_{y=2}^{y=3} ((4 + 6y^2) - (1 + 3y^2)) dy \\
 &= \int_{y=2}^{y=3} (3 + 3y^2) dy = \left[? \right]_{y=2}^{y=3}
 \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned}
 \iint_{[1,2] \times [2,3]} (2x + 3y^2) \, dx \, dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) \, dx \right) dy \\
 &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \\
 &= \int_{y=2}^{y=3} ((4 + 6y^2) - (1 + 3y^2)) dy \\
 &= \int_{y=2}^{y=3} (3 + 3y^2) dy = [3y + y^3]_{y=2}^{y=3}
 \end{aligned}$$

Example




Compute $\iint_{[1,2] \times [2,3]} (2x + 3y^2) dx dy$.

For (x, y) in $[1, 2] \times [2, 3]$, y takes values between $c = 2$ and $d = 3$. For a fixed value $y = y_0$, x takes values between $a = 1$ and $b = 2$.

$$\begin{aligned}
 \iint_{[1,2] \times [2,3]} (2x + 3y^2) dx dy &= \int_{y=2}^{y=3} \left(\int_{x=1}^{x=2} (2x + 3y^2) dx \right) dy \\
 &= \int_{y=2}^{y=3} [x^2 + 3y^2 x]_{x=1}^{x=2} dy \\
 &= \int_{y=2}^{y=3} ((4 + 6y^2) - (1 + 3y^2)) dy \\
 &= \int_{y=2}^{y=3} (3 + 3y^2) dy = [3y + y^3]_{y=2}^{y=3} \\
 &= 36 - 14 = 22 .
 \end{aligned}$$


More General Regions

What makes iterated integrals work over rectangular regions?




More General Regions

What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other.



More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



• fix x ,

More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



- fix x ,
- integrate with respect to y ,

More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



- fix x ,
- integrate with respect to y ,

More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



- fix x ,
- integrate with respect to y ,

More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



- fix x ,
- integrate with respect to y ,

More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



- fix x ,
- integrate with respect to y ,
- to obtain function that depends only on x ,

More General Regions


What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :



- fix x ,
- integrate with respect to y ,
- to obtain function that depends only on x ,
- then integrate the so obtained function in x .

More General Regions

What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :




- fix x ,
- integrate with respect to y ,
- to obtain function that depends only on x ,
- then integrate the so obtained function in x .

So far used rectangular regions;

More General Regions

What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :




- fix x ,
- integrate with respect to y ,
- to obtain function that depends only on x ,
- then integrate the so obtained function in x .

So far used rectangular regions; this also works if **slices are intervals whose endpoints depend continuously** on the location of the slice.

More General Regions

What makes iterated integrals work over rectangular regions? Slices with respect to one variable are intervals in the other. If variable is x :




- fix x ,
- integrate with respect to y ,
- to obtain function that depends only on x ,
- then integrate the so obtained function in x .

So far used rectangular regions; this also works if slices are intervals whose endpoints depend continuously on the location of the slice.


- Regions of type I: vertical slices are segments.
- Regions of type II: horizontal slices are segments.

We call such regions curvilinear trapezoids.

Strategy: Curvilinear Trapezoids (Type I)




Strategy: Curvilinear Trapezoids (Type I)




- Identify the **leftmost point(s)**, with **x-coordinate $x = a$** and the rightmost point(s), $x = b$.

Strategy: Curvilinear Trapezoids (Type I)




- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.

Strategy: Curvilinear Trapezoids (Type I)




- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.
- Draw a vertical slice at a value x between a and b .

Strategy: Curvilinear Trapezoids (Type I)



- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.
- Draw a vertical slice at a value x between a and b .
- Find the lowest point on that slice, $(x, f(x))$ and the highest point, $(x, g(x))$.

Strategy: Curvilinear Trapezoids (Type I)




- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.
- Draw a vertical slice at a value x between a and b .
- Find the lowest point on that slice, $(x, f(x))$ and the highest point, $(x, g(x))$.

The region is the region bounded by:

- the vertical lines $x = a$ and $x = b$;

Strategy: Curvilinear Trapezoids (Type I)




- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.
- Draw a vertical slice at a value x between a and b .
- Find the lowest point on that slice, $(x, f(x))$ and the highest point, $(x, g(x))$.

The region is the region bounded by:

- the vertical lines $x = a$ and $x = b$;
- the graphs of $y = f(x)$ and $y = g(x)$, with $f, g: [a, b] \rightarrow \mathbb{R}$.

Strategy: Curvilinear Trapezoids (Type I)




- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.
- Draw a vertical slice at a value x between a and b .
- Find the lowest point on that slice, $(x, f(x))$ and the highest point, $(x, g(x))$.

The region is the region bounded by:

- the vertical lines $x = a$ and $x = b$;
- the graphs of $y = f(x)$ and $y = g(x)$, with $f, g: [a, b] \rightarrow \mathbb{R}$.

$$\mathcal{R} = \{(x, y) | a \leq x \leq b, f(x) \leq y \leq g(x)\}.$$

Strategy: Curvilinear Trapezoids (Type I)



- Identify the leftmost point(s), with x -coordinate $x = a$ and the rightmost point(s), $x = b$.
- Draw a vertical slice at a value x between a and b .
- Find the lowest point on that slice, $(x, f(x))$ and the highest point, $(x, g(x))$.


The region is the region bounded by:

- the vertical lines $x = a$ and $x = b$;
- the graphs of $y = f(x)$ and $y = g(x)$, with $f, g: [a, b] \rightarrow \mathbb{R}$.


$$\mathcal{R} = \{(x, y) | a \leq x \leq b, f(x) \leq y \leq g(x)\}.$$

$$\iint_{\mathcal{R}} f(x, y) \, dx \, dy = \int_{x=a}^{x=b} \left(\int_{y=f(x)}^{y=g(x)} f(x, y) \, dy \right) dx$$

Strategy: Curvilinear Trapezoids (Type II)




Strategy: Curvilinear Trapezoids (Type II)




- Identify the **lowest point(s)**, with **y-coordinate $y = c$** and the topmost point(s), $y = d$.

Strategy: Curvilinear Trapezoids (Type II)




- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.

Strategy: Curvilinear Trapezoids (Type II)




- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.
- Draw a generic horizontal slice at some value y between c and d .

Strategy: Curvilinear Trapezoids (Type II)



- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.
- Draw a generic horizontal slice at some value y between c and d .
- Find the lowest point on that slice, $(f(y), y)$ and the topmost point, $(g(y), y)$.

Strategy: Curvilinear Trapezoids (Type II)




- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.
- Draw a generic horizontal slice at some value y between c and d .
- Find the lowest point on that slice, $(f(y), y)$ and the topmost point, $(g(y), y)$.

The region is bounded by:

- horizontal lines $y = c$ and $y = d$

Strategy: Curvilinear Trapezoids (Type II)




- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.
- Draw a generic horizontal slice at some value y between c and d .
- Find the lowest point on that slice, $(f(y), y)$ and the topmost point, $(g(y), y)$.

The region is bounded by:

- horizontal lines $y = c$ and $y = d$
- graphs of $x = f(y)$ and $x = g(y)$, with $f, g: [c, d] \rightarrow \mathbb{R}$:

Strategy: Curvilinear Trapezoids (Type II)




- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.
- Draw a generic horizontal slice at some value y between c and d .
- Find the lowest point on that slice, $(f(y), y)$ and the topmost point, $(g(y), y)$.

The region is bounded by:

- horizontal lines $y = c$ and $y = d$
- graphs of $x = f(y)$ and $x = g(y)$, with $f, g: [c, d] \rightarrow \mathbb{R}$:

$$\mathcal{R} = \{(x, y) \mid c \leq y \leq d, f(y) \leq x \leq g(y)\}.$$

Strategy: Curvilinear Trapezoids (Type II)



- Identify the lowest point(s), with y -coordinate $y = c$ and the topmost point(s), $y = d$.
- Draw a generic horizontal slice at some value y between c and d .
- Find the lowest point on that slice, $(f(y), y)$ and the topmost point, $(g(y), y)$.


The region is bounded by:

- horizontal lines $y = c$ and $y = d$
- graphs of $x = f(y)$ and $x = g(y)$, with $f, g: [c, d] \rightarrow \mathbb{R}$:

$$\mathcal{R} = \{(x, y) \mid c \leq y \leq d, f(y) \leq x \leq g(y)\}.$$

$$\iint_{\mathcal{R}} f(x, y) \, dx \, dy = \int_{y=c}^{y=d} \left(\int_{x=f(y)}^{x=g(y)} f(x, y) \, dx \right) dy$$


Strategy for Computing a Double Integral



Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.
 - Plot the curve(s) enclosing \mathcal{R} .

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.
 - Plot the curve(s) enclosing \mathcal{R} .
 - Identify the region \mathcal{R} .

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.
 - Plot the curve(s) enclosing \mathcal{R} .
 - Identify the region \mathcal{R} .
 - Chop \mathcal{R} into curvilinear trapezoids; the trapezoids are allowed to intersect only on their boundaries.

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.
 - Plot the curve(s) enclosing \mathcal{R} .
 - Identify the region \mathcal{R} .
 - Chop \mathcal{R} into curvilinear trapezoids; the trapezoids are allowed to intersect only on their boundaries.
 - By possible subdivision ensure trapezoids have smooth boundaries.

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.
 - Plot the curve(s) enclosing \mathcal{R} .
 - Identify the region \mathcal{R} .
 - Chop \mathcal{R} into curvilinear trapezoids; the trapezoids are allowed to intersect only on their boundaries.
 - By possible subdivision ensure trapezoids have smooth boundaries.
 - Integrate f over the obtained curvilinear trapezoids & collect terms.

Strategy for Computing a Double Integral




Problem

Find the integral $\iint_{\mathcal{R}} f(x, y) dx dy$ over a region \mathcal{R} enclosed by a set of smooth curves.

- We present a strategy for approaching the above problem.
- The tractability of this strategy depends on the concrete description of f and the enclosing curves.
 - Plot the curve(s) enclosing \mathcal{R} .
 - Identify the region \mathcal{R} .
 - Chop \mathcal{R} into curvilinear trapezoids; the trapezoids are allowed to intersect only on their boundaries.
 - By possible subdivision ensure trapezoids have smooth boundaries.
 - Integrate f over the obtained curvilinear trapezoids & collect terms.
- Our strategy will be augmented/combined later with variable changes (via the multivariable substitution rule).


Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$.

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$.

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$.

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$.

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. **Identify the region.**

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

The two curves intersect when ?

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute


$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

The two curves intersect when

$$\begin{aligned} x^2 &= 2x \\ x(x - 2) &= 0 \\ x &= 0 \text{ or } 2. \end{aligned}$$

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.


$$x^2 = 2x$$

The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The **intersection points** are therefore **(0, ?)** and **(2,)**.

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.


$$x^2 = 2x$$

The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The **intersection points** are therefore **(0, 0)** and **(2,)**.

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.


$$x^2 = 2x$$

The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The **intersection points are** therefore $(0, 0)$ and $(2, ?)$.

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.


$$x^2 = 2x$$

The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The **intersection points** are therefore $(0, 0)$ and $(2, 4)$.

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.


$$x^2 = 2x$$

The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The intersection points are therefore $(0, 0)$ and $(2, 4)$. We can plot the function $\frac{1}{8} (x^2 + y^2)$ as above.

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$x^2 = 2x$$


The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The intersection points are therefore $(0, 0)$ and $(2, 4)$. We can plot the function $\frac{1}{8} (x^2 + y^2)$ as above. Our integral is

$$\int_{x=?}^{x=?} \left(\int \frac{1}{8} (x^2 + y^2) \, dy \right) dx$$

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$x^2 = 2x$$


The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The intersection points are therefore $(0, 0)$ and $(2, 4)$. We can plot the function $\frac{1}{8} (x^2 + y^2)$ as above. Our integral is

$$\int_{x=0}^{x=2} \left(\int \frac{1}{8} (x^2 + y^2) \, dy \right) dx$$

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$x^2 = 2x$$


The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The intersection points are therefore $(0, 0)$ and $(2, 4)$. We can plot the function $\frac{1}{8} (x^2 + y^2)$ as above. Our integral is

$$\int_{x=0}^{x=2} \left(\int_{y=?}^{y=?} \frac{1}{8} (x^2 + y^2) \, dy \right) dx$$

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$x^2 = 2x$$


The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The intersection points are therefore $(0, 0)$ and $(2, 4)$. We can plot the function $\frac{1}{8} (x^2 + y^2)$ as above. Our integral is

$$\int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx$$

Example



Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$x^2 = 2x$$



The two curves intersect when $x(x - 2) = 0$

$$x = 0 \text{ or } 2.$$

The intersection points are therefore $(0, 0)$ and $(2, 4)$. We can plot the function $\frac{1}{8} (x^2 + y^2)$ as above. Our integral is

$$\int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx = \frac{1}{8} \int_{x=0}^{x=2} \left[\quad \right]_{y=x^2}^{y=2x} dx$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx = \frac{1}{8} \int_{x=0}^{x=2} \left[x^2 y + \frac{y^3}{3} \right]_{y=x^2}^{y=2x} dx$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\begin{aligned} \int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx &= \frac{1}{8} \int_{x=0}^{x=2} \left[x^2 y + \frac{y^3}{3} \right]_{y=x^2}^{y=2x} dx \\ &= \frac{1}{8} \int_0^2 \left(2x^3 + \frac{8}{3}x^3 - x^4 - \frac{x^6}{3} \right) dx \end{aligned}$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\begin{aligned} \int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx &= \frac{1}{8} \int_{x=0}^{x=2} \left[x^2 y + \frac{y^3}{3} \right]_{y=x^2}^{y=2x} dx \\ &= \frac{1}{8} \int_0^2 \left(2x^3 + \frac{8}{3}x^3 - x^4 - \frac{x^6}{3} \right) dx \end{aligned}$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) dx dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\begin{aligned} \int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) dy \right) dx &= \frac{1}{8} \int_{x=0}^{x=2} \left[x^2 y + \frac{y^3}{3} \right]_{y=x^2}^{y=2x} dx \\ &= \frac{1}{8} \int_0^2 \left(2x^3 + \frac{8}{3}x^3 - x^4 - \frac{x^6}{3} \right) dx \\ &= \frac{1}{8} \left[? \right]_{x=0}^{x=2} \end{aligned}$$

Example




Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\begin{aligned} \int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx &= \frac{1}{8} \int_{x=0}^{x=2} \left[x^2 y + \frac{y^3}{3} \right]_{y=x^2}^{y=2x} dx \\ &= \frac{1}{8} \int_0^2 \left(2x^3 + \frac{8}{3}x^3 - x^4 - \frac{x^6}{3} \right) dx \\ &= \frac{1}{8} \left[-\frac{1}{21}x^7 - \frac{1}{5}x^5 + \frac{7}{6}x^4 \right]_{x=0}^{x=2} \end{aligned}$$

Example





Let \mathcal{R} be the region bounded by $y = 2x$ and $y = x^2$. Compute

$$\iint_{\mathcal{R}} \frac{1}{8} (x^2 + y^2) \, dx \, dy$$

Plot $y = 2x$. Plot $y = x^2$. Identify the region.

$$\begin{aligned} \int_{x=0}^{x=2} \left(\int_{y=x^2}^{y=2x} \frac{1}{8} (x^2 + y^2) \, dy \right) dx &= \frac{1}{8} \int_{x=0}^{x=2} \left[x^2 y + \frac{y^3}{3} \right]_{y=x^2}^{y=2x} dx \\ &= \frac{1}{8} \int_0^2 \left(2x^3 + \frac{8}{3}x^3 - x^4 - \frac{x^6}{3} \right) dx \\ &= \frac{1}{8} \left[-\frac{1}{21}x^7 - \frac{1}{5}x^5 + \frac{7}{6}x^4 \right]_{x=0}^{x=2} \\ &= \frac{27}{35} \end{aligned}$$


Example




Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Example




Plot $x - 1$.




Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Example




Plot $x - 1$.




Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Example




Plot $x - 1$. Plot $y^2 = 2x + 6$.




Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Example





Plot $x - 1$. Plot $y^2 = 2x + 6$.



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. **Identify the region.**

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. **The two curves**

intersect when ?

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute



$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

intersect when

$$\begin{aligned} (x - 1)^2 &= 2x + 6 \\ x^2 - 2x + 1 &= 2x + 6 \\ x^2 - 4x - 5 &= 0 \\ x &= -1 \text{ or } 5. \end{aligned}$$

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $y = x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$



intersect when $x^2 - 2x + 1 = 2x + 6$

$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, ?)$ and $(5, ?)$.

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $y = x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$



$$\text{intersect when } x^2 - 2x + 1 = 2x + 6$$

$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$.

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$



intersect when $x^2 - 2x + 1 = 2x + 6$

$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, ?)$.

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$



$$\text{intersect when } x^2 - 2x + 1 = 2x + 6$$

$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$.

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$



intersect when $x^2 - 2x + 1 = 2x + 6$

$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$. The function can be plotted as above.

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$

intersect when $x^2 - 2x + 1 = 2x + 6$



$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$. The function can be plotted as above. The integral becomes:

$$\int_{y=?}^{y=?} \int \left(2 + \frac{1}{4}xy \right) dx dy$$

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$

intersect when $x^2 - 2x + 1 = 2x + 6$



$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$. The function can be plotted as above. The integral becomes:

$$\int_{y=-2}^{y=4} \int \left(2 + \frac{1}{4}xy \right) dx dy$$

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$

intersect when $x^2 - 2x + 1 = 2x + 6$



$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$. The function can be plotted as above. The integral becomes:

$$\int_{y=-2}^{y=4} \int_{x=?}^{x=?} \left(2 + \frac{1}{4}xy \right) dx dy$$

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$

intersect when $x^2 - 2x + 1 = 2x + 6$



$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$. The function can be plotted as above. The integral becomes:

$$\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy$$

Example



Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

Plot $x - 1$. Plot $y^2 = 2x + 6$. Identify the region. The two curves

$$(x - 1)^2 = 2x + 6$$

intersect when $x^2 - 2x + 1 = 2x + 6$



$$x^2 - 4x - 5 = 0$$

$$x = -1 \text{ or } 5.$$

The two intersection points are $(-1, -2)$ and $(5, 4)$. The function can be plotted as above. The integral becomes:

$$\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy = \int_{y=-2}^{y=4} \left[\quad \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy = \int_{y=-2}^{y=4} \left[2x + \frac{x^2 y}{8} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\begin{aligned} \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy &= \int_{y=-2}^{y=4} \left[2x + \frac{x^2 y}{8} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy \\ &= \int_{y=-2}^{y=4} ? \end{aligned}$$

dy

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\begin{aligned} \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy &= \int_{y=-2}^{y=4} \left[2x + \frac{x^2 y}{8} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy \\ &= \int_{y=-2}^{y=4} \left(-\frac{1}{32}y^5 + \frac{1}{2}y^3 - \frac{3}{4}y^2 + y + 8 \right) dy \end{aligned}$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\begin{aligned} \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy &= \int_{y=-2}^{y=4} \left[2x + \frac{x^2 y}{8} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy \\ &= \int_{y=-2}^{y=4} \left(-\frac{1}{32}y^5 + \frac{1}{2}y^3 \right. \\ &\quad \left. -\frac{3}{4}y^2 + y + 8 \right) dy \\ &= \left[? \right]_{-2}^4 \end{aligned}$$

Example





Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\begin{aligned} \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy &= \int_{y=-2}^{y=4} \left[2x + \frac{x^2 y}{8} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy \\ &= \int_{y=-2}^{y=4} \left(-\frac{1}{32}y^5 + \frac{1}{2}y^3 - \frac{3}{4}y^2 + y + 8 \right) dy \\ &= \left[-\frac{1}{192}y^6 + \frac{1}{8}y^4 - \frac{1}{4}y^3 + \frac{1}{2}y^2 + 8y \right]_{-2}^4 \end{aligned}$$

Example




Let \mathcal{R} be the region bounded by $y = x - 1$ and $y^2 = 2x + 6$. Compute

$$\iint_{\mathcal{R}} \left(2 + \frac{1}{4}xy \right) dx dy.$$

$$\begin{aligned} \int_{y=-2}^{y=4} \int_{x=\frac{y^2-6}{2}}^{x=y+1} \left(2 + \frac{1}{4}xy \right) dx dy &= \int_{y=-2}^{y=4} \left[2x + \frac{x^2 y}{8} \right]_{x=\frac{y^2-6}{2}}^{x=y+1} dy \\ &= \int_{y=-2}^{y=4} \left(-\frac{1}{32}y^5 + \frac{1}{2}y^3 \right. \\ &\quad \left. - \frac{3}{4}y^2 + y + 8 \right) dy \\ &= \left[-\frac{1}{192}y^6 + \frac{1}{8}y^4 \right. \\ &\quad \left. - \frac{1}{4}y^3 + \frac{1}{2}y^2 + 8y \right]_{-2}^4 = 45 \end{aligned}$$


Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Example




Plot $x = -1$.

Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Example




Plot $x = -1$.

Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Example




Plot $x = -1$. Plot $y = -1$.

Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Example




Plot $x = -1$. Plot $y = -1$.

Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$.

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$.

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. **Plot $x = y - y^3$.**

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. **Plot $x = y - y^3$.**

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. **Identify the region.**

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. **Identify the region.**

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. **Compute the intersection points: the four points lying on the boundary of our region have coordinates:**

?

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. **Compute the intersection points: the four points lying on the boundary of our region have coordinates: $(-1, -1)$, $(0, -1)$, $(-1, 0)$, $(0, 1)$.**

Example




Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1)$, $(0, -1)$, $(-1, 0)$, $(0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated.

Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for


$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1), (0, -1), (-1, 0), (0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated. **The integral becomes:**

$$\iint_{\mathcal{R}_1} f dA + \iint_{\mathcal{R}_2} f dA = \int \int f? + \int \int f?$$

Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for


$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1), (0, -1), (-1, 0), (0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated. **The integral becomes:**

$$\iint_{\mathcal{R}_1} f dA + \iint_{\mathcal{R}_2} f dA = \int \int f dx dy + \int \int f dx dy$$

Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for


$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1), (0, -1), (-1, 0), (0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated. **The integral becomes:**

$$\iint_{\mathcal{R}_1} f dA + \iint_{\mathcal{R}_2} f dA = \int_{y=?}^{y=?} \int f dx dy + \int_{y=?}^{y=?} \int f dx dy$$

Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for


$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1), (0, -1), (-1, 0), (0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated. **The integral becomes:**

$$\iint_{\mathcal{R}_1} f dA + \iint_{\mathcal{R}_2} f dA = \int_{y=0}^{y=1} \int_{x=-1}^{x=y-y^3} f dx dy + \int_{y=-1}^{y=0} \int_{x=-1}^{x=y-y^3} f dx dy$$

Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for


$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1), (0, -1), (-1, 0), (0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated. **The integral becomes:**

$$\iint_{\mathcal{R}_1} f dA + \iint_{\mathcal{R}_2} f dA = \int_{y=0}^{y=1} \int_{x=?}^{x=?} f dx dy + \int_{y=-1}^{y=0} \int_{x=?}^{x=?} f dx dy$$

Example



Let \mathcal{R} be region bounded by $y = (x + 1)^2$, $x = y - y^3$, the line $x = -1$ and the line $y = -1$. Set-up iterated integrals for

$$\iint_{\mathcal{R}} f dA.$$

Plot $x = -1$. Plot $y = -1$. Plot $y = (x + 1)^2$. Plot $x = y - y^3$. Identify the region. Compute the intersection points: the four points lying on the boundary of our region have coordinates:

$(-1, -1)$, $(0, -1)$, $(-1, 0)$, $(0, 1)$. Split into two curvilinear trapezoids: $\mathcal{R} = \mathcal{R}_1 \cup \mathcal{R}_2$, where $\mathcal{R}_1, \mathcal{R}_2$ are as indicated. **The integral becomes:**

$$\iint_{\mathcal{R}_1} f dA + \iint_{\mathcal{R}_2} f dA = \int_{y=0}^{y=1} \int_{x=\sqrt{y}-1}^{x=y-y^3} f dx dy + \int_{y=-1}^{y=0} \int_{x=-1}^{x=y-y^3} f dx dy$$

Example

$$\iint_{[0,\infty)\times[0,\infty)} e^{-x-y} dx dy$$

Example

$$\int_0^1 \int_{3y}^3 e^{x^2} dx dy$$