

Calculus I

Lecture 25

Volumes of Solids of Revolution

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1

Volumes

Outline

1 Volumes

2 Volumes by Cylindrical Shells

License to use and redistribute

These lecture slides and their L^AT_EX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/> and the links therein.

License to use and redistribute


These lecture slides and their L^AT_EX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,


as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/> and the links therein.


Volumes




Volumes of solids are found/defined via integration.




- How do we find the volume of a solid S ?




- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.




- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.




- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.




- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.




- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.
- Consider the part of S between two planes P_{x_1} and P_{x_2} .



- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.
- Consider the part of S between two planes P_{x_1} and P_{x_2} .
- Approximate this part of S :
- Pick a sample point x^* between x_1 and x_2 . Use a solid that has the same constant cross-sectional area $A(x^*)$ between x_1 and x_2 .




- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.
- Consider the part of S between two planes P_{x_1} and P_{x_2} .
- Approximate this part of S :
- Pick a sample point x^* between x_1 and x_2 . Use a solid that has the same constant cross-sectional area $A(x^*)$ between x_1 and x_2 .
- Let Δx be the distance from x_1 to x_2 .



Approx. volume of slab:

$$A(x^*)\Delta x$$

- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.
- Consider the part of S between two planes P_{x_1} and P_{x_2} .
- Approximate this part of S :
- Pick a sample point x^* between x_1 and x_2 . Use a solid that has the same constant cross-sectional area $A(x^*)$ between x_1 and x_2 .
- Let Δx be the distance from x_1 to x_2 .




Approx. volume of slab:

$$A(x^*)\Delta x$$

Approx. volume of S :

$$V \approx \sum_{i=1}^n A(x_i^*)\Delta x$$

- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.
- Consider the part of S between two planes P_{x_1} and P_{x_2} .
- Approximate this part of S :
- Pick a sample point x^* between x_1 and x_2 . Use a solid that has the same constant cross-sectional area $A(x^*)$ between x_1 and x_2 .
- Let Δx be the distance from x_1 to x_2 .



Approx. volume of slab:

$$A(x^*)\Delta x$$

Approx. volume of S :

$$V \approx \sum_{i=1}^n A(x_i^*)\Delta x$$

Exact volume of S :

$$V = \lim_{n \rightarrow \infty} \sum_{i=1}^n A(x_i^*)\Delta x$$

- How do we find the volume of a solid S ?
- Let P_x be the plane perpendicular to the x -axis and passing through the point x .
- The intersection of P_x with S is called a cross-section.
- Let $A(x)$ be the area of this cross-section.
- Consider the part of S between two planes P_{x_1} and P_{x_2} .
- Approximate this part of S :
- Pick a sample point x^* between x_1 and x_2 . Use a solid that has the same constant cross-sectional area $A(x^*)$ between x_1 and x_2 .
- Let Δx be the distance from x_1 to x_2 .


Definition (Volume)

Let S be a solid that lies between $x = a$ and $x = b$. If the cross-sectional area of S in the plane P_x is a continuous function $A(x)$, then the volume of S is

$$V = \lim_{n \rightarrow \infty} \sum_{i=1}^n A(x_i^*) \Delta x = \int_a^b A(x) dx$$


Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.




Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.



Example


Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the **point** $(x, 0)$ has radius \sqrt{x} .



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = ?$



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.



Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.




$$V = \int_{?}^{?} A(x)dx$$

Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.




$$V = \int_{?}^{?} A(x)dx$$

Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.




$$V = \int_{?}^{?} A(x)dx$$

Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.
- The x coords. of the solid are between 0 and 1, so its volume is




$$V = \int_0^1 A(x)dx$$

Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.
- The x coords. of the solid are between 0 and 1, so its volume is




$$V = \int_0^1 A(x)dx = \int_0^1 \pi x dx$$

Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.

- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.
- The x coords. of the solid are between 0 and 1, so its volume is




$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 \pi x \, dx \\ &= \left[\frac{\pi x^2}{2} \right]_0^1 \end{aligned}$$

Example

Find the volume of the solid obtained by rotating about the x -axis the region under the curve $y = \sqrt{x}$ from 0 to 1.


- The cross-sections of this solid are all circles.
- The circular cross-section through the point $(x, 0)$ has radius \sqrt{x} .
- The area of the cross-section is $A(x) = \pi (\sqrt{x})^2 = \pi x$.
- The volume of a single approximating section is $A(x)\Delta x$.
- The x coords. of the solid are between 0 and 1, so its volume is



$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 \pi x \, dx \\ &= \left[\pi \frac{x^2}{2} \right]_0^1 = \frac{\pi}{2} \end{aligned}$$


Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis **the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.**




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section:




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section: washer, center: $(x, 0)$.



Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section: washer, center: $(x, 0)$. Area: $A(x) = \text{Area outer disk} - \text{Area inner disk}$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area: $A(x) = \text{Area outer disk} - \text{Area inner disk}$

Inner disk radius: ? area: .

Outer disk radius: , area: .




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: .

Outer disk radius: , area: .




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: ? .

Outer disk radius: , area:




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: , area:




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: ? , area:




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:


$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area:



Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .
Outer disk radius: $x^2 + 1$, area: ?




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$V = \int A(x)dx$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$V = \int_{?}^{?} A(x) dx$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.
Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$V = \int_0^1 A(x)dx$$



Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.


Cross-section: washer, center: $(x, 0)$. Area:

$$A(x) = \text{Area outer disk} - \text{Area inner disk}$$

Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$V = \int_0^1 A(x) dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx$$



Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.


Cross-section: washer, center: $(x, 0)$. Area:

$$A(x) = \text{Area outer disk} - \text{Area inner disk}$$

Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$V = \int_0^1 A(x) dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx$$



Example (Typical Cross-Section is a Washer)


Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$V = \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx$$



Example (Typical Cross-Section is a Washer)


Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x) dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \end{aligned}$$



Example (Typical Cross-Section is a Washer)


Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x) dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[? + ? + ? \right]_0^1 \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + ? + ? \right]_0^1 \end{aligned}$$



Example (Typical Cross-Section is a Washer)


Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:

$A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + x^3 + x \right]_0^1 \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + \frac{x^3}{3} + ? \right]_0^1 \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + \frac{x^3}{3} + ? \right]_0^1 \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + \frac{x^3}{3} + x \right]_0^1 \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .

Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + \frac{x^3}{3} + x \right]_0^1 \\ &= \pi \left(\frac{1}{5} + \frac{1}{3} + 1 \right) \end{aligned}$$




Example (Typical Cross-Section is a Washer)

Find the volume of the solid obtained by rotating about the x -axis the region bounded by $y = x^2 + 1$, $y = x$, $x = 0$, and $x = 1$.

Cross-section: washer, center: $(x, 0)$. Area:
 $A(x) = \text{Area outer disk} - \text{Area inner disk}$
Inner disk radius: x , area: πx^2 .


Outer disk radius: $x^2 + 1$, area: $\pi(x^2 + 1)^2$.

$$\begin{aligned} V &= \int_0^1 A(x)dx = \int_0^1 (\pi(x^2 + 1)^2 - \pi x^2) dx \\ &= \pi \int_0^1 (x^4 + x^2 + 1) dx \\ &= \pi \left[\frac{x^5}{5} + \frac{x^3}{3} + x \right]_0^1 \\ &= \pi \left(\frac{1}{5} + \frac{1}{3} + 1 \right) = \frac{23}{15}\pi \end{aligned}$$




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.



Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at ?

radius: ?

area: $A(x) =$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at ?

radius: ?

area: $A(x) =$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at ?

radius: ?

area: $A(x) =$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: ?

area: $A(x) =$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: ?

area: $A(x) =$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) =$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = ?$




Example (Rotation About a Line Parallel to the x-axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2$




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.



Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$V = \int_{?}^{?} A(x)dx$$



Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$V = \int_0^2 A(x) dx$$



Example (Rotation About a Line Parallel to the x -axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$V = \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx$$



Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \end{aligned}$$



Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \end{aligned}$$



Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[? - ? + ? \right]_0^2 \end{aligned}$$



Example (Rotation About a Line Parallel to the x -axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[\frac{x^5}{5} - ? + ? \right]_0^2 \end{aligned}$$



Example (Rotation About a Line Parallel to the x -axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[\frac{x^5}{5} - ? + ? \right]_0^2 \end{aligned}$$



Example (Rotation About a Line Parallel to the x -axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[\frac{x^5}{5} - x^4 + ? \right]_0^2 \end{aligned}$$



Example (Rotation About a Line Parallel to the x -axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[\frac{x^5}{5} - x^4 + ? \right]_0^2 \end{aligned}$$



Example (Rotation About a Line Parallel to the x-axis)


Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.

$$\begin{aligned} V &= \int_0^2 A(x) dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[\frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \end{aligned}$$




Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.



$$\begin{aligned}V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\&= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\&= \pi \left[\frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \\&= \pi \left(\frac{2^5}{5} - 2^4 + 4 \cdot \frac{2^3}{3} \right)\end{aligned}$$


Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,

radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.



$$\begin{aligned} V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\ &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\ &= \pi \left[\frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \\ &= \pi \left(\frac{2^5}{5} - 2^4 + 4 \cdot \frac{2^3}{3} \right) \end{aligned}$$


Example (Rotation About a Line Parallel to the x -axis)

Find the volume of the solid obtained by rotating about the line $y = 1$ the region bounded by $y = -x^2 + 2x + 1$ and $y = 1$.

Cross-section: a circle centered at $(x, 1)$,


radius: $(-x^2 + 2x + 1) - 1$,

area: $A(x) = \pi ((-x^2 + 2x + 1) - 1)^2 = \pi (-x^2 + 2x)^2$.




$$\begin{aligned}
 V &= \int_0^2 A(x)dx = \int_0^2 \pi (-x^2 + 2x)^2 dx \\
 &= \pi \int_0^2 (x^4 - 4x^3 + 4x^2) dx \\
 &= \pi \left[\frac{x^5}{5} - x^4 + \frac{4x^3}{3} \right]_0^2 \\
 &= \pi \left(\frac{2^5}{5} - 2^4 + 4 \cdot \frac{2^3}{3} \right) \\
 &= \pi \left(\frac{32}{5} - 16 + \frac{32}{3} \right) = \frac{16}{15}\pi.
 \end{aligned}$$

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the **x-axis** around ...




- ... the x-axis.




- ... the y-axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...




- ... the x -axis.




- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.





- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...



- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.



Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...



- ... the x -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.



- ... the y -axis.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- ... the y -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
 - Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
 - $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
 - We understand the problem.
- ... the y -axis.
 - Approx. with washers:

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
 - Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
 - $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
 - We understand the problem.
- ... the y -axis.
 - Approx. with washers: need **inner rad. x_i** .

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...



- ... the x -axis.
 - Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
 - $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
 - We understand the problem.
- ... the y -axis.
 - Approx. with washers: need inner rad. x_i & outer rad. x_o .



Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...



- ... the x -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.



- ... the y -axis.
- Approx. with washers: need inner rad. x_i & outer rad. x_o .
- x_i and x_o : solutions to cubic:
 $-x^3 + 2x^2 - y = 0$.

Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.
- ... the y -axis.
- Approx. with washers: need inner rad. x_i & outer rad. x_o .
- x_i and x_o : solutions to cubic: $-x^3 + 2x^2 - y = 0$. Solving for x requires lots of algebra.



Find the volume obtained by rotating the region bounded by $y = 2x^2 - x^3$ and the x -axis around ...





- ... the x -axis.
- Approximate the volume using circular cylinders with radius $2x^2 - x^3$ and height Δx .
- $V = \int_0^2 \pi(2x^2 - x^3)^2 dx$.
- We understand the problem.
- ... the y -axis.
- Approx. with washers: need inner rad. x_i & outer rad. x_o .
- x_i and x_o : solutions to cubic: $-x^3 + 2x^2 - y = 0$. Solving for x requires lots of algebra.
- We show a simpler technique.



- Consider the solid obtained by rotating around the y -axis the region bounded above by $y = 2x^2 - x^3$ and below by the x -axis.




- Consider the solid obtained by rotating around the y -axis the region bounded above by $y = 2x^2 - x^3$ and below by the x -axis.
- Approximate this solid by nested cylindrical shells.




- Consider the solid obtained by rotating around the y -axis the region bounded above by $y = 2x^2 - x^3$ and below by the x -axis.
- Approximate this solid by nested cylindrical shells.
- Cylindrical shells are solids obtained by taking a cylinder and removing from its center another cylinder of equal height but smaller radius.


- Consider a cylindrical shell with:




- Consider a cylindrical shell with:
- outer radius r_2 ,




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .




- Consider a cylindrical shell with:
 - outer radius r_2 ,
 - inner radius r_1 ,
 - height h .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}}$$




- Consider a cylindrical shell with:
 - outer radius r_2 ,
 - inner radius r_1 ,
 - height h .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h$$




- Consider a cylindrical shell with:
 - outer radius r_2 ,
 - inner radius r_1 ,
 - height h .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h$$




- Consider a cylindrical shell with:
 - outer radius r_2 ,
 - inner radius r_1 ,
 - height h .
- $$V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$$




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let $\Delta r = r_2 - r_1$.







- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let $\Delta r = r_2 - r_1$.
- Let $r = \frac{r_2 + r_1}{2}$.




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h.$
- Let $\Delta r = r_2 - r_1$.
- Let $r = \frac{r_2 + r_1}{2}$.
- Then $V_{\text{shell}} = 2\pi rh\Delta r$.




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h$.
- Let $\Delta r = r_2 - r_1$.
- Let $r = \frac{r_2 + r_1}{2}$.
- Then $V_{\text{shell}} = 2\pi r h \Delta r$.




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h$.
- Let $\Delta r = r_2 - r_1$.
- Let $r = \frac{r_2 + r_1}{2}$.
- Then $V_{\text{shell}} = 2\pi r h \Delta r$.




- Consider a cylindrical shell with:
- outer radius r_2 ,
- inner radius r_1 ,
- height h .
- $V_{\text{shell}} = V_{\text{outer cyl.}} - V_{\text{inner cyl.}} = \pi r_2^2 h - \pi r_1^2 h = \pi(r_2 - r_1)(r_2 + r_1)h$.
- Let $\Delta r = r_2 - r_1$.
- Let $r = \frac{r_2 + r_1}{2}$.
- Then $V_{\text{shell}} = 2\pi rh\Delta r$.




Consider a solid obtained by rotating the region under $f(x)$ around the y axis.



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
Approximate the volume by cylindrical shells.




Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
Approximate the volume by cylindrical shells.




Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
Approximate the volume by cylindrical shells.

$$V_{\text{shell}} = 2\pi rh\Delta r$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
Approximate the volume by cylindrical shells.
Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).

$$V_{\text{shell}} = 2\pi r h \Delta r = 2\pi r f(r) \Delta r.$$




Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
Approximate the volume by cylindrical shells.
Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$




Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi r f(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$




Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
Approximate the volume by cylindrical shells.
Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to ∞ to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to ∞ to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to ∞ to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to ∞ to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be
 $h = f(r)$ (r =average outer & inner radius).


$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to ∞ to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x = \int_a^b 2\pi x f(x) dx.$$



Consider a solid obtained by rotating the region under $f(x)$ around the y axis.
 Approximate the volume by cylindrical shells.
 Select the height of an individual shell to be $h = f(r)$ (r =average outer & inner radius).

$$V_{\text{shell}} = 2\pi rh\Delta r = 2\pi rf(r)\Delta r.$$

Suppose there are n cylindrical shells and let x_1, \dots, x_n be the averages of outer and inner radii. The shell volume sum is:

$$V_{\text{approx}} = \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x.$$

Take the limit as the number of shells goes to ∞ to get

$$V = \lim_{n \rightarrow \infty} V_{\text{approx}} = \lim_{n \rightarrow \infty} \sum_{i=1}^n 2\pi x_i f(x_i) \Delta x = \int_a^b 2\pi x f(x) dx.$$


The endpoints of integration are the endpoints of the rotated region.

Definition (Volume by Cylindrical Shells)

The volume of the solid obtained by rotating around the y -axis the region under the curve $y = f(x)$ from a to b is


$$V = \int_a^b 2\pi x f(x) dx.$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.

Example




Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.


Example



Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: **outer radius x** ; height: **$?$** ; circumference: **$?$** ; infinitesimal volume: **$?$** .


Example



Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: ? ; circumference: ? ; infinitesimal volume: ? .


Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: ? ; infinitesimal volume: ? .


Example



Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: ? ; infinitesimal volume: ? .


Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: ?


Example



Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: ?


Example



Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.


Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.


Example



Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.


Example



Find the volume of the solid obtained by rotating about the y-axis the region bounded by $y = 2x^2 - x^3$ and the x-axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; **infinitesimal** volume: $2\pi x(2x^2 - x^3)dx$.

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$V = \int (2\pi x)(2x^2 - x^3)dx$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$V = \int_{?}^{?} (2\pi x)(2x^2 - x^3)dx$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$V = \int_0^2 (2\pi x)(2x^2 - x^3)dx$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$V = \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[? - ? \right]_0^2 \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - ? \right]_0^2 \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - ? \right]_0^2 \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left(\frac{2^4}{2} - \frac{2^5}{5} \right) \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left(\frac{2^4}{2} - \frac{2^5}{5} \right) \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left(\frac{2^4}{2} - \frac{2^5}{5} \right) = 2\pi \left(8 - \frac{32}{5} \right) \end{aligned}$$

Example




Find the volume of the solid obtained by rotating about the y -axis the region bounded by $y = 2x^2 - x^3$ and the x -axis.

Cylindrical shell: outer radius x ; height: $2x^2 - x^3$; circumference: $2\pi x$; infinitesimal volume: $2\pi x(2x^2 - x^3)dx$.

$$\begin{aligned} V &= \int_0^2 (2\pi x)(2x^2 - x^3)dx = 2\pi \int_0^2 (2x^3 - x^4)dx \\ &= 2\pi \left[\frac{x^4}{2} - \frac{x^5}{5} \right]_0^2 = 2\pi \left(\frac{2^4}{2} - \frac{2^5}{5} \right) = 2\pi \left(8 - \frac{32}{5} \right) = \frac{16}{5}\pi. \end{aligned}$$


Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and **the x -axis**.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about **the line $x = 1$** the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.



Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius ? ; height:

? ; circumference: ? ;

infinitesimal volume: ? .




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

? ; circumference: ? ;

infinitesimal volume: ? .



Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.


Cylindrical shell: outer radius $x - 1$; height:

?

; circumference: ?

;

infinitesimal volume: ?



Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.


Cylindrical shell: outer radius $x - 1$; height:

$$|x^3 - 3x^2 + 2x|$$

; circumference: ?

;

infinitesimal volume: ?




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: ? ;

infinitesimal volume: ? .




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: ?

infinitesimal volume: ?




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: ?




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;

infinitesimal volume: ?




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;

infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x); \text{ circumference: } 2\pi(x - 1); \\ \text{infinitesimal volume: } 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx.$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.



Example (Rotated About a Line Other Than the y -axis)


Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;

infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$V = \int_{?}^{?} 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$V = \int_{?}^{?} 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$V = \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (?) dx \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (?) dx \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-? + ? - ? + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + ? - ? + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + ? - ? + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + x^4 - ? + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + x^4 - ? + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + ? \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.

$$\begin{aligned} V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\ &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\ &= 2\pi \left[-\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \end{aligned}$$




Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.




$$\begin{aligned}
 V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\
 &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\
 &= 2\pi \left[-\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \\
 &= 2\pi \left(\left(-\frac{2^5}{5} + 2^4 - \frac{5}{3} \cdot 2^3 + 2^2 \right) \right. \\
 &\quad \left. - \left(-\frac{1^5}{5} + 1^4 - \frac{5}{3} \cdot 1^3 + 1^2 \right) \right)
 \end{aligned}$$

Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.




$$\begin{aligned}
 V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\
 &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\
 &= 2\pi \left[-\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \\
 &= 2\pi \left(\left(-\frac{2^5}{5} + 2^4 - \frac{5}{3} \cdot 2^3 + 2^2 \right) \right. \\
 &\quad \left. - \left(-\frac{1^5}{5} + 1^4 - \frac{5}{3} \cdot 1^3 + 1^2 \right) \right)
 \end{aligned}$$

Example (Rotated About a Line Other Than the y -axis)

Find the volume obtained by rotating about the line $x = 1$ the region to the right of $x = 1$ bounded by $y = x^3 - 3x^2 + 2x$ and the x -axis.

Cylindrical shell: outer radius $x - 1$; height:

$|x^3 - 3x^2 + 2x| = -(x^3 - 3x^2 + 2x)$; circumference: $2\pi(x - 1)$;
infinitesimal volume: $2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx$.



$$\begin{aligned}
 V &= \int_1^2 2\pi(x - 1)(-x^3 + 3x^2 - 2x)dx \\
 &= 2\pi \int_1^2 (-x^4 + 4x^3 - 5x^2 + 2x)dx \\
 &= 2\pi \left[-\frac{x^5}{5} + x^4 - \frac{5x^3}{3} + x^2 \right]_1^2 \\
 &= 2\pi \left(\left(-\frac{2^5}{5} + 2^4 - \frac{5}{3} \cdot 2^3 + 2^2 \right) \right. \\
 &\quad \left. - \left(-\frac{1^5}{5} + 1^4 - \frac{5}{3} \cdot 1^3 + 1^2 \right) \right) = \frac{4}{15}\pi.
 \end{aligned}$$

	Rotate about ...	
	... a horizontal line	... a vertical line
y is a function of x	Cross-sections $\int \cdot dx$	Cylindrical shells $\int \cdot dx$
x is a function of y	Cylindrical shells $\int \cdot dy$	Cross-sections $\int \cdot dy$

- $\int \cdot dx$ means integrate with respect to x .
- $\int \cdot dy$ means integrate with respect to y .
- Some equations express y as a function of x and x as a function of y . In such cases, you may use either method.