

Calculus I

Lecture 24

Area between Graphs

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

- 1 Integration and symmetry
- 2 More About Areas

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Symmetry



Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on $[-a, a]$.

① If f is even (that is, $f(-x) = f(x)$), then $\int_{-a}^a f(x) dx = 2 \int_0^a f(x) dx$.

② If f is odd (that is, $f(-x) = -f(x)$), then

$$\int_{-a}^a f(x) dx = \int_0^a f(x) dx + \int_{-a}^0 f(x) dx = 0.$$




Example

Since $f(x) = x^6 + 1$ satisfies $f(-x) = f(x)$, it is even, and so

$$\begin{aligned}\int_{-2}^2 (x^6 + 1) \, dx &= 2 \int_0^2 (x^6 + 1) \, dx \\ &= 2 \left[\frac{1}{7} x^7 + x \right]_0^2 \\ &= 2 \left(\frac{128}{7} + 2 \right) \\ &= \frac{284}{7}.\end{aligned}$$

Example




Since $f(x) = \frac{\tan x - x}{1 - 2x^2 + 2x^4}$ satisfies $f(-x) = -f(x)$, it is odd, and so


$$\int_{-1}^1 \frac{\tan x - x}{1 - 2x^2 + 2x^4} dx = 0.$$

More About Areas


Suppose two curves, $y = f(x)$ and $y = g(x)$, are given. How do we find the area bounded by those curves between the endpoints $x = a$ and $x = b$?



The Area Under a Curve




$$\text{rectangle area} = f(x) \cdot \Delta x$$




$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b f(x) dx$$

The Area Between Two Curves



$$\text{rectangle area} = (f(x) - g(x)) \cdot \Delta x$$



$$\# \text{ rectangles} = n \rightarrow \infty$$

$$A = \int_a^b [f(x) - g(x)] dx$$

Definition (The Area Between Two Curves)


The area between two curves $y = f(x)$ and $y = g(x)$ bounded by the endpoints $x = a$ and $x = b$ is

$$\int_a^b |f(x) - g(x)| dx.$$

Note that we use the absolute value, because in general we don't know which curve is above the other.

Example


Find the area of the region bounded above by $y = x^2 + 1$, bounded below by $y = x$, and bounded on its sides by $x = 0$ and $x = 1$.



- 1 Graph the functions.
- 2 Identify the region.
- 3 Integrate.

$$\begin{aligned} A &= \int_0^1 |(x^2 + 1) - x| dx \\ &= \int_0^1 (x^2 - x + 1) dx \\ &= \left[\frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1 \\ &= \frac{1}{3} - \frac{1}{2} + 1 = \frac{5}{6}. \end{aligned}$$

Example



Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$.

$$x^2 = 2x - x^2$$


$$0 = 2x - 2x^2 = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$\begin{aligned} A &= \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx \\ &= 2 \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 2 \left(\frac{1}{2} - \frac{1}{3} \right) = \frac{1}{3}. \end{aligned}$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.

Example



Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, $x = 0$ and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

$$A = A_1 + A_2$$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx$$

$$+ \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

$$= [\sin x + \cos x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{\pi/2}$$

$$= 2\sqrt{2} - 2.$$

- 1 Find the point of intersection.
- 2 Graph the functions.
- 3 Identify the region.
- 4 Integrate.