Precalculus Lecture 12 Equations with Logarithms and Exponents

Todor Miley

https://github.com/tmilev/freecalc

2020

Equations involving logarithms

- Equations involving logarithms
- Equations involving exponents

- Equations involving logarithms
- Equations involving exponents
- 3 Inverse function problems and exponents

- Equations involving logarithms
- Equations involving exponents
- 3 Inverse function problems and exponents
- Basic exponential inequalities

License to use and redistribute

These lecture slides and their LATEX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share to copy, distribute and transmit the work,
- to Remix to adapt, change, etc., the work,
- to make commercial use of the work.

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides: https://github.com/tmilev/freecalc
- Should the link be outdated/moved, search for "freecalc project".
- Creative Commons license CC BY 3.0:
 https://creativecommons.org/licenses/by/3.0/us/
 and the links therein.

$$\log_3(2x^2+1) = 2$$

Solve the equation.

$$\log_3(2x^2+1) = 2$$

$$3^{\log_3(2x^2+1)} = 3^2$$

Solve the equation.

$$\log_3(2x^2 + 1) = 2
3^{\log_3(2x^2+1)} = 3^2
2x^2 + 1 = 9$$

Solve the equation.

$$\begin{array}{rcl} \log_3(2x^2+1) & = & 2 \\ 3^{\log_3(2x^2+1)} & = & 3^2 \\ 2x^2+1 & = & 9 \\ 2x^2 & = & 8 \end{array}$$

Solve the equation.

the equation.

$$\log_{3}(2x^{2} + 1) = 2$$

$$3^{\log_{3}(2x^{2} + 1)} = 3^{2}$$

$$2x^{2} + 1 = 9$$

$$2x^{2} = 8$$

$$x^{2} = \frac{8}{2} = 4$$

Solve the equation.

the equation.
$$\log_{3}(2x^{2} + 1) = 2$$

$$3^{\log_{3}(2x^{2} + 1)} = 3^{2}$$

$$2x^{2} + 1 = 9$$

$$2x^{2} = 8$$

$$x^{2} = \frac{8}{2} = 4$$

$$x = \pm \sqrt{4} = \pm 2$$

$$\log_3(2x^2+1) = 2$$
 | Exponentiate base 3 $3^{\log_3(2x^2+1)} = 3^2$ $2x^2+1 = 9$ $2x^2 = 8$ $x^2 = \frac{8}{2} = 4$ $x = \pm\sqrt{4} = \pm 2$ $x = 2$ or $x = -2$ | final answer

The logarithmic property $\log_a(xy) = \log_a x + \log_a y$ holds only for positive x, y. Failure to check the positivity of x, y can result in extraneous (fake) solutions to logarithmic equations.

Example

$$\log_2(x+2) + \log_2(x-1) = 2$$

The logarithmic property $\log_a(xy) = \log_a x + \log_a y$ holds only for positive x, y. Failure to check the positivity of x, y can result in extraneous (fake) solutions to logarithmic equations.

Example

$$\log_2(x+2) + \log_2(x-1) = 2$$

$$\log_2((x+2)(x-1)) = 2$$

$$(x+2)(x-1) = 2^2$$

$$x^2 + x - 2 = 4$$

$$x^2 + x - 6 = 0$$

$$(x-2)(x+3) = 0$$

$$x = 2 \quad \text{or} \quad x = 3$$

$$x = -3 \text{ not a solution (outside of domain)}$$

$$16^{4t} = 8^{t-2}$$

Solve for t.

$$16^{4t} = 8^{t-2}$$

 $16^{4t} = 8^{t-2}$ Find a common base: $(?)^{4t} = (?)^{t-2}$

$$\begin{array}{rcl} & 16^{4t} & = & 8^{t-2} \\ \text{Find a common base:} & \left(2^4\right)^{4t} & = & \left(2^3\right)^{t-2} \end{array}$$

Find a common base:
$$(2^4)^{4t} = 8^{t-2}$$

 $2^{16t} = 2^{3t-6}$

Find a common base:
$$(2^4)^{4t} = 8^{t-2}$$

 $2^{16t} = 2^{3t-6}$

Find a common base:
$$(2^4)^{4t} = 8^{t-2}$$

 $2^{16t} = 2^{3t-6}$
 $16t = 3t-6$
 $13t = -6$

$$2^{1-5x} = 12$$

Solve the equation.

$$2^{1-5x} = 12 \log_2(2^{1-5x}) = \log_2 12$$

apply log2

Solve the equation.

$$2^{1-5x} = 12$$

 $\log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12$

apply log₂

$$2^{1-5x} = 12$$
 apply $\log_2 \log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = ?$

$$2^{1-5x} = 12$$
 | apply $\log_2 \log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = \log_2(4\cdot 3)$

$$2^{1-5x} = 12$$
 | apply \log_2
 $\log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$
 $1-5x = \log_2 4 + \log_2 3$

$$2^{1-5x} = 12$$
 | apply $\log_2 \log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = \log_2(4\cdot 3)$
 $1-5x = \log_2 4 + \log_2 3$
 $1-5x = ? + \log_2 3$

$$2^{1-5x} = 12$$
 | apply $\log_2 \log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$
 $1-5x = \log_2 4 + \log_2 3$
 $1-5x = 2 + \log_2 3$

$$2^{1-5x} = 12$$
 | apply \log_2
 $\log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$
 $1-5x = \log_2 4 + \log_2 3$
 $1-5x = 2 + \log_2 3$
 $5x = 1 - (2 + \log_2 3)$

$$2^{1-5x} = 12$$
 | apply \log_2
 $\log_2(2^{1-5x}) = \log_2 12$
 $1-5x = \log_2 12 = \log_2(4 \cdot 3)$
 $1-5x = \log_2 4 + \log_2 3$
 $1-5x = 2 + \log_2 3$
 $5x = 1 - (2 + \log_2 3)$
 $x = \frac{-1}{2}$

$$\begin{array}{rclcrcl} 2^{1-5x} & = & 12 & & | \ \mathsf{apply} \ \mathsf{log}_2 \\ \mathsf{log}_2(2^{1-5x}) & = & \mathsf{log}_2 \ \mathsf{12} \\ & 1-5x & = & \mathsf{log}_2 \ \mathsf{12} = \mathsf{log}_2(4 \cdot 3) \\ & 1-5x & = & \mathsf{log}_2 \ \mathsf{4} + \mathsf{log}_2 \ \mathsf{3} \\ & 1-5x & = & 2+\mathsf{log}_2 \ \mathsf{3} \\ & 5x & = & 1-(2+\mathsf{log}_2 \ \mathsf{3}) \\ & x & = & \frac{-1-\mathsf{log}_2 \ \mathsf{3}}{2} \end{array}$$

$$\begin{array}{rclcrcl} 2^{1-5x} & = & 12 & & & | \ \mathsf{apply} \ \mathsf{log}_2 \\ \mathsf{log}_2(2^{1-5x}) & = & \mathsf{log}_2 \ \mathsf{12} \\ & 1-5x & = & \mathsf{log}_2 \ \mathsf{12} = \mathsf{log}_2(4 \cdot 3) \\ & 1-5x & = & \mathsf{log}_2 \ \mathsf{4} + \mathsf{log}_2 \ \mathsf{3} \\ & 1-5x & = & 2+\mathsf{log}_2 \ \mathsf{3} \\ & 5x & = & 1-(2+\mathsf{log}_2 \ \mathsf{3}) \\ & x & = & \frac{-1-\mathsf{log}_2 \ \mathsf{3}}{5} \end{array}$$

$$e^{x-3} = 2e^{2x-1}$$

Solve the equation.

$$e^{x-3} = 2e^{2x-1}$$
 $\frac{e^{x-3}}{e^{2x-1}} = 2$

Divide by e^{2x-1}

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

Divide by
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

Divide by
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

$$e^{-x-2} = 2$$

Divide by
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$

$$\frac{e^{x-3}}{e^{2x-1}} = 2$$

$$e^{x-3-(2x-1)} = 2$$

$$e^{-x-2} = 2$$

Divide by
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$
 Divide $\frac{e^{x-3}}{e^{2x-1}} = 2$ $e^{x-3-(2x-1)} = 2$ $e^{-x-2} = 2$ Apply $-x-2 = \ln 2$

Divide by
$$e^{2x-1}$$

$$e^{x-3} = 2e^{2x-1}$$
 $\frac{e^{x-3}}{e^{2x-1}} = 2$
 $e^{x-3-(2x-1)} = 2$
 $e^{-x-2} = 2$
 $-x-2 = \ln 2$

Divide by
$$e^{2x-1}$$
Apply In

Solve the equation.

$$e^{x-3} = 2e^{2x-1}$$
 Divide
 $\frac{e^{x-3}}{e^{2x-1}} = 2$
 $e^{x-3-(2x-1)} = 2$
 $e^{-x-2} = 2$ Apply
 $-x-2 = \ln 2$
 $-x = \ln 2 + 2$

Divide by e^{2x-1}

Apply In

$$e^{x-3} = 2e^{2x-1}$$
 Divide by e^{2x-1}
 $\frac{e^{x-3}}{e^{2x-1}} = 2$
 $e^{x-3-(2x-1)} = 2$
 $e^{-x-2} = 2$ Apply In
 $-x-2 = \ln 2$
 $-x = \ln 2 + 2$
 $x = -(\ln 2 + 2)$

$$e^{x-3}=2e^{2x-1}$$
 Divide by e^{2x-1}
 $\frac{e^{x-3}}{e^{2x-1}}=2$
 $e^{x-3-(2x-1)}=2$
 $e^{-x-2}=2$ Apply In
 $-x-2=\ln 2$
 $-x=\ln 2+2$
 $x=-(\ln 2+2)$
 $x=-\ln 2-2$ Final answer

$$e^{x-3}=2e^{2x-1}$$
 Divide by e^{2x-1} $\frac{e^{x-3}}{e^{2x-1}}=2$ $e^{x-3-(2x-1)}=2$ Apply In $-x-2=\ln 2$ $-x=\ln 2+2$ $x=-(\ln 2+2)$ $x=-\ln 2-2$ Final answer $x\approx -2.693$ Calculator

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

Solve.

$$3^{2x+5} - 5 \cdot 2^{-x+1}$$

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$
$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+5}$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} =$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} =$$

$$(\log_2 3)(2x+5)+x-1 = \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 4$$

$$(\log_2 3)(2x + 5) + x - 1 = \log_2 5$$

$$x(+)+ = \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Apply \log_2

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 3$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 +) + = \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = \xi$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 4$$

$$(\log_2 3)(2x+5)+x-1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 3$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 3$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1}{1}$$

Common base

$$a = b^{\log_b a}$$

Apply \log_2

2020

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

$$x = \frac{\log_2 5 + 1 - 5\log_2 3}{2}$$

Common base

$$a = b^{\log_b a}$$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} = 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

Common base

$$a = b^{\log_b a}$$

Apply log₂

 $x = \frac{\log_2 5 + 1 - 5\log_2 3}{2\log_2 3 + 1}$

Solve.

$$3^{2x+5} = 5 \cdot 2^{-x+1}$$

$$\left(2^{\log_2 3}\right)^{2x+5} \ = \ 5 \cdot 2^{-x+1}$$

$$2^{(\log_2 3)(2x+5)} = 5 \cdot 2^{-x+1}$$

$$\frac{2^{(\log_2 3)(2x+5)}}{2^{-x+1}} = 5$$

$$2^{(\log_2 3)(2x+5)-(-x+1)} = 5$$

$$(\log_2 3)(2x+5) + x - 1 = \log_2 5$$

$$x(2\log_2 3 + 1) + 5\log_2 3 = \log_2 5 + 1$$

 $x = \frac{\log_2 5 + 1 - 5\log_2 3}{2\log_2 3 + 1}$

$$x \approx -1.1038$$

Common base

$$a = b^{\log_b a}$$

Apply log₂

Calculator

$$e^{5-3x} = 10$$

$$e^{5-3x} = 10$$
 apply $\ln (e^{5-3x}) = \ln 10$

$$e^{5-3x}=10$$
 apply In $\ln(e^{5-3x})=\ln 10$ $5-3x=\ln 10$

$$e^{5-3x} = 10$$
 apply In $\ln(e^{5-3x}) = \ln 10$
 $5-3x = \ln 10$
 $3x = 5 - \ln 10$

$$e^{5-3x} = 10$$
 apply In $\ln(e^{5-3x}) = \ln 10$ $5-3x = \ln 10$ $3x = 5 - \ln 10$ $x = \frac{5 - \ln 10}{3}$

$$e^{5-3x}=10$$
 apply In $\ln(e^{5-3x})=\ln 10$ $5-3x=\ln 10$ $3x=5-\ln 10$ $x=\frac{5-\ln 10}{3}$ Calculator: $x\approx 0.8991$.

Example (Solving an exponential word problem)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for *t*:

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for t: c(t)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for t: c(t) =

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for t: c(t) = r(t)

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for
$$t$$
: $c(t) = r(t)$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for
$$t$$
: $c(t) = r(t)$
 $48 \cdot 2^t =$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for
$$t$$
: $c(t) = r(t)$

$$48 \cdot 2^{t} =$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for
$$t$$
: $c(t) = r(t)$
 $48 \cdot 2^t = 6 \cdot 4^t$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for
$$t$$
: $c(t) = r(t)$

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$
 $48 \cdot 2^t = 6 \cdot 4^t$
 $8 \cdot 2^t = 4^t$

Find a common base: $2^{?} \cdot 2^{t} = 2^{?}$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$
 $48 \cdot 2^t = 6 \cdot 4^t$
 $8 \cdot 2^t = 4^t$

Find a common base: $2^3 \cdot 2^t = 2^?$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$
 $48 \cdot 2^t = 6 \cdot 4^t$
 $8 \cdot 2^t = 4^t$

Find a common base: $2^3 \cdot 2^t = 2^?$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$
 $48 \cdot 2^t = 6 \cdot 4^t$
 $8 \cdot 2^t = 4^t$

Find a common base: $2^3 \cdot 2^t = 2^{2t}$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Solve for
$$t$$
: $c(t) = r(t)$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

Find a common base:
$$2^3 \cdot 2^t = 2^{2t}$$

$$2^{t+3} = 2^{2t}$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$

$$48 \cdot 2^t = 6 \cdot 4^t$$

$$8 \cdot 2^t = 4^t$$

Find a common base: $2^3 \cdot 2^t = 2^{2t}$

$$2^{t+3} = 2^{2t}$$

$$t + 3 = 2t$$

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$
Find a common base: $2^{3} \cdot 2^{t} = 2^{2t}$

$$2^{t+3} = 2^{2t}$$

$$t+3=2t$$

t = 3.

A farmer buys 48 chickens and 6 rabbits. The chicken population doubles each year, and the rabbit population doubles every six months. When does the farmer have the same number of chickens as rabbits?

Let c(t) denote the number of chickens after t years, and let r(t) denote the number of rabbits after t years.

Solve for
$$t$$
: $c(t) = r(t)$

$$48 \cdot 2^{t} = 6 \cdot 4^{t}$$

$$8 \cdot 2^{t} = 4^{t}$$
Find a common base: $2^{3} \cdot 2^{t} = 2^{2t}$

$$2^{t+3} = 2^{2t}$$

$$t+3 = 2t$$

Therefore the chicken and rabbit populations are equal after 3 years.

t=3

$$9^x = 2 \cdot 3^x + 63$$

$$9^x = 2 \cdot 3^x + 63$$
$$9^x - 2 \cdot 3^x - 63 = 0$$

$$9^{x} = 2 \cdot 3^{x} + 63$$

 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $2^{x} - 2u - 63 = 0$

$$9^{x} = 2 \cdot 3^{x} + 63$$

 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $2u - 63 = 0$

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$

$$9^{x} = 2 \cdot 3^{x} + 63$$

 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$
(?)(?) = 0

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
Substitute $u = 3^{x}$

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $u = 9 \text{ or } u = -7$

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $u = 9 \text{ or } u = -7$
 $3^{x} = 9 \text{ or } 3^{x} = -7$

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $u = 9 \text{ or } u = -7$
 $3^{x} = 9 \text{ or } 3^{x} = -7$
 $x = ?$

$$9^{x} = 2 \cdot 3^{x} + 63$$

 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $u = 9 \text{ or } u = -7$
 $3^{x} = 9 \text{ or } 3^{x} = -7$
 $x = 2$

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $u = 9 \text{ or } u = -7$
 $3^{x} = 9 \text{ or } 3^{x} = -7$
 $x = 2$
?

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$ | Substitute $u = 3^{x}$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $u = 9 \text{ or } u = -7$
 $3^{x} = 9 \text{ or } 3^{x} = -7$
 $x = 2$ no real solution

Solve for x.

$$9^{x} = 2 \cdot 3^{x} + 63$$
 $9^{x} - 2 \cdot 3^{x} - 63 = 0$
 $u^{2} - 2u - 63 = 0$
 $(u - 9)(u + 7) = 0$
 $y = 0$
Substitute $y = 3^{x}$
 $y = 0$
Substitute $y = 3^{x}$
 $y = 0$
Substitute $y = 0$
 $y = 0$
Substitute $y = 0$
Su

Therefore x = 2 is the solution.

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^{x} = u$.

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = ?$.

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2-3u-4=0$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
(?) (?) = 0

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2-3u-4=0$$

$$(u-4)(u+1)=0$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$

$$(u - 4) (u + 1) = 0$$

$$u = 4$$

or

$$u = -1$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

Set $e^x = u$. Then $e^{2x} = u^2$.

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

or

$$u=4$$

$$e^{x} = 4$$
 or

$$u = -1$$

$$e^{x} = -1$$

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

$$u = 4$$
 or $u = -1$
 $e^x = 4$ or $e^x = -1$
 $x = \ln 4$ or no real solution

Solve the equation

$$e^{2x} - 3e^x - 4 = 0$$

$$u^2 - 3u - 4 = 0$$
$$(u - 4) (u + 1) = 0$$

$$u=4$$
 or $u=-1$
 $e^x=4$ or $e^x=-1$
 $x=\ln 4$ or no real solution
 $x\approx 1.3863$

$$4^{x+1} - 2^{x+2} - 3 = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = ?$$
.

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
.

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
.

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = ?$,

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$,

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$, $2^{x+2} = ?$.

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(?)$$
 $(?)$ $=$ 0

$$4^{x+1} - 2^{x+2} - 3 = 0$$

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.

$$4u^2 - 4u - 3 = 0$$

$$(2u-3)(2u+1) = 0$$

Solve the equation

Set
$$u = 2^x$$
. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$
$$(2u - 3)(2u + 1) = 0$$

 $4^{x+1} - 2^{x+2} - 3 = 0$

2u - 3 = 0 or 2u + 1 = 0

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = 0$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right)$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$
or no real solution

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln(?)}{\ln?}$$
or no real solution

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln\left(\frac{3}{2}\right)}{\ln 2} \text{ or no real solution}$$

$$4^{x+1} - 2^{x+2} - 3 = 0$$
Set $u = 2^x$. Then $4^{x+1} = 4u^2$, $2^{x+2} = 4u$.
$$4u^2 - 4u - 3 = 0$$

$$(2u - 3)(2u + 1) = 0$$

$$2u - 3 = 0 \text{ or } 2u + 1 = 0$$

$$u = \frac{3}{2} \text{ or } u = -\frac{1}{2}$$

$$2^x = \frac{3}{2} \text{ or } 2^x = -\frac{1}{2}$$

$$x = \log_2\left(\frac{3}{2}\right) = \frac{\ln\left(\frac{3}{2}\right)}{\ln 2} \approx 0.58496 \text{ or no real solution}$$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1}$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1}$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

> 2 terms
$$\Rightarrow$$

transfer one side
 $3^{2x} = u$
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$
$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$
$$u - 2 - 63u^{-1} = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$
$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$
$$4 - 2 - 63 \cdot 3^{-1} = 0$$

> 2 terms
$$\Rightarrow$$

transfer one side
 $3^{2x} = u$
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(?)(?) = 0$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

 $3^{2x} = 2 + 63 \cdot 3^{-2x}$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

> 2 terms
$$\Rightarrow$$

transfer one side
 $3^{2x} = u$
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$
Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

> 2 terms
$$\Rightarrow$$

transfer one side
 $3^{2x} = u$
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$
Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$
or $u = -7$

> 2 terms
$$\Rightarrow$$

transfer one side
 $3^{2x} = u$
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$
Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$
or
$$u = -7$$
or no real solution

> 2 terms
$$\Rightarrow$$

transfer one side
 $3^{2x} = u$
 $3^{-2x} = (3^{2x})^{-1} = u^{-1}$
Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$
or no real solution

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

> 2 terms
$$\Rightarrow$$
 transfer one side $3^{2x} = u$ $3^{-2x} = (3^{2x})^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$2x = ?$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$2x = 2$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$

$$2x = 2$$

$$x = 1$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$ Multiply $\cdot u$

$$3^{2x} = 2 + 63 \cdot 3^{-2x}$$

$$3^{2x} - 2 - 63 \cdot 3^{-2x} = 0$$

$$u - 2 - 63u^{-1} = 0$$

$$u^{2} - 2u - 63 = 0$$

$$(u - 9)(u + 7) = 0$$

$$u - 9 = 0 \text{ or } u + 7 = 0$$

$$u = 9 \text{ or } u = -7$$

$$3^{2x} = 9 \text{ or no real solution}$$

$$2x = \log_{3} 9$$


$$2x = 2$$

$$x = 1$$

$$>$$
 2 terms \Rightarrow transfer one side $3^{2x} = u$ $3^{-2x} = \left(3^{2x}\right)^{-1} = u^{-1}$ Multiply $\cdot u$


Find
$$f^{-1}(x)$$
 for

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.




Find
$$f^{-1}(x)$$
 for

$$f(x)=\frac{e^{x}-e^{-x}}{e^{x}+e^{-x}}.$$



$$\frac{e^x-e^{-x}}{e^x+e^{-x}}=y$$


Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{(u - ?)}{(u + ?)} = y$$

Set $u = e^x$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.




$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{(u - ?)}{(u + ?)} = y$$

Set
$$u = e^x$$

 $e^{-x} =$?

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{\left(u - \frac{1}{u}\right)}{\left(u + \frac{1}{u}\right)} = y$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.

$$y = f(x)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{\left(u - \frac{1}{u}\right) u}{\left(u + \frac{1}{u}\right) u} = y$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.


$$y = f(x)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$
$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$
$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$


$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$


$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$


$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$


$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$


$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$


$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$


$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$2x = \ln\left(\frac{1 + y}{1 - y}\right)$$

$$\begin{vmatrix} \text{Set } u = e^x \\ e^{-x} = \frac{1}{e^x} = \frac{1}{u} \end{vmatrix}$$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$


$$e^{2x} = \frac{1 + y}{1 - y}$$

$$2x = \ln\left(\frac{1 + y}{1 - y}\right)$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



answer
$$f^{-1}(y) = \frac{1}{2} \ln \left(\frac{1+y}{1-y} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{\left(u - \frac{1}{u}\right)u}{\left(u + \frac{1}{u}\right)u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$


$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2}\ln\left(\frac{1 + y}{1 - y}\right)$$

Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for $f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$.



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln\left(\frac{1 + y}{1 - y}\right)$$


Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Find
$$f^{-1}(x)$$
 for

$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}.$$

f = tanh = hyperbolic tangent function.



Final answer, relabeled:

$$f^{-1}(x) = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right)$$

$$\frac{e^{x} - e^{-x}}{e^{x} + e^{-x}} = y$$

$$\frac{(u - \frac{1}{u}) u}{(u + \frac{1}{u}) u} = y$$

$$\frac{u^{2} - 1}{u^{2} + 1} = y$$

$$u^{2} - 1 = y(u^{2} + 1)$$

$$u^{2}(1 - y) = 1 + y$$

$$u^{2} = \frac{1 + y}{1 - y}$$

$$(e^{x})^{2} = \frac{1 + y}{1 - y}$$

$$e^{2x} = \frac{1 + y}{1 - y}$$

$$x = \frac{1}{2} \ln\left(\frac{1 + y}{1 - y}\right)$$


Set
$$u = e^x$$

 $e^{-x} = \frac{1}{e^x} = \frac{1}{u}$

Take In


Solve the inequality. $2^{-3x-5} < 7$

$$2^{-3x-5} < 7$$




Solve the inequality. 2^{-3x-5} < 7





Solve the inequality. $2^{-3x-5} < 7$


$$2^{-3x-5}$$
 < 7



Solve the inequality.

$$2^{-3x-5}$$
 < 7


$$\log_2 2^{-3x-5} < \log_2 7$$



Solve the inequality.

$$2^{-3x-5}$$
 < 7


$$\log_2 2^{-3x-5} < \log_2 7$$



Solve the inequality.

$$2^{-3x-5}$$
 < 7


$$\frac{\log_2 2^{-3x-5}}{-3x-5} < \log_2 7$$



Solve the inequality.

$$2^{-3x-5}$$
 < 7

$$\begin{array}{rcl} \log_2 2^{-3x-5} & < & \log_2 7 \\ -3x-5 & < & \log_2 7 \\ -3x & < & \log_2 7 + 5 \end{array}$$



Solve the inequality.

$$2^{-3x-5} < 7$$


$$\log_{2} 2^{-3x-5} < \log_{2} 7$$

$$-3x-5 < \log_{2} 7$$

$$-3x < \log_{2} 7+5$$

$$x > -\frac{\log_{2} 7+5}{3}$$

Logarithms preserve inequalities: apply log₂



Solve the inequality.

$$2^{-3x-5} < 7$$


$$\log_{2} 2^{-3x-5} < \log_{2} 7$$

$$-3x-5 < \log_{2} 7$$

$$-3x < \log_{2} 7+5$$

$$x > -\frac{\log_{2} 7+5}{3}$$

Logarithms preserve inequalities: apply log₂



Solve the inequality.


$$2^{-3x-5} < 7$$

$$\log_2 2^{-3x-5} < \log_2 7$$

$$\begin{array}{rcl}
-3x - 5 & < \log_2 7 \\
-3x & < \log_2 7 + 5 \\
x & > -\frac{\log_2 7 + 5}{3}
\end{array}$$

$$x \in \left(-\frac{5 + \log_2 7}{3}, \infty\right)$$

Logarithms preserve inequalities: apply log₂



Solve the inequality.


$$2^{-3x-5} < 7$$

$$\log_2 2^{-3x-5} < \log_2 7$$

$$\begin{array}{rcl}
-3x - 5 & < \log_2 7 \\
-3x & < \log_2 7 + 5 \\
x & > -\frac{\log_2 7 + 5}{3}
\end{array}$$

$$x \in \left(-\frac{5+\log_2 7}{3}, \infty\right)$$

Logarithms preserve inequalities: apply log₂



Solve the inequality. $2^{-3x-5} < 7$

$$\log_2 2^{-3x-5} < \log_2 7 \\ -3x-5 < \log_2 7$$

$$-3x < \log_2 7 + 5$$

$$x > -\frac{\log_2 7 + 5}{3}$$

$$x \in \left(-\frac{5 + \log_2 7}{3}, \infty\right)$$

Logarithms preserve inequalities: apply log₂

