

Precalculus

Lecture 18

Todor Milev

`https://github.com/tmilev/freecalc`

2020

Outline

1 Lines

- Slope-intercept Form
- Line intersection

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:

<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$

Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$(1, -2, 3) \in ?$

$(0, 5) \in$

$(0, 5, -2, 4, 0) \in$

$(0, 1, 2, 3, \dots, n) \in$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in ?$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in ?$$

$$(0, 1, 2, 3, \dots, n) \in$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in \mathbb{R}^5$$

$$(0, 1, 2, 3, \dots, n) \in$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in \mathbb{R}^5$$

$$(0, 1, 2, 3, \dots, n) \in ?$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ Definition (\mathbb{R}^2)

The set of ordered pairs of real numbers is denoted by \mathbb{R}^2 .

Definition (\mathbb{R}^3)

The set of ordered triples of real numbers is denoted by \mathbb{R}^3 .

Definition (\mathbb{R}^n)

The set of ordered n -tuples of real numbers is denoted by \mathbb{R}^n .

Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in \mathbb{R}^5$$

$$(0, 1, 2, 3, \dots, n) \in \mathbb{R}^{n+1}$$

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.
- A set of pairs of numbers (x, y) is called a line in \mathbb{R}^2 if it is the set of solutions to some linear equation.

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.
- A set of pairs of numbers (x, y) is called a line in \mathbb{R}^2 if it is the set of solutions to some linear equation.
- A set of points in the plane will be called a line if it is the graph of some linear equation.

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.
 - A set of pairs of numbers (x, y) is called a line in \mathbb{R}^2 if it is the set of solutions to some linear equation.
 - A set of points in the plane will be called a line if it is the graph of some linear equation.
-
- To introduce the Cartesian coordinate system we used informal, intuitive notions of point, lines and the plane.

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.
 - A set of pairs of numbers (x, y) is called a line in \mathbb{R}^2 if it is the set of solutions to some linear equation.
 - A set of points in the plane will be called a line if it is the graph of some linear equation.
-
- To introduce the Cartesian coordinate system we used informal, intuitive notions of point, lines and the plane.
 - We could (and often do in more advanced subjects) remove this informality by *defining* \mathbb{R}^2 to be the Euclidean plane.

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.
 - A set of pairs of numbers (x, y) is called a line in \mathbb{R}^2 if it is the set of solutions to some linear equation.
 - A set of points in the plane will be called a line if it is the graph of some linear equation.
-
- To introduce the Cartesian coordinate system we used informal, intuitive notions of point, lines and the plane.
 - We could (and often do in more advanced subjects) remove this informality by *defining* \mathbb{R}^2 to be the Euclidean plane.

Definition

- An equation of the form $ax + by + c = 0$, where a, b, c are constants such that a and b are not simultaneously zero, is called a linear equation.
 - A set of pairs of numbers (x, y) is called a line in \mathbb{R}^2 if it is the set of solutions to some linear equation.
 - A set of points in the plane will be called a line if it is the graph of some linear equation.
-
- To introduce the Cartesian coordinate system we used informal, intuitive notions of point, lines and the plane.
 - We could (and often do in more advanced subjects) remove this informality by *defining* \mathbb{R}^2 to be the Euclidean plane.

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$? \quad (y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we **plug in $(1, 3)$** and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by **$x = 1$** , **$y = 3$** is:

$$y - 3 = x - 1.$$

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$? \quad (y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides **become zero** when $x = 1, y = 3$.

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$? \quad (y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides **become zero** when $x = 1, y = 3$.

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$? \quad (y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity

$$(6 - 3) \neq (2 - 1)$$

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$? \quad (y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity

$$(6 - 3) \neq (2 - 1)$$

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$? \quad (y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above **we don't get an identity**

$$(6 - 3) \neq (2 - 1)$$

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$(2 - 1)(y - 3) = ? \quad (x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity, but that can be easily fixed:

$$(2 - 1)(6 - 3) \neq (2 - 1)$$

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$(2 - 1)(y - 3) = (6 - 3)(x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity, but that can be easily fixed:

$$(2 - 1)(6 - 3) = (6 - 3)(2 - 1)$$

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$(2 - 1)(y - 3) = (6 - 3)(x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity, but that can be easily fixed:

$$(2 - 1)(6 - 3) = (6 - 3)(2 - 1)$$

- Perhaps the last modification caused $x = 1, y = 3$ to no longer be solutions?

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$(2 - 1)(y - 3) = (6 - 3)(x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity, but that can be easily fixed:

$$(2 - 1)(6 - 3) = (6 - 3)(2 - 1)$$

- Perhaps the last modification caused $x = 1, y = 3$ to no longer be solutions? No - both sides are still zero when $x = 1, y = 3$.

Example

Find an equation of the line passing through $(1, 3)$ and $(2, 6)$.

$$(2 - 1)(y - 3) = (6 - 3)(x - 1)$$

- It suffices to manufacture a linear equation such that when we plug in $(1, 3)$ and $(2, 6)$ we get an identity.
- A (very simple) equation satisfied by $x = 1, y = 3$ is:

$$y - 3 = x - 1.$$

This is so because both sides become zero when $x = 1, y = 3$.

- If we plug in $x = 2$ and $y = 6$ in the above we don't get an identity, but that can be easily fixed:

$$(2 - 1)(6 - 3) = (6 - 3)(2 - 1)$$

- Perhaps the last modification caused $x = 1, y = 3$ to no longer be solutions? No - both sides are still zero when $x = 1, y = 3$.

Example

Find an equation of the line passing through (x_1, y_1) and (x_2, y_2) .

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1)$$

- It suffices to manufacture a linear equation such that when we plug in (x_1, y_1) and (x_2, y_2) we get an identity.
- A (very simple) equation satisfied by $x = x_1, y = y_2$ is:

$$y - y_2 = x - x_1.$$

This is so because both sides become zero when $x = x_1, y = y_1$.

- If we plug in $x = x_2$ and $y = y_2$ in the above we don't get an identity (necessarily), but that can be easily fixed:

$$(x_2 - x_1)(y_2 - y_1) = (y_2 - y_1)(x_2 - x_1)$$

- Perhaps the last modification caused $x = x_1, y = y_1$ to no longer be solutions? No - both sides are still zero when $x = x_1, y = y_1$.

Example

Find an equation of the line passing through (x_1, y_1) and (x_2, y_2) .

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1)$$

- It suffices to manufacture a linear equation such that when we plug in (x_1, y_1) and (x_2, y_2) we get an identity.
- A (very simple) equation satisfied by $x = x_1, y = y_2$ is:

$$y - y_2 = x - x_1.$$

This is so because both sides become zero when $x = x_1, y = y_1$.

- If we plug in $x = x_2$ and $y = y_2$ in the above we don't get an identity (necessarily), but that can be easily fixed:

$$(x_2 - x_1)(y_2 - y_1) = (y_2 - y_1)(x_2 - x_1)$$

- Perhaps the last modification caused $x = x_1, y = y_1$ to no longer be solutions? No - both sides are still zero when $x = x_1, y = y_1$.

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$y - y_1 = m(x - x_1)$$

point-slope form

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$y - y_1 = m(x - x_1)$$

point-slope form

$$y = m(x - x_1) + y_1$$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$y - y_1 = m(x - x_1)$$

point-slope form

$$y = m(x - x_1) + y_1$$

$$y = mx + y_1 - mx_1$$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$y - y_1 = m(x - x_1)$$

| *point-slope form*

$$y = m(x - x_1) + y_1$$

$$y = mx + y_1 - mx_1$$

Set $b = y_1 - mx_1$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$y - y_1 = m(x - x_1) \quad \left| \text{point-slope form} \right.$$

$$y = m(x - x_1) + y_1$$

$$y = mx + y_1 - mx_1$$

Set $b = y_1 - mx_1$

$$y = mx + b \quad \left| \text{slope-intercept form} \right.$$

Proposition

Let (x_1, y_1) and (x_2, y_2) be points and L be the line between them. Then L has equation

$$(x_2 - x_1)(y - y_1) = (y_2 - y_1)(x - x_1).$$

If $x_1 \neq x_2$, set $m = \frac{y_2 - y_1}{x_2 - x_1}$; then L has also equations

$$y - y_1 = \frac{y_2 - y_1}{x_2 - x_1}(x - x_1)$$

$$y - y_1 = m(x - x_1)$$

point-slope form

$$y = m(x - x_1) + y_1$$

$$y = mx + y_1 - mx_1$$

Set $b = y_1 - mx_1$

$$y = mx + b$$


slope-intercept form

Definition (non-vertical line, slope-intercept form)

A line that is the graph of an equation of the form

$$y = mx + b$$

is called a *non-vertical* line.




Definition (non-vertical line, slope-intercept form)

A line that is the graph of an equation of the form

$$y = mx + b$$

is called a *non-vertical* line.

- The equation above is called the slope-intercept form of the (non-vertical) line.




Definition (non-vertical line, slope-intercept form)

A line that is the graph of an equation of the form

$$y = mx + b$$

is called a *non-vertical* line.

- The equation above is called the slope-intercept form of the (non-vertical) line.
- The number m is called the slope of the line.




Definition (non-vertical line, slope-intercept form)

A line that is the graph of an equation of the form

$$y = mx + b$$

is called a *non-vertical* line.


- The equation above is called the slope-intercept form of the (non-vertical) line.
- The number m is called the slope of the line.
- The number b is the y intercept of the line.



Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)


$y = mx + b$, m - is called slope, b is called y -intercept.



Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y -intercept.




- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y -intercept.




- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points.

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y -intercept.




- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. **The run is assumed positive.**

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y -intercept.




- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points.


Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.




- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.



Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)


$y = mx + b$, m - is called slope, b is called y-intercept.



- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$y_2 = mx_2 + b$$



$$y_1 = mx_1 + b$$



Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y -intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl} y_2 & = & mx_2 + b \\ y_1 & = & mx_1 + b \\ \hline y_2 - y_1 & = & mx_2 + b - mx_1 - b \end{array}$$

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y -intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & mx_2 + \cancel{b} - mx_1 - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1)
 \end{array}$$

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & \cancel{mx_2} + \cancel{b} - \cancel{mx_1} - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1)
 \end{array}$$

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & mx_2 + \cancel{b} - mx_1 - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1) \\
 m & = & \frac{y_2 - y_1}{x_2 - x_1}
 \end{array}$$

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & mx_2 + \cancel{b} - mx_1 - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1) \\
 m & = & \frac{y_2 - y_1}{x_2 - x_1}
 \end{array}$$

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & mx_2 + \cancel{b} - mx_1 - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1) \\
 m & = & \frac{y_2 - y_1}{x_2 - x_1}
 \end{array}$$

Geometric Interpretation of Slope

Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.





- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the run of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the rise of the line between the two points. Negative rise is allowed.

$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & mx_2 + \cancel{b} - mx_1 - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1) \\
 m & = & \frac{y_2 - y_1}{x_2 - x_1} \\
 m & = & \frac{\text{rise}}{\text{run}}
 \end{array}$$

Geometric Interpretation of Slope


Definition (non-vertical line, slope-intercept form)

$y = mx + b$, m - is called slope, b is called y-intercept.




- Fix pts. (x_1, y_1) , (x_2, y_2) on the line with $x_2 > x_1$.
- Call $x_2 - x_1$ the **run** of the line between the points. The run is assumed positive.
- Call $y_2 - y_1$ the **rise** of the line between the two points. Negative rise is allowed.


$$\begin{array}{rcl}
 y_2 & = & mx_2 + b \\
 y_1 & = & mx_1 + b \\
 \hline
 y_2 - y_1 & = & mx_2 + \cancel{b} - mx_1 - \cancel{b} \\
 y_2 - y_1 & = & m(x_2 - x_1) \\
 m & = & \frac{y_2 - y_1}{x_2 - x_1} \\
 m & = & \frac{\text{rise}}{\text{run}}
 \end{array}$$




- If two linear functions have positive slopes, the one with the bigger slope increases faster.




- If two linear functions have positive slopes, the one with the bigger slope increases faster.
- $y = 2x$ increases twice as fast as $y = x$.




- If two linear functions have positive slopes, the one with the bigger slope increases faster.
- $y = 2x$ increases twice as fast as $y = x$.
- $y = 3x$ increases three times as fast as $y = x$.




- If two linear functions have positive slopes, the one with the bigger slope increases faster.
- $y = 2x$ increases twice as fast as $y = x$.
- $y = 3x$ increases three times as fast as $y = x$.
- $y = \frac{x}{2}$ increases half as fast as $y = x$.




- If two linear functions have positive slopes, the one with the bigger slope increases faster.
- $y = 2x$ increases twice as fast as $y = x$.
- $y = 3x$ increases three times as fast as $y = x$.
- $y = \frac{x}{2}$ increases half as fast as $y = x$.
- $y = \frac{x}{3}$ increases one third as fast as $y = x$.




- If two linear functions have negative slopes, the one with the lower slope decreases faster.




- If two linear functions have negative slopes, the one with the lower slope decreases faster.
- $y = -2x$ decreases twice as fast as $y = -x$.



- If two linear functions have negative slopes, the one with the lower slope decreases faster.
- $y = -2x$ decreases twice as fast as $y = -x$.
- $y = -3x$ decreases three times as fast as $y = -x$.




- If two linear functions have negative slopes, the one with the lower slope decreases faster.
- $y = -2x$ decreases twice as fast as $y = -x$.
- $y = -3x$ decreases three times as fast as $y = -x$.
- $y = -\frac{1}{2}x$ decreases half as fast as $y = -x$.



- If two linear functions have negative slopes, the one with the lower slope decreases faster.
- $y = -2x$ decreases twice as fast as $y = -x$.
- $y = -3x$ decreases three times as fast as $y = -x$.
- $y = -\frac{1}{2}x$ decreases half as fast as $y = -x$.
- $y = -\frac{1}{3}x$ decreases one third as fast as $y = -x$.


Definition (Vertical line)

A line of the form $x = a$ is called a vertical line.



Definition (Vertical line)


A line of the form $x = a$ is called a vertical line.



- y does not participate directly in the equation $x = a$.

Definition (Vertical line)


A line of the form $x = a$ is called a vertical line.



- y does not participate directly in the equation $x = a$.
- Therefore the equation cannot be rewritten in slope-intercept form ($y = ?x + ?$).

Definition (Vertical line)

A line of the form $x = a$ is called a vertical line.



- y does not participate directly in the equation $x = a$.
- Therefore the equation cannot be rewritten in slope-intercept form ($y = ?x + ?$).
- Consequently the notion of a slope is not undefined for vertical lines.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.


- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
----------	-----	-------------

$$x + y = 1$$

$$2x - y = 3$$

$$y = 2$$

$$x = -1$$


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation

pt.

another pt.

$$x + y = 1$$

?

?

$$2x - y = 3$$

$$y = 2$$

$$x = -1$$


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
----------	-----	-------------

$x + y = 1$	$(1, 0)$	$(0, 1)$
-------------	----------	----------

$2x - y = 3$		
--------------	--	--

$y = 2$		
---------	--	--

$x = -1$		
----------	--	--


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
----------	-----	-------------

$x + y = 1$	$(1, 0)$	$(0, 1)$
-------------	----------	----------

$$2x - y = 3$$

$$y = 2$$

$$x = -1$$


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
----------	-----	-------------

$x + y = 1$	$(1, 0)$	$(0, 1)$
-------------	----------	----------

$2x - y = 3$?	?
--------------	---	---

$$y = 2$$

$$x = -1$$


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$		
$x = -1$		


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$		
$x = -1$		


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$?	?
$x = -1$		


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$	$(0, 2)$	$(1, 2)$
$x = -1$		


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$	$(0, 2)$	$(1, 2)$
$x = -1$		


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$	$(0, 2)$	$(1, 2)$
$x = -1$?	?


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$	$(0, 2)$	$(1, 2)$
$x = -1$	$(-1, 0)$	$(-1, 1)$


Other points can be used as well.

Plotting Lines from line equation

To plot a line from its equation $ax + by = c$ do the following.

- If $b = 0$, the line is vertical through $x = \frac{c}{a}$. Suppose $b \neq 0$.
 - Plug in arbitrary number for x and find y from $y = \frac{c-ax}{b}$.
 - Use same procedure to find a second point on the line.
 - If $a \neq 0$: can also plug in values for y to find x .
 - Draw a line between the two dots.

Example



Plot the line with the given equation.

equation	pt.	another pt.
$x + y = 1$	$(1, 0)$	$(0, 1)$
$2x - y = 3$	$(0, -3)$	$(\frac{3}{2}, 0)$
$y = 2$	$(0, 2)$	$(1, 2)$
$x = -1$	$(-1, 0)$	$(-1, 1)$

Other points can be used as well.

Example

Find an equation of a line passing through the indicated pairs of points.

- $(1, 2)$ and $(2, -1)$.
- $(1, 1)$ and $(2, -2)$.
- $(0, 1)$ and $(1, 0)$.
- $(3, 5)$ and $(7, -11)$.

Example

Find an equation of the line passing through $(1, 2)$ with slope $-\frac{1}{2}$.

To find the intersection of two lines (if they do intersect) with equations $a_1x + b_1y + c_1 = 0$ and $a_2x + b_2y + c_2 = 0$ we need to solve the system of equations

$$\begin{cases} a_1x + b_1y + c_1 = 0 \\ a_2x + b_2y + c_2 = 0 \end{cases}$$

Example

Find the intersection of the following lines.

- ① $x - y = 3$ and $x + 2y = 10$.
- ② $3x - y = 3$ and $x = 1 - 3y$.
- ③ Line $x = 3$ and $x = 1 - 2y$
- ④ Line through $(2, 0)$ and $(1, 2)$ and line through $(3, 7)$ and $(2, 5)$.
- ⑤ Line through $(3, -1)$ and $(-1, 3)$ and line through $(1, 1)$ and $(2, 3)$.

Definition

Two lines are parallel if they have no common point.

Proposition

Two non-vertical lines are parallel if and only if they have equal slopes and different y intercepts.

Proof \Leftarrow .

- Suppose the two lines have different y intercepts and have the same slope m .
- Then the lines have equations as shown below.

$$\left| \begin{array}{l} y = mx + b_1 \\ y = mx + b_2 \end{array} \right.$$

- System has no solutions as $b_1 \neq b_2 \Rightarrow$ the lines don't intersect. \square

Definition

Two lines are parallel if they have no common point.

Proposition

Two non-vertical lines are parallel if and only if they have equal slopes and different y intercepts.

Proof \Rightarrow .

- Suppose the two lines have different slopes.
- Suppose the lines have equations as shown below.

$$\begin{array}{r} y = m_1 x + b_1 \\ - \quad y = m_2 x + b_2 \\ \hline 0 = (m_1 - m_2)x + b_1 - b_2 \\ (m_1 - m_2)x = b_2 - b_1 \quad \Bigg| \text{ Div. by } m_1 - m_2 \neq 0 \\ x = \frac{b_2 - b_1}{m_1 - m_2} \end{array}$$

- The system has solution $x = \frac{b_2 - b_1}{m_1 - m_2}$, $y = m_1 \frac{b_2 - b_1}{m_1 - m_2} + b_1 \Rightarrow$ the lines intersect.

