

Calculus III

Lecture 10

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1 Multivariable Chain Rule

Outline

- 1 Multivariable Chain Rule
- 2 Directional Derivatives via the Chain Rule

Outline

- 1 Multivariable Chain Rule
- 2 Directional Derivatives via the Chain Rule
- 3 Gradient

Outline

- 1 Multivariable Chain Rule
- 2 Directional Derivatives via the Chain Rule
- 3 Gradient
- 4 Differential Operators
 - Differential Operators Variable Changes

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
`https://creativecommons.org/licenses/by/3.0/us/`
and the links therein.

Multivariable Chain Rule Motivation

Recall:

- f , differentiable function,
- $\mathbf{u} = (u_1, u_2, u_3)$, unit vector,
- $P(x_0, y_0, z_0)$, point.

What is the rate of change of f at P in the direction \mathbf{u} ?

Multivariable Chain Rule Motivation

Recall:

- f , differentiable function,
- $\mathbf{u} = (u_1, u_2, u_3)$, unit vector,
- $P(x_0, y_0, z_0)$, point.

What is the rate of change of f at P in the direction \mathbf{u} ?

Directional derivative

$$(D_{\mathbf{u}}f)(P) = \left. \frac{d}{dt} \right|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Multivariable Chain Rule Motivation

Recall:

- f , differentiable function,
- $\mathbf{u} = (u_1, u_2, u_3)$, unit vector,
- $P(x_0, y_0, z_0)$, point.

What is the rate of change of f at P in the direction \mathbf{u} ?

Directional derivative

$$(D_{\mathbf{u}}f)(P) = \left. \frac{d}{dt} \right|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

More general, if

- $w = w(x, y, z)$;
- $x = x(t)$, $y = y(t)$, $z = z(t)$,

and all the functions are differentiable, how do we compute $\frac{dw}{dt}$?

Chain Rule

Differentials

$$dw = w_x(x, y, z)dx + w_y(x, y, z)dy + w_z(x, y, z)dz$$

and

$$dx = x'(t)dt \quad dy = y'(t)dt \quad dz = z'(t)dt .$$

Chain Rule

Differentials

$$dw = w_x(x, y, z)dx + w_y(x, y, z)dy + w_z(x, y, z)dz$$

and

$$dx = x'(t)dt \quad dy = y'(t)dt \quad dz = z'(t)dt .$$

Then

$$d(w) = (w_x x'(t) + w_y y'(t) + w_z z'(t)) dt$$

Chain Rule

Differentials

$$dw = w_x(x, y, z)dx + w_y(x, y, z)dy + w_z(x, y, z)dz$$

and

$$dx = x'(t)dt \quad dy = y'(t)dt \quad dz = z'(t)dt.$$

Then

$$d(w) = (w_x x'(t) + w_y y'(t) + w_z z'(t)) dt$$

Therefore

$$\frac{d}{dt}(w(x(t), y(t), z(t))) = \frac{\partial w}{\partial x}(x, y, z) \frac{dx}{dt} + \frac{\partial w}{\partial y}(x, y, z) \frac{dy}{dt} + \frac{\partial w}{\partial z}(x, y, z) \frac{dz}{dt}$$

Chain Rule

Differentials

$$dw = w_x(x, y, z)dx + w_y(x, y, z)dy + w_z(x, y, z)dz$$

and

$$dx = x'(t)dt \quad dy = y'(t)dt \quad dz = z'(t)dt.$$

Then

$$d(w) = (w_x x'(t) + w_y y'(t) + w_z z'(t)) dt$$

Therefore

$$\frac{d}{dt}(w(x(t), y(t), z(t))) = \frac{\partial w}{\partial x}(x, y, z) \frac{dx}{dt} + \frac{\partial w}{\partial y}(x, y, z) \frac{dy}{dt} + \frac{\partial w}{\partial z}(x, y, z) \frac{dz}{dt}$$


Derivative of composition of functions \implies Chain Rule

Algebra of Chain rule - Tree Diagrams

- $w = w(x, y, z)$;
- $x = x(t)$, $y = y(t)$, $z = z(t)$,

$$\frac{dw}{dt}(t) = \frac{\partial w}{\partial x}(x, y, z) \frac{dx}{dt}(t) + \frac{\partial w}{\partial y}(x, y, z) \frac{dy}{dt}(t) + \frac{\partial w}{\partial z}(x, y, z) \frac{dz}{dt}(t)$$

Alternative way of arranging terms - tree diagram:



More General Chain Rule


More general formula:

- $w = F(x, y, z)$;
- $x = f(u, v)$, $y = g(u, v)$, $z = h(u, v)$.

$$w = F(f(u, v), g(u, v), h(u, v)) = G(u, v)$$

To compute $\frac{\partial w}{\partial u} = \frac{\partial G}{\partial u}$:

- arrange variables in a tree diagram:



Example: powerexponential

Let $f(x) = x^x$. Compute $f'(x)$.

- Calculus I method: logarithmic differentiation or $x^x = e^{x \ln x}$.
- Calculus III method: chain rule.

Example: powerexponential

Let $f(x) = x^x$. Compute $f'(x)$.

- Calculus I method: logarithmic differentiation or $x^x = e^{x \ln x}$.
- Calculus III method: chain rule.

Let $w = w(u, v) = u^v$ and $u = u(x) = x$, $v = v(x) = x$.

Then $f(x) = w(u(x), v(x))$ and

Directional Derivatives via the Chain Rule

- Let f differentiable function,
- Let $\mathbf{u} = (u_1, u_2, u_3)$, unit vector,
- Let $P(x_0, y_0, z_0)$, point.

Directional Derivatives via the Chain Rule

- Let f differentiable function,
- Let $\mathbf{u} = (u_1, u_2, u_3)$, unit vector,
- Let $P(x_0, y_0, z_0)$, point.

What is the rate of change of f at P in the direction \mathbf{u} ? Answer was studied: directional derivative.

Directional Derivatives via the Chain Rule

- Let f differentiable function,
- Let $\mathbf{u} = (u_1, u_2, u_3)$, unit vector,
- Let $P(x_0, y_0, z_0)$, point.

What is the rate of change of f at P in the direction \mathbf{u} ? Answer was studied: directional derivative.

$$(D_{\mathbf{u}}f)_{(x,y,z)=(x_0,y_0,z_0)} = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\left\{ \begin{array}{lcl} x & = & x_0 + tu_1 \\ y & = & y_0 + tu_2 \\ z & = & z_0 + tu_3 \end{array} \right. .$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\left\{ \begin{array}{l} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{array} \right. .$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{dw}{dt}\bigg|_{t=0}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} .$$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt}\bigg|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} \end{aligned}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and $\left\{ \begin{array}{l} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{array} \right.$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt}\bigg|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \end{aligned}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases}.$$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt}\bigg|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \end{aligned}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases}.$$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt}\bigg|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \end{aligned}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt} \Big|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and $\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} .$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt} \Big|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right) \Big|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \\ &= \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) \cdot \mathbf{u} \end{aligned}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} .$$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt}\bigg|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \\ &= \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) \cdot \mathbf{u} \end{aligned}$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt} \Big|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and $\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases}.$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt} \Big|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right) \Big|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \\ &= \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) \cdot \mathbf{u} \end{aligned}$$

Definition (∇f ("nabla of f "))

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right).$$

$$(D_{\mathbf{u}}f)(x_0, y_0, z_0) = \frac{d}{dt}\bigg|_{t=0} f(x_0 + tu_1, y_0 + tu_2, z_0 + tu_3)$$

Let $w = f(x, y, z)$ and
$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} .$$

$$\begin{aligned} (D_{\mathbf{u}}f)(x_0, y_0, z_0) &= \frac{dw}{dt}\bigg|_{t=0} \\ &= \left(\frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt} + \frac{\partial f}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} \\ &= \frac{\partial f}{\partial x} u_1 + \frac{\partial f}{\partial y} u_2 + \frac{\partial f}{\partial z} u_3 \\ &= \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) \cdot \mathbf{u} = \nabla f \cdot \mathbf{u} \end{aligned}$$

Definition (∇f ("nabla of f "))

$$\nabla f = \left(\frac{\partial f}{\partial x}, \frac{\partial f}{\partial y}, \frac{\partial f}{\partial z} \right) .$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1) .$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

The partial derivatives are

$$\begin{aligned} \frac{\partial f}{\partial x} &= \\ \frac{\partial f}{\partial y} &= \\ \frac{\partial f}{\partial z} &= \end{aligned}$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

The partial derivatives are

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{2x}{x^2 + 2y^2 - z^2} \\ \frac{\partial f}{\partial y} &= \frac{4y}{x^2 + 2y^2 - z^2} \\ \frac{\partial f}{\partial z} &= \frac{-2z}{x^2 + 2y^2 - z^2} \end{aligned}$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

The partial derivatives are

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{2x}{x^2 + 2y^2 - z^2} & \frac{\partial f}{\partial x}(2, 1, -1) &= \\ \frac{\partial f}{\partial y} &= \frac{4y}{x^2 + 2y^2 - z^2} & \frac{\partial f}{\partial y}(2, 1, -1) &= \\ \frac{\partial f}{\partial z} &= \frac{-2z}{x^2 + 2y^2 - z^2} & \frac{\partial f}{\partial z}(2, 1, -1) &= \end{aligned}$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

The partial derivatives are

$$\begin{aligned} \frac{\partial f}{\partial x} &= \frac{2x}{x^2 + 2y^2 - z^2} & \frac{\partial f}{\partial x}(2, 1, -1) &= \frac{4}{5} \\ \frac{\partial f}{\partial y} &= \frac{4y}{x^2 + 2y^2 - z^2} & \frac{\partial f}{\partial y}(2, 1, -1) &= \frac{4}{5} \\ \frac{\partial f}{\partial z} &= \frac{-2z}{x^2 + 2y^2 - z^2} & \frac{\partial f}{\partial z}(2, 1, -1) &= \frac{2}{5} \end{aligned}$$

$$\nabla f|_{(x,y,z)=(2,1,-1)} = \left(\frac{4}{5}, \frac{4}{5}, \frac{2}{5} \right)$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

$$\nabla f|_{(x,y,z)=(2,1,-1)} = (f_x, f_y, f_z)|_{(x,y,z)=(2,1,-1)} = \left(\frac{4}{5}, \frac{4}{5}, \frac{2}{5}\right)$$

$$(D_{\mathbf{u}}f)|_{(x,y,z)=(2,1,-1)} = \nabla f|_{(x,y,z)=(2,1,-1)} \cdot \mathbf{u} =$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

$$\nabla f|_{(x,y,z)=(2,1,-1)} = (f_x, f_y, f_z)|_{(x,y,z)=(2,1,-1)} = \left(\frac{4}{5}, \frac{4}{5}, \frac{2}{5}\right)$$

$$(D_{\mathbf{u}}f)|_{(x,y,z)=(2,1,-1)} = \nabla f|_{(x,y,z)=(2,1,-1)} \cdot \mathbf{u} = \frac{\sqrt{6}}{5}$$

Example

Find the directional derivative of $f(x, y, z) = \ln(x^2 + 2y^2 - z^2)$ at $P(2, 1, -1)$ in the direction $\mathbf{v} = (-1, 2, 1)$.

A unit vector in the direction of \mathbf{v} is

$$\mathbf{u} = \frac{1}{|\mathbf{v}|} \mathbf{v} = \frac{1}{\sqrt{6}}(-1, 2, 1).$$

$$\nabla f|_{(x,y,z)=(2,1,-1)} = (f_x, f_y, f_z)|_{(x,y,z)=(2,1,-1)} = \left(\frac{4}{5}, \frac{4}{5}, \frac{2}{5}\right)$$

$$(D_{\mathbf{u}}f)|_{(x,y,z)=(2,1,-1)} = \nabla f|_{(x,y,z)=(2,1,-1)} \cdot \mathbf{u} = \frac{\sqrt{6}}{5}$$

$(D_{\mathbf{u}}f)(2, 1, -1) > 0$ implies that if we start at $(2, 1, -1)$ and move in the direction \mathbf{u} , then f is increasing.

Gradient

- Let f be a differentiable function.
- At a given point P , in which direction does f increase the fastest?

Gradient

- Let f be a differentiable function.
- At a given point P , in which direction does f increase the fastest?
- What is that maximal rate of increase?

Gradient

- Let f be a differentiable function.
- At a given point P , in which direction does f increase the fastest?
- What is that maximal rate of increase?
- It can be shown that if the maximal rate of increase is strictly positive, then it is achieved in exactly one direction.

Gradient

- Let f be a differentiable function.
- At a given point P , in which direction does f increase the fastest?
- What is that maximal rate of increase?
- It can be shown that if the maximal rate of increase is strictly positive, then it is achieved in exactly one direction.

Definition

The *gradient vector* of f at P is the unique vector that has

- magnitude equal to the maximal rate of increase of f at P .
- if the magnitude is not zero, then the direction is the one in which f increases the fastest.

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
- if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
 - if $\text{magn.} \neq 0$, its **direction is the one in which f increases fastest**.
-
- **The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$.**

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
 - if magn. $\neq 0$, its direction is the one in which f increases fastest.
-
- Recall that $\nabla f = (f_x, f_y, f_z)$.
 - The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:
$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u}$$

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
 - if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.
-
- Recall that $\nabla f = (f_x, f_y, f_z)$.
 - The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:
$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha$$
where α is the angle between ∇f and \mathbf{u} .

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
 - if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.
-
- Recall that $\nabla f = (f_x, f_y, f_z)$.
 - The increase of f in **unit direction \mathbf{u}** is $D_{\mathbf{u}}f$. We have:
$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha = |\nabla f| \cos \alpha,$$
where α is the angle between ∇f and \mathbf{u} .

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
 - if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.
-
- Recall that $\nabla f = (f_x, f_y, f_z)$.
 - The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:

$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha = |\nabla f| \cos \alpha,$$
 where α is the angle between ∇f and \mathbf{u} .
 - If $|\nabla f| \neq 0$, then $(D_{\mathbf{u}}f)$ is maximal when $\cos \alpha = 1$, i.e., $\alpha = 0$.

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
- if $\text{mag.} \neq 0$, its direction is the one in which f increases fastest.
- Recall that $\nabla f = (f_x, f_y, f_z)$.
- The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:

$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha = |\nabla f| \cos \alpha,$$
 where α is the angle between ∇f and \mathbf{u} .
- If $|\nabla f| \neq 0$, then $(D_{\mathbf{u}}f)$ is maximal when $\cos \alpha = 1$, i.e., $\alpha = 0$.
- Therefore the maximum of $D_{\mathbf{u}}f$ is achieved for $\mathbf{u} = \frac{\nabla f}{|\nabla f|}$.

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
 - if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.
-
- Recall that $\nabla f = (f_x, f_y, f_z)$.
 - The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:

$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha = |\nabla f| \cos \alpha,$$
 where α is the angle between ∇f and \mathbf{u} .
 - If $|\nabla f| \neq 0$, then $(D_{\mathbf{u}}f)$ is maximal when $\cos \alpha = 1$, i.e., $\alpha = 0$.
 - Therefore the maximum of $D_{\mathbf{u}}f$ is achieved for $\mathbf{u} = \frac{\nabla f}{|\nabla f|}$.
 - The maximum of $D_{\mathbf{u}}f$ is then $|\nabla f| = |(f_x, f_y, f_z)|$.

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
- if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.
- Recall that $\nabla f = (f_x, f_y, f_z)$.
- The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:

$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha = |\nabla f| \cos \alpha,$$
 where α is the angle between ∇f and \mathbf{u} .
- If $|\nabla f| \neq 0$, then $(D_{\mathbf{u}}f)$ is maximal when $\cos \alpha = 1$, i.e., $\alpha = 0$.
- Therefore the maximum of $D_{\mathbf{u}}f$ is achieved for $\mathbf{u} = \frac{\nabla f}{|\nabla f|}$.
- The maximum of $D_{\mathbf{u}}f$ is then $|\nabla f| = |(f_x, f_y, f_z)|$.

Theorem (Coordinate Computation of gradient vector)

The gradient vector of f equals $\nabla f = (f_x, f_y, f_z)$.

Definition

The *gradient vector* of f at P is the unique vector such that:

- its magnitude equals the maximal rate of increase of f at P ;
- if $\text{magn.} \neq 0$, its direction is the one in which f increases fastest.
- Recall that $\nabla f = (f_x, f_y, f_z)$.
- The increase of f in unit direction \mathbf{u} is $D_{\mathbf{u}}f$. We have:

$$(D_{\mathbf{u}}f) = (\nabla f) \cdot \mathbf{u} = |\nabla f| \cdot |\mathbf{u}| \cos \alpha = |\nabla f| \cos \alpha,$$
 where α is the angle between ∇f and \mathbf{u} .
- If $|\nabla f| \neq 0$, then $(D_{\mathbf{u}}f)$ is maximal when $\cos \alpha = 1$, i.e., $\alpha = 0$.
- Therefore the maximum of $D_{\mathbf{u}}f$ is achieved for $\mathbf{u} = \frac{\nabla f}{|\nabla f|}$.
- The maximum of $D_{\mathbf{u}}f$ is then $|\nabla f| = |(f_x, f_y, f_z)|$.

Theorem (Coordinate Computation of gradient vector)

The *gradient vector* of f equals $\nabla f = (f_x, f_y, f_z)$.

- In view of preceding thm., the gradient of f is denoted by ∇f .

Covariant Derivative

- f : a differentiable function.
- Directional derivative $D_{\mathbf{u}}f$ = rate of change along straight line.

Covariant Derivative

- f : a differentiable function.
- Directional derivative $D_{\mathbf{u}}f$ = rate of change along straight line.
- Let γ : a smooth parametric curve.

Covariant Derivative

- f : a differentiable function.
- Directional derivative $D_{\mathbf{u}}f$ = rate of change along straight line.
- Let γ : a smooth parametric curve.
- Question: How does f change as we move along γ ?

Covariant Derivative

- f : a differentiable function.
- Directional derivative $D_{\mathbf{u}}f$ = rate of change along straight line.
- Let γ : a smooth parametric curve.
- Question: How does f change as we move along γ ?

Definition

The rate of change of $f(\gamma(t))$ with respect to t is called the *covariant derivative* of f along γ and is denoted by $\nabla_{\gamma'} f$.

Covariant Derivative

- f : a differentiable function.
- Directional derivative $D_{\mathbf{u}}f$ = rate of change along straight line.
- Let γ : a smooth parametric curve.
- Question: How does f change as we move along γ ?

Definition

The rate of change of $f(\gamma(t))$ with respect to t is called the *covariant derivative* of f along γ and is denoted by $\nabla_{\gamma'} f$.

We can compute the covariant derivative using the chain rule:

$$(\nabla_{\gamma'(t_0)} f)(\gamma(t_0)) = \frac{d}{dt}\bigg|_{t=t_0} f(\gamma(t)) = (\nabla f)_{\gamma(t_0)} \cdot \gamma'(t_0) .$$

Covariant Derivative

- f : a differentiable function.
- Directional derivative $D_{\mathbf{u}}f$ = rate of change along straight line.
- Let γ : a smooth parametric curve.
- Question: How does f change as we move along γ ?

Definition

The rate of change of $f(\gamma(t))$ with respect to t is called the *covariant derivative* of f along γ and is denoted by $\nabla_{\gamma'} f$.

We can compute the covariant derivative using the chain rule:

$$(\nabla_{\gamma'(t_0)} f)(\gamma(t_0)) = \frac{d}{dt}\bigg|_{t=t_0} f(\gamma(t)) = (\nabla f)_{\gamma(t_0)} \cdot \gamma'(t_0) .$$

If \mathbf{u} is a unit vector, $\gamma(t_0) = P$ and $\gamma'(t_0) = \mathbf{u}$, then:

$$(D_{\mathbf{u}} f)(P) = (\nabla f)_P \cdot \mathbf{u} = (\nabla f)_{\gamma(t_0)} \cdot \gamma'(t_0) = \frac{d}{dt}\bigg|_{t=t_0} f(\gamma(t)) .$$

Gradient in Polar Coordinates

$\mathbf{e}_r = \mathbf{e}_r(P)$ and $\mathbf{e}_\theta = \mathbf{e}_\theta(P)$ are the polar fundamental directions at P

$$(\nabla f)_P = a\mathbf{e}_r + b\mathbf{e}_\theta$$

Gradient in Polar Coordinates

$\mathbf{e}_r = \mathbf{e}_r(P)$ and $\mathbf{e}_\theta = \mathbf{e}_\theta(P)$ are the polar fundamental directions at P

$$(\nabla f)_P = a\mathbf{e}_r + b\mathbf{e}_\theta$$

\mathbf{e}_r and \mathbf{e}_θ perpendicular unit vectors \implies

$$a = (\nabla f)_P \cdot \mathbf{e}_r = (D_{\mathbf{e}_r} f)(P)$$

$$b = (\nabla f)_P \cdot \mathbf{e}_\theta = (D_{\mathbf{e}_\theta} f)(P)$$

Gradient in Polar Coordinates

$\mathbf{e}_r = \mathbf{e}_r(P)$ and $\mathbf{e}_\theta = \mathbf{e}_\theta(P)$ are the polar fundamental directions at P

$$(\nabla f)_P = a\mathbf{e}_r + b\mathbf{e}_\theta$$

\mathbf{e}_r and \mathbf{e}_θ perpendicular unit vectors \implies

$$a = (\nabla f)_P \cdot \mathbf{e}_r = (D_{\mathbf{e}_r} f)(P)$$

$$b = (\nabla f)_P \cdot \mathbf{e}_\theta = (D_{\mathbf{e}_\theta} f)(P)$$

To compute $(D_{\mathbf{e}_r} f)(P)$ we use the line through $P(r_0, \theta_0)$ with direction \mathbf{e}_r , which in polar coordinates is given by $(r, \theta) = (t, \theta_0)$. Therefore

$$a = (D_{\mathbf{e}_r} f)(P) = \left. \frac{d}{dt} \right|_{t=r_0} f(t, \theta_0) = \frac{\partial f}{\partial r}(P).$$

Gradient in Polar Coordinates

$\mathbf{e}_r = \mathbf{e}_r(P)$ and $\mathbf{e}_\theta = \mathbf{e}_\theta(P)$ are the polar fundamental directions at P

$$(\nabla f)_P = a\mathbf{e}_r + b\mathbf{e}_\theta$$

\mathbf{e}_r and \mathbf{e}_θ perpendicular unit vectors \implies

$$a = (\nabla f)_P \cdot \mathbf{e}_r = (D_{\mathbf{e}_r} f)(P)$$

$$b = (\nabla f)_P \cdot \mathbf{e}_\theta = (D_{\mathbf{e}_\theta} f)(P)$$

To compute $(D_{\mathbf{e}_\theta} f)(P)$ we use the circle centered at the origin and passing through $P(r_0, \theta_0)$. The polar parametrization of this circle that has *unit* tangent at P is given by $(r, \theta) = (r_0, \frac{1}{r_0}t)$. Therefore

$$b = (D_{\mathbf{e}_\theta} f)(P) = \left. \frac{d}{dt} \right|_{t=\theta_0} f\left(r_0, \frac{1}{r_0}t\right) = \frac{1}{r_0} \frac{\partial f}{\partial \theta}(P).$$

Gradient in Polar Coordinates

$\mathbf{e}_r = \mathbf{e}_r(P)$ and $\mathbf{e}_\theta = \mathbf{e}_\theta(P)$ are the polar fundamental directions at P

$$(\nabla f)_P = a\mathbf{e}_r + b\mathbf{e}_\theta$$

\mathbf{e}_r and \mathbf{e}_θ perpendicular unit vectors \implies

$$a = (\nabla f)_P \cdot \mathbf{e}_r = (D_{\mathbf{e}_r} f)(P)$$

$$b = (\nabla f)_P \cdot \mathbf{e}_\theta = (D_{\mathbf{e}_\theta} f)(P)$$

From the previous computations:

$$(\nabla f)_P = \frac{\partial f}{\partial r}(P)\mathbf{e}_r + \frac{1}{r_0} \frac{\partial f}{\partial \theta}(P)\mathbf{e}_\theta ,$$

or

$$\nabla = \mathbf{e}_r \frac{\partial}{\partial r} + \mathbf{e}_\theta \frac{1}{r} \frac{\partial}{\partial \theta} .$$

Application

Let f be a function on the plane such that f depends only on the distance to a fixed point, O .

Application

Let f be a function on the plane such that f depends only on the distance to a fixed point, O .

In a polar coordinate system with origin at O we get $f(P) = g(r)$

$$\nabla = \mathbf{e}_r \frac{\partial}{\partial r} + \mathbf{e}_\theta \frac{1}{r} \frac{\partial}{\partial \theta} .$$

$$\nabla f = g'(r) \mathbf{e}_r = g'(r) \hat{\mathbf{r}} = \frac{g'(r)}{r} \mathbf{r} .$$

Application

Let f be a function on the plane such that f depends only on the distance to a fixed point, O .

In a polar coordinate system with origin at O we get $f(P) = g(r)$

$$\nabla = \mathbf{e}_r \frac{\partial}{\partial r} + \mathbf{e}_\theta \frac{1}{r} \frac{\partial}{\partial \theta} .$$

$$\nabla f = g'(r) \mathbf{e}_r = g'(r) \hat{\mathbf{r}} = \frac{g'(r)}{r} \mathbf{r} .$$

Example: $f(P) = |OP|^{-1} = r^{-1} = g(r)$. Then

$$\nabla f = g'(r) \mathbf{e}_r = -r^{-2} \mathbf{e}_r = -\frac{1}{r^3} \mathbf{r}$$

Application

Let f be a function on the plane such that f depends only on the distance to a fixed point, O .

In a polar coordinate system with origin at O we get $f(P) = g(r)$

$$\nabla = \mathbf{e}_r \frac{\partial}{\partial r} + \mathbf{e}_\theta \frac{1}{r} \frac{\partial}{\partial \theta} .$$

$$\nabla f = g'(r) \mathbf{e}_r = g'(r) \hat{\mathbf{r}} = \frac{g'(r)}{r} \mathbf{r} .$$

Example: $f(P) = |OP|^{-1} = r^{-1} = g(r)$. Then

$$\nabla f = g'(r) \mathbf{e}_r = -r^{-2} \mathbf{e}_r = -\frac{1}{r^3} \mathbf{r}$$

Problem: Let \mathbf{X} be a vector field of the form

$$\mathbf{X} = h(r) \mathbf{r}$$

for some continuous function h . Show that \mathbf{X} is a *gradient field*: there exists a smooth function f such that $\mathbf{X} = \nabla f$.

Gravity and Gradient

- Let an object move along surface $z = f(x, y)$.
- Let gravity \mathbf{G} be constant, $\mathbf{G} = -mg \mathbf{k}$.
- Normal to surface:

$$\mathbf{n} = (-f_x(x_0, y_0), -f_y(x_0, y_0), 1) = -\nabla f + \mathbf{k}$$

- Let \mathbf{F} be the component of \mathbf{G} effectively acting on the object. Object is restricted to the surface $\Rightarrow \mathbf{F}$ is the component of \mathbf{G} tangent to the surface.

$$\begin{aligned}\mathbf{F} &= \text{orth}_n \mathbf{G} = -mg \text{orth}_n \mathbf{k} \\ \text{orth}_n \mathbf{k} &= \mathbf{k} - \text{proj}_n \mathbf{k} = \mathbf{k} - \frac{\mathbf{k} \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n} = \mathbf{k} - \frac{1}{|\mathbf{n}|^2} (-\nabla f + \mathbf{k})\end{aligned}$$

Gravity and Gradient

- Let an object move along surface $z = f(x, y)$.
- Let gravity \mathbf{G} be constant, $\mathbf{G} = -mg \mathbf{k}$.
- Normal to surface:

$$\mathbf{n} = (-f_x(x_0, y_0), -f_y(x_0, y_0), 1) = -\nabla f + \mathbf{k}$$

- Let \mathbf{F} be the component of \mathbf{G} effectively acting on the object. Object is restricted to the surface $\Rightarrow \mathbf{F}$ is the component of \mathbf{G} tangent to the surface.

$$\begin{aligned}\mathbf{F} &= \text{orth}_n \mathbf{G} = -mg \text{orth}_n \mathbf{k} \\ \text{orth}_n \mathbf{k} &= \mathbf{k} - \text{proj}_n \mathbf{k} = \mathbf{k} - \frac{\mathbf{k} \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n} = \mathbf{k} - \frac{1}{|\mathbf{n}|^2} (-\nabla f + \mathbf{k})\end{aligned}$$

Horizontal component of \mathbf{F} :

$$\frac{mg}{1 + |\nabla f|^2} (-\nabla f)$$

Gravity and Gradient

- Let an object move along surface $z = f(x, y)$.
- Let gravity \mathbf{G} be constant, $\mathbf{G} = -mg \mathbf{k}$.
- Normal to surface:

$$\mathbf{n} = (-f_x(x_0, y_0), -f_y(x_0, y_0), 1) = -\nabla f + \mathbf{k}$$

- Let \mathbf{F} be the component of \mathbf{G} effectively acting on the object. Object is restricted to the surface $\Rightarrow \mathbf{F}$ is the component of \mathbf{G} tangent to the surface.

$$\begin{aligned}\mathbf{F} &= \text{orth}_n \mathbf{G} = -mg \text{orth}_n \mathbf{k} \\ \text{orth}_n \mathbf{k} &= \mathbf{k} - \text{proj}_n \mathbf{k} = \mathbf{k} - \frac{\mathbf{k} \cdot \mathbf{n}}{|\mathbf{n}|^2} \mathbf{n} = \mathbf{k} - \frac{1}{|\mathbf{n}|^2} (-\nabla f + \mathbf{k})\end{aligned}$$

Horizontal component of \mathbf{F} :

$$\frac{mg}{1 + |\nabla f|^2} (-\nabla f)$$

Gravity pulls object in the direction of fastest descent.

Differential operators definition

- Let D be an open set in the plane.

Differential operators definition

- Let D be an open set in the plane.
- Let $\mathcal{C}^\infty(D)$ denote the set of infinitely differentiable f-ns over D .

Differential operators definition

- Let D be an open set in the plane.
- Let $\mathcal{C}^\infty(D)$ denote the set of infinitely differentiable f-ns over D .

Definition

The two-variable differential operators $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ are the maps from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ given by: $\frac{\partial}{\partial x}(f) = \frac{\partial f}{\partial x}$ and $\frac{\partial}{\partial y}(f) = \frac{\partial f}{\partial y}$ for every function $f \in \mathcal{C}^\infty(D)$.

Differential operators definition

- Let D be an open set in the plane.
- Let $\mathcal{C}^\infty(D)$ denote the set of infinitely differentiable f-ns over D .

Definition

The two-variable differential operators $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ are the maps from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ given by: $\frac{\partial}{\partial x}(f) = \frac{\partial f}{\partial x}$ and $\frac{\partial}{\partial y}(f) = \frac{\partial f}{\partial y}$ for every function $f \in \mathcal{C}^\infty(D)$.

- The operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is defined via

$$\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}(f) = \underbrace{\frac{\partial}{\partial x} \cdots \frac{\partial}{\partial x}}_{n \text{ times}} \underbrace{\frac{\partial}{\partial y} \cdots \frac{\partial}{\partial y}}_{m \text{ times}}(f)$$

Differential operators definition

- Let D be an open set in the plane.
- Let $\mathcal{C}^\infty(D)$ denote the set of infinitely differentiable f-ns over D .

Definition

The two-variable differential operators $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ are the maps from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ given by: $\frac{\partial}{\partial x}(f) = \frac{\partial f}{\partial x}$ and $\frac{\partial}{\partial y}(f) = \frac{\partial f}{\partial y}$ for every function $f \in \mathcal{C}^\infty(D)$.

- The operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is defined via

$$\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}(f) = \underbrace{\frac{\partial}{\partial x} \cdots \frac{\partial}{\partial x}}_{n \text{ times}} \underbrace{\frac{\partial}{\partial y} \cdots \frac{\partial}{\partial y}}_{m \text{ times}}(f)$$

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

- For $n = 0, m = 0$, the operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is by definition equal to 1.

Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

- For $n = 0, m = 0$, the operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is by definition equal to 1.
- A function g in \mathcal{C}^∞ gives rise to a differential operator via multiplication: $(g \cdot f)(x) = (gf)(x) = g(x)f(x)$.

Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

- For $n = 0, m = 0$, the operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is by definition equal to 1.
- A function g in \mathcal{C}^∞ gives rise to a differential operator via multiplication: $(g \cdot f)(x) = (gf)(x) = g(x)f(x)$.
- Functions are by definition zero-order differential operators.

Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

- For $n = 0, m = 0$, the operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is by definition equal to 1.
- A function g in \mathcal{C}^∞ gives rise to a differential operator via multiplication: $(g \cdot f)(x) = (gf)(x) = g(x)f(x)$.
- Functions are by definition zero-order differential operators.
- The number $m + n$ is defined to be the order of the differential operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$.

Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

- For $n = 0, m = 0$, the operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is by definition equal to 1.
- A function g in \mathcal{C}^∞ gives rise to a differential operator via multiplication: $(g \cdot f)(x) = (gf)(x) = g(x)f(x)$.
- Functions are by definition zero-order differential operators.
- The number $m + n$ is defined to be the order of the differential operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$.
- The order of a differential operator ξ is the largest order of the differential operators appearing in the expression of ξ via $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$.


Differential operator notation

Definition (Smooth finite order differential operators)

A differential operator over D is a map from $\mathcal{C}^\infty(D)$ to $\mathcal{C}^\infty(D)$ obtained by sums of operators $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ with coefficients in $\mathcal{C}^\infty(D)$.

- For $n = 0, m = 0$, the operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$ is by definition equal to 1.
- A function g in \mathcal{C}^∞ gives rise to a differential operator via multiplication: $(g \cdot f)(x) = (gf)(x) = g(x)f(x)$.
- Functions are by definition zero-order differential operators.
- The number $m + n$ is defined to be the order of the differential operator $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$.
- The order of a differential operator ξ is the largest order of the differential operators appearing in the expression of ξ via $\frac{\partial^n}{\partial x^n} \frac{\partial^m}{\partial y^m}$.
- Analogous definitions exist for functions in n variables.


Example (Derivatives in polar coordinates)



Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\frac{\partial z}{\partial r} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r}$$

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\frac{\partial z}{\partial r} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r} = \frac{\partial f}{\partial x} \cos \theta + \frac{\partial f}{\partial y} \sin \theta$$

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\frac{\partial z}{\partial r} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r} = \frac{\partial f}{\partial x} \cos \theta + \frac{\partial f}{\partial y} \sin \theta$$

Recall that $\frac{\partial z}{\partial x} = \left(\frac{\partial f}{\partial x} \right) (x, y)$ and $\frac{\partial z}{\partial y} = \left(\frac{\partial f}{\partial y} \right) (x, y)$.

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\begin{aligned} \frac{\partial z}{\partial r} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r} = \frac{\partial f}{\partial x} \cos \theta + \frac{\partial f}{\partial y} \sin \theta \\ &= \cos \theta \frac{\partial z}{\partial x} + \sin \theta \frac{\partial z}{\partial y} . \end{aligned}$$

Recall that $\frac{\partial z}{\partial x} = \left(\frac{\partial f}{\partial x} \right) (x, y)$ and $\frac{\partial z}{\partial y} = \left(\frac{\partial f}{\partial y} \right) (x, y)$.

Example (Derivatives in polar coordinates)



Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.


- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\begin{aligned} \frac{\partial z}{\partial r} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r} = \frac{\partial f}{\partial x} \cos \theta + \frac{\partial f}{\partial y} \sin \theta \\ &= \cos \theta \frac{\partial z}{\partial x} + \sin \theta \frac{\partial z}{\partial y} . \end{aligned}$$

The above is true for all differentiable $z = f(x, y)$,

Recall that $\frac{\partial z}{\partial x} = \left(\frac{\partial f}{\partial x} \right) (x, y)$ and $\frac{\partial z}{\partial y} = \left(\frac{\partial f}{\partial y} \right) (x, y)$.

Example (Derivatives in polar coordinates)



Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.


- Compute $\frac{\partial z}{\partial r}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial r}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\begin{aligned} \frac{\partial z}{\partial r} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial r} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial r} = \frac{\partial f}{\partial x} \cos \theta + \frac{\partial f}{\partial y} \sin \theta \\ &= \cos \theta \frac{\partial z}{\partial x} + \sin \theta \frac{\partial z}{\partial y} . \end{aligned}$$

The above is true for all differentiable $z = f(x, y)$, therefore

$$\frac{\partial}{\partial r} = \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} .$$


Example (Derivatives in polar coordinates)



Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\frac{\partial z}{\partial \theta} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta}$$

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\frac{\partial z}{\partial \theta} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta} = \frac{\partial f}{\partial x} (-r \sin \theta) + \frac{\partial f}{\partial y} r \cos \theta$$

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\frac{\partial z}{\partial \theta} = \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta} = \frac{\partial f}{\partial x} (-r \sin \theta) + \frac{\partial f}{\partial y} r \cos \theta$$

Recall that $\frac{\partial z}{\partial x} = \left(\frac{\partial f}{\partial x} \right) (x, y)$ and $\frac{\partial z}{\partial y} = \left(\frac{\partial f}{\partial y} \right) (x, y)$.

Example (Derivatives in polar coordinates)




Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\begin{aligned} \frac{\partial z}{\partial \theta} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta} = \frac{\partial f}{\partial x} (-r \sin \theta) + \frac{\partial f}{\partial y} r \cos \theta \\ &= -r \sin \theta \frac{\partial z}{\partial x} + r \cos \theta \frac{\partial z}{\partial y} . \end{aligned}$$

Recall that $\frac{\partial z}{\partial x} = \left(\frac{\partial f}{\partial x}\right)(x, y)$ and $\frac{\partial z}{\partial y} = \left(\frac{\partial f}{\partial y}\right)(x, y)$.

Example (Derivatives in polar coordinates)



Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.


- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\begin{aligned}\frac{\partial z}{\partial \theta} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta} = \frac{\partial f}{\partial x} (-r \sin \theta) + \frac{\partial f}{\partial y} r \cos \theta \\ &= -r \sin \theta \frac{\partial z}{\partial x} + r \cos \theta \frac{\partial z}{\partial y}.\end{aligned}$$

The above is true for all differentiable $z = f(x, y)$,

Recall that $\frac{\partial z}{\partial x} = \left(\frac{\partial f}{\partial x} \right) (x, y)$ and $\frac{\partial z}{\partial y} = \left(\frac{\partial f}{\partial y} \right) (x, y)$.

Example (Derivatives in polar coordinates)



Let $x = r \cos \theta$, $y = r \sin \theta$ and $z = f(x, y)$.

- Compute $\frac{\partial z}{\partial \theta}$ via $\frac{\partial z}{\partial x}$ and $\frac{\partial z}{\partial y}$.
- Express the differential operator $\frac{\partial}{\partial \theta}$ via $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

$$\begin{aligned} \frac{\partial z}{\partial \theta} &= \frac{\partial f}{\partial x} \frac{\partial x}{\partial \theta} + \frac{\partial f}{\partial y} \frac{\partial y}{\partial \theta} = \frac{\partial f}{\partial x} (-r \sin \theta) + \frac{\partial f}{\partial y} r \cos \theta \\ &= -r \sin \theta \frac{\partial z}{\partial x} + r \cos \theta \frac{\partial z}{\partial y} . \end{aligned}$$

The above is true for all differentiable $z = f(x, y)$, therefore

$$\frac{\partial}{\partial \theta} = -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y} .$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.
We computed previously that

$$\begin{aligned}\frac{\partial}{\partial r} &= \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} \\ \frac{\partial}{\partial \theta} &= -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y} .\end{aligned}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.
We computed previously that

$$\begin{aligned}\frac{\partial}{\partial r} &= \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} \\ \frac{\partial}{\partial \theta} &= -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y} .\end{aligned}$$

This is a linear system in $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$.

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.
We computed previously that

$$\begin{aligned}\frac{\partial}{\partial r} &= \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} \\ \frac{\partial}{\partial \theta} &= -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y} .\end{aligned}$$

This is a linear system in $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$. To solve the system, **eliminate $\frac{\partial}{\partial x}$ by multiplying the first equality by $r \sin \theta$, the second by $\cos \theta$ and adding the two.**

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.
We computed previously that

$$\begin{aligned}\frac{\partial}{\partial r} &= \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} \\ \frac{\partial}{\partial \theta} &= -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y} .\end{aligned}$$

This is a linear system in $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$. To solve the system, eliminate $\frac{\partial}{\partial x}$ by multiplying the first equality by $r \sin \theta$, the second by $\cos \theta$ and adding the two. Similarly **eliminate $\frac{\partial}{\partial y}$ by multiplying the first equality by $-r \cos \theta$ and the second by $\sin \theta$** and adding the two.

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.
We computed previously that

$$\begin{aligned}\frac{\partial}{\partial r} &= \cos \theta \frac{\partial}{\partial x} + \sin \theta \frac{\partial}{\partial y} \\ \frac{\partial}{\partial \theta} &= -r \sin \theta \frac{\partial}{\partial x} + r \cos \theta \frac{\partial}{\partial y} .\end{aligned}$$

This is a linear system in $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$. To solve the system, eliminate $\frac{\partial}{\partial x}$ by multiplying the first equality by $r \sin \theta$, the second by $\cos \theta$ and adding the two. Similarly eliminate $\frac{\partial}{\partial y}$ by multiplying the first equality by $-r \cos \theta$ and the second by $\sin \theta$ and adding the two. Finally :

$$\begin{aligned}\frac{\partial}{\partial x} &= \cos \theta \frac{\partial}{\partial r} - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \\ \frac{\partial}{\partial y} &= \sin \theta \frac{\partial}{\partial r} + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta}\end{aligned}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Recall that

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Recall that

$$\tan \theta = \frac{r \sin \theta}{r \cos \theta}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Recall that

$$\tan \theta = \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Recall that

$$\tan \theta = \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right)\end{aligned}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$


Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x}$$



Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.


Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\tan \theta = \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x}$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

$$r = \sqrt{x^2 + y^2}$$

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} ?$$




Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta}$$



Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.


Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\tan \theta = \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x}$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

$$r = \sqrt{x^2 + y^2}$$

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} ?$$




Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$

$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$




Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$




$$\begin{aligned}\frac{\partial z}{\partial x} &= \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right) \\ &= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}\end{aligned}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$




$$\begin{aligned}\frac{\partial z}{\partial x} &= \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right) \\ &= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}\end{aligned}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$


$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$


$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} ? + \frac{\partial z}{\partial \theta} ?$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$


$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} ?$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$


$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} ?$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$


$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \frac{x}{x^2 + y^2}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$

$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \frac{x}{x^2 + y^2}$$


$$= \sin \theta \frac{\partial z}{\partial r} + \frac{\cos \theta}{r} \frac{\partial z}{\partial \theta}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$

$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \frac{x}{x^2 + y^2}$$


$$= \sin \theta \frac{\partial z}{\partial r} + \frac{\cos \theta}{r} \frac{\partial z}{\partial \theta}$$

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$

$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \frac{x}{x^2 + y^2}$$

$$= \sin \theta \frac{\partial z}{\partial r} + \frac{\cos \theta}{r} \frac{\partial z}{\partial \theta}$$


The above hold for all z ,

Example (Partial Derivatives in Polar Coordinates)

Express $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$ via $\frac{\partial}{\partial r}$ and $\frac{\partial}{\partial \theta}$, where $x = r \cos \theta$, $y = r \sin \theta$.

Suppose $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$. Recall that

$$\begin{aligned}\tan \theta &= \frac{r \sin \theta}{r \cos \theta} = \frac{y}{x} \\ \theta &= \arctan\left(\frac{y}{x}\right) \\ r &= \sqrt{x^2 + y^2}\end{aligned}$$



$$\frac{\partial z}{\partial x} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial x} = \frac{\partial z}{\partial r} \frac{x}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \left(\frac{-y}{x^2 + y^2} \right)$$

$$= \cos \theta \frac{\partial z}{\partial r} - \frac{\sin \theta}{r} \frac{\partial z}{\partial \theta}$$

$$\frac{\partial z}{\partial y} = \frac{\partial z}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial z}{\partial \theta} \frac{\partial \theta}{\partial y} = \frac{\partial z}{\partial r} \frac{y}{\sqrt{x^2 + y^2}} + \frac{\partial z}{\partial \theta} \frac{x}{x^2 + y^2}$$

$$= \sin \theta \frac{\partial z}{\partial r} + \frac{\cos \theta}{r} \frac{\partial z}{\partial \theta}$$

The above hold for all z , therefore

$$\begin{aligned}\frac{\partial}{\partial x} &= \cos \theta \frac{\partial}{\partial r} - \frac{1}{r} \sin \theta \frac{\partial}{\partial \theta} \\ \frac{\partial}{\partial y} &= \sin \theta \frac{\partial}{\partial r} + \frac{1}{r} \cos \theta \frac{\partial}{\partial \theta}\end{aligned}$$

The Laplace Operator

Definition

The n -variable Laplace operator is the differential operator:

$$\Delta = \sum_{j=1}^n \frac{\partial^2}{\partial x_j^2} = \frac{\partial^2}{\partial x_1^2} + \cdots + \frac{\partial^2}{\partial x_n^2} \quad .$$

In particular the two-variable Laplace operator is:

$$\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$$

The Laplace operator is named after Pierre Laplace (1749-1827).

Example

Express the Laplace operator $\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$ in polar coordinates.

$$\Delta = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2}{\partial \theta^2} .$$

Harmonic Functions

Recall that $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$

Definition (Harmonic function definition)

Functions f such that $\Delta f = 0$ are called *harmonic* functions.

Example

The function $f(x, y) = \ln(x^2 + y^2)$ is a harmonic function.

Harmonic Functions

Recall that $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$

Definition (Harmonic function definition)

Functions f such that $\Delta f = 0$ are called *harmonic* functions.

Example

The function $f(x, y) = \ln(x^2 + y^2)$ is a harmonic function. Rewrite in polar coordinates: $f(x, y) = g(r, \theta) = \ln(r^2) = 2 \ln r$. Then

$$\begin{aligned}\Delta g &= \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial g}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 g}{\partial \theta^2} = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial(2 \ln r)}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2(2 \ln r)}{\partial \theta^2} \\ &= \frac{1}{r} \frac{\partial}{\partial r} \left(r \cdot \frac{2}{r} \right) = \frac{1}{r} \frac{\partial}{\partial r} (2) = 0.\end{aligned}$$

Harmonic Functions

Recall that $\Delta = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2}$

Definition (Harmonic function definition)

Functions f such that $\Delta f = 0$ are called *harmonic* functions.

Example

The function $f(x, y) = \ln(x^2 + y^2)$ is a harmonic function. Rewrite in polar coordinates: $f(x, y) = g(r, \theta) = \ln(r^2) = 2 \ln r$. Then

$$\begin{aligned}\Delta g &= \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial g}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 g}{\partial \theta^2} = \frac{1}{r} \frac{\partial}{\partial r} \left(r \frac{\partial (2 \ln r)}{\partial r} \right) + \frac{1}{r^2} \frac{\partial^2 (2 \ln r)}{\partial \theta^2} \\ &= \frac{1}{r} \frac{\partial}{\partial r} \left(r \cdot \frac{2}{r} \right) = \frac{1}{r} \frac{\partial}{\partial r} (2) = 0.\end{aligned}$$

Fact: The only harmonic functions independent of θ are of the form $g(r, \theta) = c_1 \ln r + c_2$.