

Calculus I

Lecture 1

Construcing Functions

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1 A Catalog of Essential Functions

- Linear Functions
- Polynomials
- Power Functions
- Rational Functions
- Algebraic Functions
- Transcendental Functions

Outline

- 1 A Catalog of Essential Functions
 - Linear Functions
 - Polynomials
 - Power Functions
 - Rational Functions
 - Algebraic Functions
 - Transcendental Functions
- 2 New Functions from Old Functions

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
`https://creativecommons.org/licenses/by/3.0/us/`
and the links therein.


Linear Functions

Definition (Linear Function)


A linear function is a function the graph of which is a line. We can write any linear function in slope-intercept form:

$$f(x) = mx + b.$$

m is called the slope, and b is called the y -intercept.






$f(x)$	Direction	y-intercept
$x + 1$ $-0.5x$ -1		






$f(x)$	Direction	y-intercept
$x + 1$ $-0.5x$ -1	↗	

- $m > 0$ means the graph of f points up (↗).




$f(x)$	Direction	y-intercept
$x + 1$ $-0.5x$ -1	 	

- $m > 0$ means the graph of f points up ().
- $m < 0$ means the graph of f points down (.




$f(x)$	Direction	y-intercept
$x + 1$	\nearrow	
$-0.5x$	\searrow	
-1	\rightarrow	

- $m > 0$ means the graph of f points up (\nearrow).
- $m < 0$ means the graph of f points down (\searrow).
- $m = 0$ means the graph of f is horizontal (\rightarrow).




$f(x)$	Direction	y-intercept
$x + 1$	\nearrow	1
$-0.5x$	\searrow	
-1	\rightarrow	

- $m > 0$ means the graph of f points up (\nearrow).
- $m < 0$ means the graph of f points down (\searrow).
- $m = 0$ means the graph of f is horizontal (\rightarrow).
- b tells us the height of the point where the graph hits the y-axis.



$f(x)$	Direction	y-intercept
$x + 1$	\nearrow	1
$-0.5x + 0$	\searrow	0
-1	\rightarrow	

- $m > 0$ means the graph of f points up (\nearrow).
- $m < 0$ means the graph of f points down (\searrow).
- $m = 0$ means the graph of f is horizontal (\rightarrow).
- b tells us the height of the point where the graph hits the y-axis.



$f(x)$	Direction	y-intercept
$x + 1$	\nearrow	1
$-0.5x$	\searrow	0
-1	\rightarrow	-1

- $m > 0$ means the graph of f points up (\nearrow).
- $m < 0$ means the graph of f points down (\searrow).
- $m = 0$ means the graph of f is horizontal (\rightarrow).
- b tells us the height of the point where the graph hits the y-axis.

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

If we interpret x as an indeterminate formal expression, rather than a number, we say that $f(x)$ is a polynomial (rather than a polynomial function).

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$					
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$?				
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes				
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	?			
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4			
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	?		
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1		
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	?	
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	?
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	?				
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes				
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	?			
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0			
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	?		
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6		
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	?	
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	?
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$?				
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative **integer** and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$?				
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes				
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	?			
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2			
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	?		
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$		
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$?	
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$?
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	3
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

Polynomials

Definition (Polynomial Function)

A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a non-negative integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .

$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	3
$3x^2 - \frac{1}{2x} + \sqrt{2}$?				

Polynomials

Definition (Polynomial Function)


A polynomial function is a function f of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where n is a **non-negative** integer and a_0, \dots, a_n are real numbers, called the coefficients. If $a_n \neq 0$ the integer n is called the degree of f .


$f(x)$	Polynomial?	Degree	a_0	a_1	a_2
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	3
$3x^2 - \frac{1}{2x} + \sqrt{2}$	No				

- Linear functions are polynomial (functions).




Linear

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.




Quadratic

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.
- And there are many more.




Cubic

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.
- And there are many more.



Quartic

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.
- And there are many more.



Quintic

Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = x^a .$$

Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = x^a .$$

x = base.

Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = x^a .$$

x = base. a = **exponent** or **power**.

Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer $(1, 2, 3, \dots)$

then x^a = polynomial function.

$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$

Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)

then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)

then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = ?$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)

then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = ?$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = x^b y^b$$

$$x^{a+b} =$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = x^b y^b$$

$$x^{a+b} = ?$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)

then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = x^b y^b$$

$$x^{a+b} = x^a x^b$$

$$x^{-a} =$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = x^b y^b$$

$$x^{a+b} = x^a x^b$$

$$x^{-a} = ?$$



Power Functions

Definition (Power Function)

Let $x > 0$, a - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

x = base. a = exponent or power. First equality = one of ways to define for non-integer a (we study $\ln x$, e^x later).

If a - positive integer ($1, 2, 3, \dots$)
then x^a = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$ when n -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = x^b y^b$$

$$x^{a+b} = x^a x^b$$

$$x^{-a} = \frac{1}{x^a}$$




- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.

- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.


- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.
- Let $x > 0$. For $n = 2m + 1$ -odd, we can extend the definition of n^{th} root to negative numbers by $\sqrt[n]{-x} := -\sqrt[n]{x}$.

- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.
- Let $x > 0$. For $n = 2m + 1$ -odd, we can extend the definition of n^{th} root to negative numbers by $\sqrt[n]{-x} := -\sqrt[n]{x}$.
- In this course, even roots of negative numbers are not defined.



- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.
- Let $x > 0$. For $n = 2m + 1$ -odd, we can extend the definition of n^{th} root to negative numbers by $\sqrt[n]{-x} := -\sqrt[n]{x}$.
- In this course, even roots of negative numbers are not defined.
- The graph of \sqrt{x} is the top half of the parabola $x = y^2$.





- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.
- Let $x > 0$. For $n = 2m + 1$ -odd, we can extend the definition of n^{th} root to negative numbers by $\sqrt[n]{-x} := -\sqrt[n]{x}$.
- In this course, even roots of negative numbers are not defined.
- The graph of \sqrt{x} is the top half of the parabola $x = y^2$. Similarly for $y = \sqrt[2m]{x}$, we graph top of $x = y^{2m}$.




- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.
- Let $x > 0$. For $n = 2m + 1$ -odd, we can extend the definition of n^{th} root to negative numbers by $^{2m+1}\sqrt{-x} := -^{2m+1}\sqrt{x}$.
- In this course, even roots of negative numbers are not defined.
- The graph of \sqrt{x} is the top half of the parabola $x = y^2$. Similarly for $y = ^{2m}\sqrt{x}$, we graph top of $x = y^{2m}$.
- The graph of the cube root $f(x) = \sqrt[3]{x}$ is the graph of the polynomial $x = y^3$.



- n - positive integer, $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$ the n^{th} root function.
 $\sqrt[n]{x} \geq 0$ for $x \geq 0$.
- For $n = 2$, we get the square root \sqrt{x} ; for $n = 3$ we get the cube root $\sqrt[3]{x}$, and so on.
- Let $x > 0$. For $n = 2m + 1$ -odd, we can extend the definition of n^{th} root to negative numbers by $\sqrt[n]{-x} := -\sqrt[n]{x}$.
- In this course, even roots of negative numbers are not defined.
- The graph of \sqrt{x} is the top half of the parabola $x = y^2$. Similarly for $y = \sqrt[2m]{x}$, we graph top of $x = y^{2m}$.
- The graph of the cube root $f(x) = \sqrt[3]{x}$ is the graph of the polynomial $x = y^3$. Similarly for $y = \sqrt[2m+1]{x}$, we graph $x = y^{2m+1}$.



$f(x) = x^{-1} = \frac{1}{x}$ is called the reciprocal function. Its graph has equation $y = \frac{1}{x}$, or $xy = 1$, and is an hyperbola with the coordinate axes as its



asymptotes.

Rational Functions

Definition (Rational Function)

A rational function is a quotient of two polynomials; that is, a function of the form

$$f(x) = \frac{g(x)}{h(x)},$$

where g and h are polynomials.


Rational Functions

Definition (Rational Function)

A rational function is a quotient of two polynomials; that is, a function of the form

$$f(x) = \frac{g(x)}{h(x)},$$

where g and h are polynomials.



Example ($x/(x^2 - 1)$)

The function

$$f(x) = \frac{x}{x^2 - 1}$$

is a rational function.

Algebraic Functions

(Algebraic Function)

A function in x that can be constructed using x , constants, and finitely many of the operations $+$, $-$, $$, $/$, and $\sqrt[n]{}$ is an algebraic function.*

Algebraic Functions

(Algebraic Function)

A function in x that can be constructed using x , constants, and finitely many of the operations $+$, $-$, $$, $/$, and $\sqrt[n]{}$ is an algebraic function.*

Outside of present course: function $f(x)$ = algebraic if it satisfies a polynomial equation with polynomial coefficients, i.e., $a_0(x) + a_1(x)f(x) + \cdots + a_n(x)(f(x))^n = 0$ for some polynomials $a_i(x)$.




Algebraic Functions

(Algebraic Function)

A function in x that can be constructed using x , constants, and finitely many of the operations $+$, $-$, $$, $/$, and $\sqrt[n]{}$ is an algebraic function.*

Outside of present course: function $f(x)$ = algebraic if it satisfies a polynomial equation with polynomial coefficients, i.e., $a_0(x) + a_1(x)f(x) + \cdots + a_n(x)(f(x))^n = 0$ for some polynomials $a_i(x)$.

Examples.



Transcendental Functions

Transcendental functions include many classes of functions.

Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as $\cos x$, $\sin x$, $\tan x$, etc.

Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as $\cos x$, $\sin x$, $\tan x$, etc.
- Exponential functions such as 2^x , $\left(\frac{1}{2}\right)^x$, 5^x , e^x , etc.

Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as $\cos x$, $\sin x$, $\tan x$, etc.
- Exponential functions such as 2^x , $\left(\frac{1}{2}\right)^x$, 5^x , e^x , etc.
- The logarithm function $\ln x$.

Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as $\cos x$, $\sin x$, $\tan x$, etc.
- Exponential functions such as 2^x , $\left(\frac{1}{2}\right)^x$, 5^x , e^x , etc.
- The logarithm function $\ln x$.
- And many more.

Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as $\cos x$, $\sin x$, $\tan x$, etc.
- Exponential functions such as 2^x , $\left(\frac{1}{2}\right)^x$, 5^x , e^x , etc.
- The logarithm function $\ln x$.
- And many more.
- Outside of the present course: by definition, a function is transcendental if it is not algebraic, i.e., if it satisfies no polynomial equation with polynomial coefficients.

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) =$$

$$(f - g)(x) =$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) =$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = ?$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = ?$$

$$\left(\frac{f}{g}\right)(x) =$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) =$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

for $g(x) \neq 0$.

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

for $g(x) \neq 0$.

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f .

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f .

$$\text{Dom}(f + g) = ?$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others.

$$\text{Dom}(f + g) = ?$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\text{Dom}(f + g) = ?$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= ? \\ \text{Dom}(f \cdot g) &= ? \\ \text{Dom}\left(\frac{f}{g}\right) &= ? \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\text{Dom}(f + g) = \text{Dom}(f) \cap \text{Dom}(g)$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

\cap stands for
set intersection

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= ? \\ \text{Dom}(f \cdot g) &= ? \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \mid g(x) \neq 0\}$$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad \Bigg| \quad \text{for } g(x) \neq 0$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\text{Dom}(f + g) = \text{Dom}(f) \cap \text{Dom}(g)$$

$$\text{Dom}(f - g) = \text{Dom}(f) \cap \text{Dom}(g)$$

$$\text{Dom}(f \cdot g) = \text{Dom}(f) \cap \text{Dom}(g)$$

$$\text{Dom}\left(\frac{f}{g}\right) = \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \mid g(x) \neq 0\}$$

\cap stands for
set intersection

right expr.
stands for set
where $g(x) \neq 0$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \mid g(x) \neq 0\} \end{aligned}$$

\cap stands for set intersection

 right expr. stands for **set** where $g(x) \neq 0$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \text{ — } g(x) \neq 0\} \end{aligned}$$

\cap stands for
set intersection

 right expr.
stands for set
where $g(x) \neq 0$

Combinations of Functions

Two functions f and g can be combined to form new functions $f + g$, $f - g$, $f \cdot g$, and $\frac{f}{g}$:

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let $\text{Dom}(f)$ denote the domain of f . The function $f + g$ is defined only if both f and g are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \mid g(x) \neq 0\} \end{aligned}$$

\cap stands for set intersection


 right expr. stands for set where $g(x) \neq 0$

Definition (Composition of f and g)

If f and g are two functions, then the composition of f and g is written $f \circ g$ and is defined by the formula

$$(f \circ g)(x) = f(g(x)).$$

Imagine f and g as machines taking some input and producing some output. Then $f \circ g$ corresponds to attaching both machines end-to-end so that the output of g becomes the input of f .




Definition (Composition of f and g)

If f and g are two functions, then the composition of f and g is written $f \circ g$ and is defined by the formula

$$(f \circ g)(x) = f(g(x)).$$

Imagine f and g as machines taking some input and producing some output. Then $f \circ g$ corresponds to attaching both machines end-to-end so that the output of g becomes the input of f .



The domain of $f \circ g$ is the set of all numbers x in the domain of g such that $g(x)$ is in the domain of f . If the domain of f is A and the domain of g is B , we write this as

$$\{x \in B \mid g(x) \in A\}.$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x)$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x))$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x})$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in \mathbb{R}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x)$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x))$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x})$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = \sqrt{3-\sqrt{x}}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3-\sqrt{x} \geq 0$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3 - \sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3 - \sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3-\sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3-\sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

$$x \leq 9$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3-\sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

$$x \leq 9$$

$$x \in ?$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3-\sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

$$x \leq 9$$

$$x \in [0, 9]$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x)$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x))$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x})$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

Example

Find $f \circ g, g \circ f, g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

Example

Find $f \circ g, g \circ f, g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

Example

Find $f \circ g$, $g \circ f$, $g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

$$x \geq -6$$

Example

Find $f \circ g, g \circ f, g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

$$x \geq -6$$

$$x \in ?$$

Example

Find $f \circ g, g \circ f, g \circ g$ and their domains, where $f(x) = \sqrt{x}$ and $g(x) = \sqrt{3-x}$.

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

$$x \geq -6$$

$$x \in [-6, 3].$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$x \neq ?$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$\begin{aligned} f(x) &= \frac{2x-1}{x+2} \\ g(x) &= \frac{2x+3}{5x-7} \end{aligned} \quad \left| \quad x \neq -2 \right.$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq ?$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x))$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right)$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right) = \frac{2\left(\frac{2x + 3}{5x - 7}\right) - 1}{\frac{2x + 3}{5x - 7} + 2}$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right) = \frac{2\left(\frac{2x + 3}{5x - 7}\right) - 1}{\frac{2x + 3}{5x - 7} + 2}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} \end{aligned}$$

Example

Give simplified formulas for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{\cancel{5x-7}}}{\frac{2x+3+(10x-14)}{\cancel{5x-7}}} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq ?$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x))$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right)$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2} \quad \left| \quad x \neq -2 \right.$$

$$g(x) = \frac{2x+3}{5x-7} \quad \left| \quad x \neq \frac{7}{5} \right.$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \quad \left| \quad x \neq \frac{11}{12}, \frac{7}{5} \right. \end{aligned}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$x \neq -2$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right) = \frac{2\left(\frac{2x + 3}{5x - 7}\right) - 1}{\frac{2x + 3}{5x - 7} + 2}$$

$$= \frac{\frac{2(2x + 3)}{5x - 7} - \frac{5x - 7}{5x - 7}}{\frac{2x + 3}{5x - 7} + \frac{2(5x - 7)}{5x - 7}} = \frac{\frac{4x + 6 - (5x - 7)}{5x - 7}}{\frac{2x + 3 + (10x - 14)}{5x - 7}} = \frac{-x + 13}{12x - 11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x - 1}{x + 2}\right) = \frac{2\left(\frac{2x - 1}{x + 2}\right) - 1}{\frac{2x - 1}{x + 2} + 2}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

=?

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq ?$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

$$(g \circ f)(x) = ?$$

$$x \neq ?$$

$$(g \circ g)(x) = ?$$

$$x \neq ?$$

Example

Give simplified f-las for $f \circ g$, $f \circ f$, $g \circ f$, $g \circ g$. Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

$$(g \circ f)(x) = \frac{7x+4}{3x-19}$$

$$x \neq -2, \frac{19}{3}$$

$$(g \circ g)(x) = \frac{19x-15}{-25x+64}$$

$$x \neq \frac{7}{5}, \frac{64}{25}$$