

# Calculus III

## Lecture 20

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Divergence Theorem

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
`https://creativecommons.org/licenses/by/3.0/us/`  
and the links therein.

## Theorem (Divergence Theorem)

Let  $D$  be a compact set in space with boundary  $S$  a piecewise smooth parametrized surface, oriented by the outward normal, and let  $\mathbf{X}$  be a smooth vector field on  $D$  given by

$$\mathbf{X}(x, y, z) = P(x, y, z)\mathbf{i} + Q(x, y, z)\mathbf{j} + R(x, y, z)\mathbf{k} \quad .$$

Then

$$\iint_S \mathbf{X} \cdot d\mathbf{S} = \iiint_D \left( \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dV$$

## Corollary (May serve as alternative definition of div)

$$\begin{aligned} (\operatorname{div} \mathbf{X})(p) &= \lim_{D \rightarrow \{p\}} \frac{1}{\operatorname{vol}(D)} \iint_S \mathbf{X} \cdot d\mathbf{S} \\ &= \lim_{D \rightarrow \{p\}} \frac{1}{\operatorname{vol}(D)} \iiint_D \left( \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} \right) dV \\ &= \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z}. \end{aligned}$$

# Divergence Theorem

- Let  $\mathbf{X} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ .
- Recall our notation


$$\begin{aligned}\operatorname{div} \mathbf{X} &= \frac{\partial P}{\partial x} + \frac{\partial Q}{\partial y} + \frac{\partial R}{\partial z} = (\partial_x, \partial_y, \partial_z) \cdot (P, Q, R) \\ \operatorname{div} \mathbf{X} &= \nabla \cdot \mathbf{X}.\end{aligned}$$

## Theorem (Divergence Theorem)

$$\iint_S \mathbf{X} \cdot d\mathbf{S} = \iiint_D \operatorname{div} \mathbf{X} dV$$

- If  $(\operatorname{div} \mathbf{X})(p) > 0$ , then  $p$  acts as a source;
- If  $(\operatorname{div} \mathbf{X})(p) < 0$ , then  $p$  acts as a sink;
- If  $\operatorname{div} \mathbf{X} \equiv 0$  on some domain  $D$ , then  $\mathbf{X}$  is incompressible on  $D$ .

# Example



Let  $S$  be the part of the paraboloid  $z = 4 - x^2 - y^2$  above the  $xy$ -plane, oriented upward, and  $\mathbf{X} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$ . Use the Divergence Theorem to compute  $\iint_S \mathbf{X} \cdot d\mathbf{S}$ .

The surface  $S$  does not enclose a region in space. However, we add the disk  $D$  of radius 2 centered at the origin in the plane  $z = 0$  to make it closed.  $R$  orients  $D$  with the downward normal, hence

$$\begin{aligned} \iint_{S \uparrow \cup D \downarrow} \mathbf{X} \cdot d\mathbf{S} &= \iiint_R \operatorname{div} \mathbf{X} \, dV = 0, \\ \iint_{S \uparrow} \mathbf{X} \cdot d\mathbf{S} &= \iint_{D \uparrow} \mathbf{X} \cdot d\mathbf{S}. \end{aligned}$$

The upward normal to  $D$  is  $\mathbf{k}$ , hence  $\mathbf{X} \cdot d\mathbf{S} = \mathbf{X} \cdot \mathbf{k} \, dS = c \, dS$ . Therefore

$$\iint_S \mathbf{X} \cdot d\mathbf{S} = \iint_D \mathbf{X} \cdot d\mathbf{S} = \iint_D c \, dS = c \cdot \operatorname{area}(D) = 4\pi c.$$

# Balloon Pressure Equilibrium

- Let  $\mathbf{F}$  be the total displacement force due pressure difference between interior and exterior of inflated balloon:

$$\mathbf{F} = \iint_S d\mathbf{F} = \iint_S p\mathbf{N} dS = \iint_S p d\mathbf{S} .$$

# Balloon Pressure Equilibrium

- Let  $\mathbf{F}$  be the total displacement force due pressure difference between interior and exterior of inflated balloon:

$$\mathbf{F} = \iint_S d\mathbf{F} = \iint_S p\mathbf{N} dS = \iint_S p d\mathbf{S}.$$

- For every unit vector  $\mathbf{u}$  we have

$$\mathbf{F} \cdot \mathbf{u} = \left( \iint_S p\mathbf{N} dS \right) \cdot \mathbf{u} = \iint_S p\mathbf{u} \cdot \mathbf{N} dS = \iiint_D \operatorname{div}(p\mathbf{u}) dV = 0$$

because  $\operatorname{div}(p\mathbf{u}) = 0$  since the vector field  $\mathbf{X} = p\mathbf{u}$  is constant on  $D$ .

- Therefore  $\mathbf{F} \cdot \mathbf{u} = 0$  for every unit vector  $\mathbf{u}$ ;



# Balloon Pressure Equilibrium

- Let  $\mathbf{F}$  be the total displacement force due pressure difference between interior and exterior of inflated balloon:

$$\mathbf{F} = \iint_S d\mathbf{F} = \iint_S p\mathbf{N} dS = \iint_S p d\mathbf{S}.$$

- For every unit vector  $\mathbf{u}$  we have

$$\mathbf{F} \cdot \mathbf{u} = \left( \iint_S p\mathbf{N} dS \right) \cdot \mathbf{u} = \iint_S p\mathbf{u} \cdot \mathbf{N} dS = \iiint_D \operatorname{div}(p\mathbf{u}) dV = 0$$

because  $\operatorname{div}(p\mathbf{u}) = 0$  since the vector field  $\mathbf{X} = p\mathbf{u}$  is constant on  $D$ .

- Therefore  $\mathbf{F} \cdot \mathbf{u} = 0$  for every unit vector  $\mathbf{u}$ ;
- Which implies  $\mathbf{F} = \mathbf{0}$ .

# Archimedes' Law from the Divergence Theorem

A solid body is submerged into a tank containing a liquid of constant density  $\rho$ . What is the buoyant force?

# Archimedes' Law from the Divergence Theorem

A solid body is submerged into a tank containing a liquid of constant density  $\rho$ . What is the buoyant force?

- Body occupies a region  $D$ , exterior boundary  $S$ ;
- Unit outward normal field  $\mathbf{N}$ ;

# Archimedes' Law from the Divergence Theorem

A solid body is submerged into a tank containing a liquid of constant density  $\rho$ . What is the buoyant force?

- Body occupies a region  $D$ , exterior boundary  $S$ ;
- Unit outward normal field  $\mathbf{N}$ ;
- Magnitude of pressure at depth  $a$  below the surface is  $p_0 + \rho ag$ , where
  - $g$  is the magnitude of the gravitational acceleration.
  - $p_0$  is the pressure at surface of liquid

# Archimedes' Law from the Divergence Theorem

A solid body is submerged into a tank containing a liquid of constant density  $\rho$ . What is the buoyant force?

- Body occupies a region  $D$ , exterior boundary  $S$ ;
- Unit outward normal field  $\mathbf{N}$ ;
- Magnitude of pressure at depth  $a$  below the surface is  $p_0 + \rho ag$ , where
  - $g$  is the magnitude of the gravitational acceleration.
  - $p_0$  is the pressure at surface of liquid
- Infinitesimal force acting on  $S$  is  $d\mathbf{F} = -(p_0 + \rho ag) \mathbf{N} dS$ ,

# Archimedes' Law from the Divergence Theorem

A solid body is submerged into a tank containing a liquid of constant density  $\rho$ . What is the buoyant force?

- Body occupies a region  $D$ , exterior boundary  $S$ ;
- Unit outward normal field  $\mathbf{N}$ ;
- Magnitude of pressure at depth  $a$  below the surface is  $p_0 + \rho ag$ , where
  - $g$  is the magnitude of the gravitational acceleration.
  - $p_0$  is the pressure at surface of liquid
- Infinitesimal force acting on  $S$  is  $d\mathbf{F} = -(p_0 + \rho ag) \mathbf{N} dS$ ,
- The total force is

$$\mathbf{F} = \iint_S d\mathbf{F} = \iint_S -(p_0 + \rho ag) \mathbf{N} dS = \iint_S -\rho ag \mathbf{N} dS = \iint_S \rho g z \mathbf{N} dS$$

# Archimedes' Law from the Divergence Theorem

A solid body is submerged into a tank containing a liquid of constant density  $\rho$ . What is the buoyant force?

- Body occupies a region  $D$ , exterior boundary  $S$ ;
- Unit outward normal field  $\mathbf{N}$ ;
- Magnitude of pressure at depth  $a$  below the surface is  $p_0 + \rho ag$ , where
  - $g$  is the magnitude of the gravitational acceleration.
  - $p_0$  is the pressure at surface of liquid
- Infinitesimal force acting on  $S$  is  $d\mathbf{F} = -(p_0 + \rho ag) \mathbf{N} dS$ ,
- The total force is

$$\mathbf{F} = \iint_S d\mathbf{F} = \iint_S -(p_0 + \rho ag) \mathbf{N} dS = \iint_S -\rho ag \mathbf{N} dS = \iint_S \rho gz \mathbf{N} dS$$

- EC: Use the Divergence Theorem to show that  $\mathbf{F} = \rho Vg \mathbf{k}$  ( $V$ : volume of the region enclosed by  $S$ .)

# Curl

Let  $\mathbf{X} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$  be a smooth vector field.

## Definition (Curl, coordinate definition)

The *curl* of a vector field  $\mathbf{X}$ , denoted by  $\mathbf{curl} \mathbf{X}$ , is defined by  

$$\mathbf{curl} \mathbf{X} = (\partial_y R - \partial_z Q)\mathbf{i} + (\partial_z P - \partial_x R)\mathbf{j} + (\partial_x Q - \partial_y P)\mathbf{k}.$$

$$\mathbf{curl} \mathbf{X} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ \partial_x & \partial_y & \partial_z \\ P & Q & R \end{vmatrix} = \nabla \times \mathbf{X}.$$

- Just like  $\text{div}$ ,  $\mathbf{curl}$  can be equipped with a coordinate-free definition (in this case the above definition becomes a theorem).



# Induced Orientation on a Boundary Curve


- Let  $S$  be smooth surface, oriented by unit normal vector  $\mathbf{n}$ .
- Let  $D$  be region in  $S$ , bounded by a curve  $C = \partial D$ .
- Let  $\mathbf{N}$  denote the unit vector field on  $C$  which is
  - tangent to  $S$ ;
  - normal to  $C$ ;
  - pointing outward of  $D$ .
- Let  $\mathbf{T}$  be unit tangent vector to  $C$  (and hence tangent to  $S$ ).
- Then  $\mathbf{N}$  orients the tangents of  $C$  and thus  $C$  itself.

## Definition

We say that  $\mathbf{T}$  is *positively oriented* if the triple  $\{\mathbf{n}, \mathbf{N}, \mathbf{T}\}$  is positively oriented in space.


- Since  $\mathbf{T}, \mathbf{n}, \mathbf{N}$  are pairwise orthogonal unit vectors, positive orientation is equivalent to  $\mathbf{T} = \mathbf{n} \times \mathbf{N}$ .
- If we view the plane tangent to  $S$  from the tip of  $\mathbf{n}$ , then  $\{\mathbf{N}, \mathbf{T}\}$  is positively oriented in that plane.

## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals ?


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ ,


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , **pointing outwards from  $D \Rightarrow \mathbf{N}$  equals ?**

## Example (Orientation of the equator of a sphere)




Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .




## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N}$


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = \mathbf{i} \times (-\mathbf{k}) = ?$


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = \mathbf{i} \times (-\mathbf{k}) = \mathbf{j}$ .


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = \mathbf{i} \times (-\mathbf{k}) = \mathbf{j}$ .


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = \mathbf{i} \times (-\mathbf{k}) = \mathbf{j}$ .
- A viewer, standing along  $\mathbf{n}$  with feet on surface, and facing in the direction of the tangent, has the surface to the left.


## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = \mathbf{i} \times (-\mathbf{k}) = \mathbf{j}$ .
- A viewer, standing along  $\mathbf{n}$  with feet on surface, and facing in the direction of the tangent, has the surface to the left.
- Change  $D$  to be lower hemisphere: we get  $\mathbf{N} = \mathbf{k}$ ,  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = -\mathbf{j}$ .

## Example (Orientation of the equator of a sphere)



Let  $S$  be the unit sphere  $x^2 + y^2 + z^2 = 1$  oriented by the outward normal  $\mathbf{n}$ ,  $D = S \cap \{z \geq 0\}$  be the upper hemisphere. Introduce an orientation on the boundary  $C = \partial D$ .

- At the point  $(1, 0, 0)$  the normal to the surface  $\mathbf{n}$  equals  $\mathbf{i}$ .
- Let  $\mathbf{T}$  be a unit tangent to  $C$  at  $(1, 0, 0)$ ; then  $\mathbf{T} = \mathbf{j}$  or  $-\mathbf{j}$ .
- Let  $\mathbf{N}$  be unit vector perpendicular to  $\mathbf{n}$  and  $\mathbf{T}$ , pointing outwards from  $D \Rightarrow \mathbf{N}$  equals  $-\mathbf{k}$ .
- $\Rightarrow$  positively oriented tangent to  $C$  is  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = \mathbf{i} \times (-\mathbf{k}) = \mathbf{j}$ .
- A viewer, standing along  $\mathbf{n}$  with feet on surface, and facing in the direction of the tangent, has the surface to the left.
- Change  $D$  to be lower hemisphere: we get  $\mathbf{N} = \mathbf{k}$ ,  $\mathbf{T} = \mathbf{n} \times \mathbf{N} = -\mathbf{j}$ .
- A viewer, standing along  $\mathbf{n}$  with feet on surface, facing in the direction of the tangent, has the surface again to the left.

- Let  $S$  be a smooth surface, oriented by the unit normal field  $\mathbf{n}$ .
- Let  $D$  be a region on  $S$ , bounded by the piecewise smooth curve  $C = \partial D$ .
- Let  $C$  have unit tangent  $\mathbf{T}$  positively oriented by  $\mathbf{n}$ .
- Let  $\mathbf{X} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$  be a smooth vector field defined in a open set around  $S$ .
- Recall that  $\mathbf{curl} \, \mathbf{X} = (R_y - Q_z)\mathbf{i} + (P_z - R_x)\mathbf{j} + (Q_x - P_y)\mathbf{k}$ .
- Recall that  $\mathbf{X} \cdot d\mathbf{r} = \mathbf{T}ds$  and  $d\mathbf{S} = \mathbf{n}dS$ .

### Theorem (Stokes)

$$\oint_C \mathbf{X} \cdot d\mathbf{r} = \iint_D \mathbf{curl} \, \mathbf{X} \cdot d\mathbf{S} \quad .$$



## Theorem (Stokes)

$$\oint_C \mathbf{X} \cdot d\mathbf{r} = \iint_D \mathbf{curl} \mathbf{X} \cdot d\mathbf{S} \quad .$$

Idea of proof:

- Use a parametrization of  $S$  to get integrals in the parameter plane.
- Apply Green's Theorem in the parameter plane.

We can use Stokes' theorem to:

- Evaluate line integrals by computing a surface integral, or
- Evaluate a surface integral by computing a line integral.

# Example

# Vector Potential

Given a smooth vector field  $\mathbf{X}$ , one can ask:

- Is  $\mathbf{X}$  the **curl** of a vector field?
- Any field  $\mathbf{G}$  such that  $\mathbf{X} = \mathbf{curl} \mathbf{G}$  is called a *vector potential* for  $\mathbf{X}$ .
- If  $\mathbf{X} = \nabla \times \mathbf{G}$  is a curl field, then  $\operatorname{div} \mathbf{X} = 0$ .
- Two vector potentials differ by a gradient field.

Surface  $D$  the part of the paraboloid  $z = 4 - x^2 - y^2$  above the  $xy$ -plane, oriented upward,  $\mathbf{X} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$

$$\iint_D \mathbf{X} \cdot d\mathbf{S}$$

Surface  $D$  the part of the paraboloid  $z = 4 - x^2 - y^2$  above the  $xy$ -plane, oriented upward,  $\mathbf{X} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$

$$\iint_D \mathbf{X} \cdot d\mathbf{S}$$

$\operatorname{div} \mathbf{X} = 0$ , hence  $\mathbf{X}$  may be the curl of a vector field  $\mathbf{G} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ .

Surface  $D$  the part of the paraboloid  $z = 4 - x^2 - y^2$  above the  $xy$ -plane, oriented upward,  $\mathbf{X} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$

$$\iint_D \mathbf{X} \cdot d\mathbf{S}$$

$\operatorname{div} \mathbf{X} = 0$ , hence  $\mathbf{X}$  may be the curl of a vector field  $\mathbf{G} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ .

$$Q_x - P_y = c, \quad P_z - R_x = b, \quad R_y - Q_z = a.$$

Surface  $D$  the part of the paraboloid  $z = 4 - x^2 - y^2$  above the  $xy$ -plane, oriented upward,  $\mathbf{X} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$

$$\iint_D \mathbf{X} \cdot d\mathbf{S}$$

$\operatorname{div} \mathbf{X} = 0$ , hence  $\mathbf{X}$  may be the curl of a vector field  $\mathbf{G} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ .

$$Q_x - P_y = c, \quad P_z - R_x = b, \quad R_y - Q_z = a.$$

One solution is  $Q = cx$ ,  $P = bz$ ,  $R = ay$ , hence  $\mathbf{G} = bz\mathbf{i} + cx\mathbf{j} + ay\mathbf{k}$  is a vector potential for  $\mathbf{X}$ . Then  $\mathbf{X} = \mathbf{curl} \mathbf{G}$  and therefore

$$\iint_D \mathbf{X} \cdot d\mathbf{S} = \iint_D \mathbf{curl} \mathbf{G} \cdot d\mathbf{S} = \oint_C \mathbf{G} \cdot d\mathbf{r} = \oint_C bz \, dx + cx \, dy + ay \, dz,$$

where  $C = \partial D$ , the circle of radius 2 centered at the origin, oriented counterclockwise;



Surface  $D$  the part of the paraboloid  $z = 4 - x^2 - y^2$  above the  $xy$ -plane, oriented upward,  $\mathbf{X} = a\mathbf{i} + b\mathbf{j} + c\mathbf{k}$

$$\iint_D \mathbf{X} \cdot d\mathbf{S}$$

$\operatorname{div} \mathbf{X} = 0$ , hence  $\mathbf{X}$  may be the curl of a vector field  $\mathbf{G} = P\mathbf{i} + Q\mathbf{j} + R\mathbf{k}$ .

$$Q_x - P_y = c, \quad P_z - R_x = b, \quad R_y - Q_z = a.$$

One solution is  $Q = cx$ ,  $P = bz$ ,  $R = ay$ , hence  $\mathbf{G} = bz\mathbf{i} + cx\mathbf{j} + ay\mathbf{k}$  is a vector potential for  $\mathbf{X}$ . Then  $\mathbf{X} = \operatorname{curl} \mathbf{G}$  and therefore

$$\iint_D \mathbf{X} \cdot d\mathbf{S} = \iint_D \operatorname{curl} \mathbf{G} \cdot d\mathbf{S} = \oint_C \mathbf{G} \cdot d\mathbf{r} = \oint_C bz \, dx + cx \, dy + ay \, dz,$$

where  $C = \partial D$ , the circle of radius 2 centered at the origin, oriented counterclockwise;  $x = 2 \cos t$ ,  $y = 2 \sin t$ ,  $z = 0$ , with  $0 \leq t \leq 2\pi$  is an orientation-compatible parametrization of  $C$

$$\oint_C \mathbf{G} \cdot d\mathbf{r} = \int_0^{2\pi} 2c \cos t (2 \cos t) \, dt = 4c \int_0^{2\pi} \cos^2 t \, dt = 4\pi c.$$

# Div, Curl, Grad

$$\operatorname{div}(\mathbf{curl} \mathbf{X}) = \nabla \cdot (\nabla \times \mathbf{X}) = 0 .$$

# Div, Curl, Grad

$$\operatorname{div}(\mathbf{curl} \mathbf{X}) = \nabla \cdot (\nabla \times \mathbf{X}) = 0 .$$

$B$ : ball centered at  $p$ , with boundary a sphere  $S$  centered at  $p$ .

$$\iiint_B \operatorname{div}(\mathbf{curl} \mathbf{X}) dV = \iint_{S=\partial B} \mathbf{curl} \mathbf{X} \cdot d\mathbf{S} = \oint_{\partial S} \mathbf{X} \cdot d\mathbf{r} = 0 ,$$

$$\operatorname{div}(\mathbf{curl} \mathbf{X})(p) = \lim_{B \rightarrow \{p\}} \frac{1}{\operatorname{vol}(B)} \iiint_B \operatorname{div}(\mathbf{curl} \mathbf{X}) dV = 0 .$$

# Div, Curl, Grad

$$\operatorname{div}(\mathbf{curl} \mathbf{X}) = \nabla \cdot (\nabla \times \mathbf{X}) = 0 .$$

$B$ : ball centered at  $p$ , with boundary a sphere  $S$  centered at  $p$ .

$$\iiint_B \operatorname{div}(\mathbf{curl} \mathbf{X}) dV = \iint_{S=\partial B} \mathbf{curl} \mathbf{X} \cdot d\mathbf{S} = \oint_{\partial S} \mathbf{X} \cdot d\mathbf{r} = 0 ,$$

$$\operatorname{div}(\mathbf{curl} \mathbf{X})(p) = \lim_{B \rightarrow \{p\}} \frac{1}{\operatorname{vol}(B)} \iiint_B \operatorname{div}(\mathbf{curl} \mathbf{X}) dV = 0 .$$

$$\mathbf{curl}(\mathbf{grad} f) = \nabla \times (\nabla f) = \mathbf{0} .$$

# Div, Curl, Grad

$$\operatorname{div}(\mathbf{curl} \mathbf{X}) = \nabla \cdot (\nabla \times \mathbf{X}) = 0 .$$

$B$ : ball centered at  $p$ , with boundary a sphere  $S$  centered at  $p$ .

$$\iiint_B \operatorname{div}(\mathbf{curl} \mathbf{X}) dV = \iint_{S=\partial B} \mathbf{curl} \mathbf{X} \cdot d\mathbf{S} = \oint_{\partial S} \mathbf{X} \cdot d\mathbf{r} = 0 ,$$

$$\operatorname{div}(\mathbf{curl} \mathbf{X})(p) = \lim_{B \rightarrow \{p\}} \frac{1}{\operatorname{vol}(B)} \iiint_B \operatorname{div}(\mathbf{curl} \mathbf{X}) dV = 0 .$$

$$\mathbf{curl}(\mathbf{grad} f) = \nabla \times (\nabla f) = \mathbf{0} .$$

$D$ : disk centered at  $p$ , in the plane normal to  $\mathbf{n}$  at  $p$ , and  $C = \partial D$

$$\iint_D \mathbf{curl}(\mathbf{grad} f) \cdot \mathbf{n} dS = \iint_D \mathbf{curl}(\mathbf{grad} f) \cdot d\mathbf{S} = \oint_C \mathbf{grad} f \cdot d\mathbf{r} = 0 ,$$

$$\mathbf{curl}(\mathbf{grad} f)(p) \cdot \mathbf{n} = \lim_{D \rightarrow \{p\}} \frac{1}{\operatorname{area}(D)} \iint_D \mathbf{curl}(\mathbf{grad} f) \cdot \mathbf{n} dS = 0 ;$$

since this is valid for all unit vectors  $\mathbf{n}$ , we conclude that