Calculus I Lecture 24 Area between Graphs

Todor Miley

https://github.com/tmilev/freecalc

2020

Lecture 24

Outline

Integration and symmetry

Outline

Integration and symmetry

More About Areas

License to use and redistribute

These lecture slides and their LATEX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share to copy, distribute and transmit the work,
- to Remix to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides: https://github.com/tmilev/freecalc
- Should the link be outdated/moved, search for "freecalc project".
- Creative Commons license CC BY 3.0:
 https://creativecommons.org/licenses/by/3.0/us/and the links therein.

License to use and redistribute

These lecture slides and their LATEX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share to copy, distribute and transmit the work,
- to Remix to adapt, change, etc., the work,
- to make commercial use of the work.

as long as you reasonably acknowledge the original project.


- Latest version of the .tex sources of the slides: https://github.com/tmilev/freecalc
- Should the link be outdated/moved, search for "freecalc project".
- Creative Commons license CC BY 3.0:
 https://creativecommons.org/licenses/by/3.0/us/
 and the links therein.

Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$





Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$






Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$




Todor Milev

Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{x} f(x) dx = 2 \int_{0}^{x} f(x) dx$.
- ② If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$




Todor Milev


Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{x} f(x) dx = 2 \int_{0}^{x} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$






Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$



Todor Milev

Lecture 24


Area between Graphs

Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$




Todor Milev

Theorem (Integrals of Symmetric Functions)

Suppose f is continuous on [-a, a].

- If f is even (that is, f(-x) = f(x)), then $\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$.
- 2 If f is odd (that is, f(-x) = -f(x)), then

$$\int_{-a}^{a} f(x) dx = \int_{0}^{a} f(x) dx + \int_{-a}^{0} f(x) dx = 0.$$




Since
$$f(x) = x^6 + 1$$
 satisfies $f(-x) = f(x)$, it is even, and so
$$\int_{-2}^{2} (x^6 + 1) dx =$$

Since
$$f(x) = x^6 + 1$$
 satisfies $f(-x) = f(x)$, it is even, and so
$$\int_{-2}^{2} (x^6 + 1) dx = 2 \int_{0}^{2} (x^6 + 1) dx$$

Since
$$f(x) = x^6 + 1$$
 satisfies $f(-x) = f(x)$, it is even, and so
$$\int_{-2}^{2} (x^6 + 1) dx = 2 \int_{0}^{2} (x^6 + 1) dx$$
$$= 2 \left[\frac{1}{7} x^7 + x \right]_{0}^{2}$$


Since
$$f(x)=x^6+1$$
 satisfies $f(-x)=f(x)$, it is even, and so
$$\int_{-2}^2 (x^6+1) \, \mathrm{d}x = 2 \int_0^2 (x^6+1) \, \mathrm{d}x$$
$$= 2 \left[\frac{1}{7}x^7+x\right]_0^2$$
$$= 2 \left(\frac{128}{7}+2\right)$$

Since
$$f(x) = x^6 + 1$$
 satisfies $f(-x) = f(x)$, it is even, and so
$$\int_{-2}^{2} (x^6 + 1) \, dx = 2 \int_{0}^{2} (x^6 + 1) \, dx$$
$$= 2 \left[\frac{1}{7} x^7 + x \right]_{0}^{2}$$
$$= 2 \left(\frac{128}{7} + 2 \right)$$
$$= \frac{284}{7}.$$



Since
$$f(x) = \frac{\tan x - x}{1 - 2x^2 + 2x^4}$$
 satisfies $f(-x) = -f(x)$, it is odd, and so

$$\int_{-1}^{1} \frac{\tan x - x}{1 - 2x^2 + 2x^4} \mathrm{d}x =$$




Since
$$f(x) = \frac{\tan x - x}{1 - 2x^2 + 2x^4}$$
 satisfies $f(-x) = -f(x)$, it is odd, and so

$$\int_{-1}^{1} \frac{\tan x - x}{1 - 2x^2 + 2x^4} dx = 0.$$

Lecture 24


More About Areas

Suppose two curves, y = f(x) and y = g(x), are given. How do we find the area bounded by those curves between the endpoints x = a and x = b?




Todor Milev Lecture 24 Area between Graphs 2020

The Area Under a Curve




The Area Between Two Curves




Lecture 24

The Area Under a Curve




 $rectangle \ area = height\cdot width$

The Area Between Two Curves




The Area Under a Curve




rectangle area = height·width

The Area Between Two Curves




The Area Under a Curve




rectangle area = height Δx

The Area Between Two Curves




Lecture 24

The Area Under a Curve




rectangle area = $height \cdot \Delta x$

The Area Between Two Curves




The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$

The Area Between Two Curves




Lecture 24

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles =
$$n = 4$$

A = $\sum_{i=1}^{4} f(x_i) \Delta x$

The Area Between Two Curves



The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = n = 8A = $\sum_{i=1}^{8} f(x_i) \Delta x$

The Area Between Two Curves



The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$



rectangles = n = 16


$$A = \sum_{i=1}^{16} f(x_i) \Delta x$$

The Area Between Two Curves



Lecture 24

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$

$$A = \lim_{n \to \infty} \sum_{i=1}^{n} f(x_i) \Delta x$$

The Area Between Two Curves



The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles =
$$n \rightarrow \infty$$

$$A = \lim_{n \to \infty} \sum_{i=1}^{n} f(x_i) \Delta x$$

The Area Between Two Curves



The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves



Lecture 24

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves




rectangle area = height⋅width

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves




rectangle area = height⋅width

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves




rectangle area = height Δx

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves




rectangle area = $height \cdot \Delta x$

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves




rectangle area = $(f(x) - g(x)) \cdot \Delta x$

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles =
$$n \to \infty$$

$$A = \int_a^b f(x) dx$$

The Area Between Two Curves




rectangle area = $(f(x) - g(x)) \cdot \Delta x$




rectangles = n = 4

$$A = \sum_{i=1}^{4} (f(x_i) - g(x_i)) \Delta x$$

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles = $n \to \infty$ A = $\int_a^b f(x) dx$

The Area Between Two Curves




rectangle area = $(f(x) - g(x)) \cdot \Delta x$




rectangles =
$$n = 8$$

$$A = \sum_{i=1}^{8} (f(x_i) - g(x_i)) \Delta x$$

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles =
$$n \to \infty$$

$$A = \int_a^b f(x) dx$$

The Area Between Two Curves




rectangle area = $(f(x) - g(x)) \cdot \Delta x$




rectangles =
$$n = 16$$

$$A = \sum_{i=1}^{16} (f(x_i) - g(x_i)) \Delta x$$

The Area Under a Curve




rectangle area = $f(x) \cdot \Delta x$




rectangles =
$$n \to \infty$$

A = $\int_a^b f(x) dx$

The Area Between Two Curves



rectangle area = $(f(x) - g(x)) \cdot \Delta x$



rectangles =
$$n \to \infty$$

$$A = \int_a^b [f(x) - g(x)] dx$$

Definition (The Area Between Two Curves)


The area between two curves y = f(x) and y = g(x) bounded by the endpoints x = a and x = b is

$$\int_a^b |f(x)-g(x)| \mathrm{d}x.$$

Note that we use the absolute value, because in general we don't know which curve is above the other.


Example

Find the area of the region bounded above by $y = x^2 + 1$, bounded below by y = x, and bounded on its sides by x = 0 and x = 1.




Todor Milev

Find the area of the region bounded above by $y = x^2 + 1$, bounded below by y = x, and bounded on its sides by x = 0 and x = 1.




Graph the functions.

Find the area of the region bounded above by $y = x^2 + 1$, bounded below by y = x, and bounded on its sides by x = 0 and x = 1.




Graph the functions.

Find the area of the region bounded above by $y = x^2 + 1$, bounded below by y = x, and bounded on its sides by x = 0 and x = 1.




Graph the functions.



- Graph the functions.
- Identify the region.

Example


Find the area of the region bounded above by $y = x^2 + 1$, bounded below by y = x, and bounded on its sides by x = 0 and x = 1.



- Graph the functions.
- Identify the region.


Lecture 24

Example



- Graph the functions.
- Identify the region.
- Integrate.


Example



- Graph the functions.
- Identify the region.
- Integrate.

$$A = \int_0^1 |(x^2 + 1) - x| dx$$
$$= \int_0^1 (x^2 - x + 1) dx$$

Example




- Graph the functions.
- Identify the region.
- Integrate.

$$A = \int_0^1 |(x^2 + 1) - x| dx$$
$$= \int_0^1 (x^2 - x + 1) dx$$
$$= \left[\frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1$$

Example

Find the area of the region bounded above by $y = x^2 + 1$, bounded below by y = x, and bounded on its sides by x = 0 and x = 1.



- Graph the functions.
- Identify the region.
- Integrate.

$$A = \int_0^1 |(x^2 + 1) - x| dx$$


$$= \int_0^1 (x^2 - x + 1) dx$$

$$= \left[\frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1$$

$$= \frac{1}{3} - \frac{1}{2} + 1$$

Lecture 24

Example



- Graph the functions.
- Identify the region.
- Integrate.


$$A = \int_0^1 |(x^2 + 1) - x| dx$$

$$= \int_0^1 (x^2 - x + 1) dx$$


$$= \left[\frac{x^3}{3} - \frac{x^2}{2} + x \right]_0^1$$

$$= \frac{1}{3} - \frac{1}{2} + 1 = \frac{5}{6}.$$

Example



Example




Find the point of

intersection.


Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$.

Example



Find the point of intersection.

$$x^2 = 2x - x^2$$




Find the point of intersection.

$$x^2 = 2x - x^2$$

$$0=2x-2x^2$$

Example




Find the point of intersection.

$$x^{2} = 2x - x^{2}$$

 $0 = 2x - 2x^{2} = 2x(1 - x)$

Example




Find the point of intersection.

$$x^{2} = 2x - x^{2}$$

 $0 = 2x - 2x^{2} = 2x(1 - x)$
 $x = 0$ or 1.

Example




$$0 = 2x - 2x^{2} = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

- Find the point of intersection.
- @ Graph the functions.

Example



Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$. $x^2 = 2x - x^2$

$$C = 2x - x^{-1}$$


$$0 = 2x - 2x^2 = 2x(1-x)$$

$$x = 0 \text{ or } 1.$$

- Find the point of intersection.
- @ Graph the functions.

Lecture 24

Example




Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$. $x^2 = 2x - x^2$

$$0 = 2x - 2x^2 = 2x(1 - x)$$

x = 0 or 1.

- Find the point of intersection.
- @ Graph the functions.

Example




Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$.

$$x^{2} = 2x - x^{2}$$

 $0 = 2x - 2x^{2} = 2x(1 - x)$
 $x = 0$ or 1.

- Find the point of intersection.
- Graph the functions.
- Identify the region.

Example




$$0 = 2x - 2x^2 = 2x(1 - x)$$

x = 0 or 1.

- Find the point of intersection.
- Graph the functions.
- Identify the region.

Example




Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$.

x = 0 or 1.

$$x^{2} = 2x - x^{2}$$
$$0 = 2x - 2x^{2} = 2x(1 - x)$$

- Find the point of intersection.
- Graph the functions.
- Identify the region.
- Integrate.

Example



Find the point of intersection.


- @ Graph the functions.
- Identify the region.
- Integrate.

$$x^2 = 2x - x^2$$

 $0 = 2x - 2x^2 = 2x(1 - x)$
 $x = 0$ or 1.

$$A = \int_0^1 (2x - 2x^2) dx$$

Example




- Find the point of intersection.
- Graph the functions.
- Identify the region.
- Integrate.

$$x^{2} = 2x - x^{2}$$

 $0 = 2x - 2x^{2} = 2x(1 - x)$
 $x = 0 \text{ or } 1$.

$$A = \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx$$

Example



- Find the point of intersection.
- Graph the functions.
- Identify the region.
- Integrate.

Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$. $x^2 = 2x - x^2$


$$0 = 2x - 2x^{2} = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$A = \int_{0}^{1} (2x - 2x^{2}) dx = 2 \int_{0}^{1} (x - x^{2}) dx$$

e point of etion.
$$= 2 \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1$$
 the functions.

Example



- Find the point of intersection.
- @ Graph the functions.
- Identify the region.
- Integrate.


Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$.

$$x^{2} = 2x - x^{2}$$

 $0 = 2x - 2x^{2} = 2x(1 - x)$
 $x = 0 \text{ or } 1.$

$$A = \int_0^1 (2x - 2x^2) dx = 2 \int_0^1 (x - x^2) dx$$
$$= 2 \left[\frac{x^2}{2} - \frac{x^3}{3} \right]_0^1 = 2 \left(\frac{1}{2} - \frac{1}{3} \right)$$

Example



- Find the point of intersection.
- @ Graph the functions.
- Identify the region.
- Integrate.

Find the area of the region enclosed by the parabolas $y = x^2$ and $y = 2x - x^2$.

$$0 = 2x - 2x^{2} = 2x(1 - x)$$

$$x = 0 \text{ or } 1.$$

$$A = \int_{0}^{1} (2x - 2x^{2}) dx = 2 \int_{0}^{1} (x - x^{2}) dx$$

$$=2\left[\frac{x^2}{2}-\frac{x^3}{3}\right]_0^1=2\left(\frac{1}{2}-\frac{1}{3}\right)=\frac{1}{3}.$$

Example



Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

Example



Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.


Find the point of intersection.

Todor Milev

Lecture 24

Area between Graphs

Example




Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

Find the point of intersection.

Example




Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

- Find the point of intersection.
- Graph the functions.

Example




Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

- Find the point of intersection.
- Graph the functions.

Example




Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

- Find the point of intersection.
- Graph the functions.

Example




Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

- Find the point of intersection.
- Graph the functions.
- Identify the region.

Example



Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.


The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

$$A=A_1+A_2$$

- Find the point of intersection.
- Graph the functions.
- Identify the region.

Todor Milev

Example



- Find the point of intersection.
- Graph the functions.
- Identify the region.
- Integrate.

Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.


The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.

$$A = A_1 + A_2$$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx$$

$$+ \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

Example



- Find the point of intersection.
- Graph the functions.
- Identify the region.
- Integrate.

Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$.

The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$.


$$A = A_1 + A_2$$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx$$

$$+ \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

$$= [\sin x + \cos x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{\pi/2}$$

Example



- Find the point of intersection.
- Graph the functions.
- Identify the region.
- Integrate.

Find the area of the region enclosed by the curves $y = \sin x$, $y = \cos x$, x = 0 and $x = \pi/2$. The only point of intersection in the interval $[0, \pi/2]$ is $(\pi/4, 1/\sqrt{2})$. $A = A_1 + A_2$

$$= \int_0^{\pi/4} (\cos x - \sin x) dx$$

$$+ \int_{\pi/4}^{\pi/2} (\sin x - \cos x) dx$$

$$= [\sin x + \cos x]_0^{\pi/4} + [-\cos x - \sin x]_{\pi/4}^{\pi/2}$$

$$= 2\sqrt{2} - 2$$