

Calculus I

Lecture 21

The Fundamental Theorem of Calculus Part I

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1 Antiderivatives

Outline

- 1 Antiderivatives
- 2 Evaluating Definite Integrals
 - The Evaluation Theorem (FTC part 2)
 - Indefinite Integrals

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:

<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Antiderivatives

Definition (Antiderivative)

A function F is called an antiderivative of f on an interval I if $F'(x) = f(x)$ for all x in I .

Example

- Let $f(x) = x^2$.

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) = \quad$, then $F'(x) = x^2 = f(x)$.

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) =$, then $F'(x) = x^2 = f(x)$.

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) = \frac{1}{3}x^3$, then $F'(x) = x^2 = f(x)$.

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) = \frac{1}{3}x^3$, then $F'(x) = x^2 = f(x)$.
- Is this the only one?

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) = \frac{1}{3}x^3$, then $F'(x) = x^2 = f(x)$.
- Is this the only one?
- No. If $G(x) = \frac{1}{3}x^3 + 1$, then $G'(x) = x^2 = f(x)$.

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) = \frac{1}{3}x^3$, then $F'(x) = x^2 = f(x)$.
- Is this the only one?
- No. If $G(x) = \frac{1}{3}x^3 + 1$, then $G'(x) = x^2 = f(x)$.
- $\frac{1}{3}x^3 + 2$ will also work.

Example

- Let $f(x) = x^2$.
- Use the Power Rule to find an antiderivative of f :
- If $F(x) = \frac{1}{3}x^3$, then $F'(x) = x^2 = f(x)$.
- Is this the only one?
- No. If $G(x) = \frac{1}{3}x^3 + 1$, then $G'(x) = x^2 = f(x)$.
- $\frac{1}{3}x^3 + 2$ will also work.
- Any function of the form $H(x) = \frac{1}{3}x^3 + C$, where C is a constant, is an antiderivative of f .

Theorem

If F is an antiderivative of f on an interval I , then an arbitrary antiderivative of f on I is of the form

$$F(x) + C$$

where C is an arbitrary constant.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) =$, then
 $F'(x) = \sin x$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) =$, then
 $F'(x) = \sin x$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) = -\cos x$, then
 $F'(x) = \sin x$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) = -\cos x$, then $F'(x) = \sin x$.
- Therefore antiderivative is of the form $G(x) = -\cos x + C$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) = -\cos x$, then $F'(x) = \sin x$.
- Therefore antiderivative is of the form $G(x) = -\cos x + C$.
- If $F(x) = \frac{x^{n+1}}{n+1}$, then $F'(x) = x^n$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) = -\cos x$, then $F'(x) = \sin x$.
- Therefore antiderivative is of the form $G(x) = -\cos x + C$.
- If $F(x) = \frac{x^{n+1}}{n+1}$, then $F'(x) = x^n$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

$$f(x) = x^n, n \geq 0$$

- If $F(x) = -\cos x$, then $F'(x) = \sin x$.

- Therefore antiderivative is of the form $G(x) = -\cos x + C$.

- If $F(x) = \frac{x^{n+1}}{n+1}$, then $F'(x) = x^n$.

Example

Find all antiderivatives of each of the following functions.

$$f(x) = \sin x$$

- If $F(x) = -\cos x$, then $F'(x) = \sin x$.
- Therefore antiderivative is of the form $G(x) = -\cos x + C$.

$$f(x) = x^n, n \geq 0$$

- If $F(x) = \frac{x^{n+1}}{n+1}$, then $F'(x) = x^n$.
- Therefore any antiderivative is of the form $G(x) = \frac{x^{n+1}}{n+1} + C$.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) =$, then $F'(x) = \frac{1}{x}$.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) =$, then $F'(x) = \frac{1}{x}$.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) = \ln |x|$, then $F'(x) = \frac{1}{x}$.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) = \ln |x|$, then $F'(x) = \frac{1}{x}$.
- This is valid for any interval on which $\frac{1}{x}$ is defined.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) = \ln |x|$, then $F'(x) = \frac{1}{x}$.
- This is valid for any interval on which $\frac{1}{x}$ is defined.
- $\frac{1}{x}$ is defined

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) = \ln |x|$, then $F'(x) = \frac{1}{x}$.
- This is valid for any interval on which $\frac{1}{x}$ is defined.
- $\frac{1}{x}$ is defined everywhere except at 0.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) = \ln |x|$, then $F'(x) = \frac{1}{x}$.
- This is valid for any interval on which $\frac{1}{x}$ is defined.
- $\frac{1}{x}$ is defined everywhere except at 0.
- The most general answer needs two different constants, one for $(-\infty, 0)$ and one for $(0, \infty)$.

Example

Find the most general antiderivative of $f(x) = \frac{1}{x}$.

- If $F(x) = \ln |x|$, then $F'(x) = \frac{1}{x}$.
- This is valid for any interval on which $\frac{1}{x}$ is defined.
- $\frac{1}{x}$ is defined everywhere except at 0.
- The most general answer needs two different constants, one for $(-\infty, 0)$ and one for $(0, \infty)$.

$$G(x) = \begin{cases} \ln |x| + C_1 & \text{if } x > 0 \\ \ln |x| + C_2 & \text{if } x < 0 \end{cases}$$

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	
$x^n (n \neq -1)$	
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula.
Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula.
Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula.
Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula.
Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	$\tan x$
$\sec x \tan x$	

Every differentiation formula gives rise to an antiderivation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	$\tan x$
$\sec x \tan x$	

Every differentiation formula gives rise to an antidifferentiation formula. Suppose $F' = f$ and $G' = g$.

Function	Particular Antiderivative
$cf(x)$	$cF(x)$
$f(x) + g(x)$	$F(x) + G(x)$
$x^n (n \neq -1)$	$\frac{x^{n+1}}{n+1}$
$\frac{1}{x}$	$\ln x $
e^x	e^x
$\cos x$	$\sin x$
$\sin x$	$-\cos x$
$\sec^2 x$	$\tan x$
$\sec x \tan x$	$\sec x$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4 \quad + 2 \quad -$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4 \quad + 2 \quad -$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2x^5 - 2\sqrt{x} + C$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \quad -$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} -$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} -$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} - \frac{x^{1/2}}{\frac{1}{2}}$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$g'(x) = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

$$g(x) = 4(-\cos x) + 2 \frac{x^5}{5} - \frac{x^{1/2}}{\frac{1}{2}} + C$$

Example

Find all functions g such that

$$g'(x) = 4 \sin x + \frac{2x^5 - \sqrt{x}}{x}.$$

Rewrite:

$$g'(x) = 4 \sin x + 2 \frac{x^5}{x} - \frac{\sqrt{x}}{x} = 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}}$$

Find the antiderivative:

$$\begin{aligned} g'(x) &= 4 \sin x + 2x^4 - \frac{1}{\sqrt{x}} \\ g(x) &= 4(-\cos x) + 2 \frac{x^5}{5} - \frac{x^{1/2}}{\frac{1}{2}} + C \\ &= -4 \cos x + \frac{2}{5} x^5 - 2\sqrt{x} + C \end{aligned}$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$f(x) =$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$f(x) = \frac{x^{-1/2}}{-\frac{1}{2}}$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$f(x) = \frac{x^{-1/2}}{-\frac{1}{2}} + C$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

To find C , use the fact that $f(1) = 1$.

$$f(1) = 1$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

To find C , use the fact that $f(1) = 1$.

$$\begin{aligned} f(1) &= 1 \\ -\frac{2}{\sqrt{1}} + C &= 1 \end{aligned}$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

To find C , use the fact that $f(1) = 1$.

$$f(1) = 1$$

$$-\frac{2}{\sqrt{1}} + C = 1$$

$$C = 3$$

Example

Find f if $f'(x) = \frac{1}{x\sqrt{x}}$ for $x > 0$, and $f(1) = 1$.

$$f'(x) = \frac{1}{x\sqrt{x}} = x^{-3/2}$$

To find C , use the fact that $f(1) = 1$.

$$\begin{aligned} f(x) &= \frac{x^{-1/2}}{-\frac{1}{2}} + C \\ &= -\frac{2}{\sqrt{x}} + C \end{aligned}$$

$$f(1) = 1$$

$$-\frac{2}{\sqrt{1}} + C = 1$$

$$C = 3$$

Therefore

$$f(x) = -\frac{2}{\sqrt{x}} + 3.$$

Theorem (The Evaluation Theorem (FTC part 2))

If f is continuous on $[a, b]$, then

$$\int_a^b f(x)dx = F(b) - F(a),$$

where F is any antiderivative of f .

$\int_a^b f(x)dx$ exists for any continuous (over $[a, b]$)
function f .

Theorem (The Evaluation Theorem (FTC part 2))

If f is continuous on $[a, b]$, then

$$\int_a^b f(x)dx = F(b) - F(a),$$

where F is any antiderivative of f .

Theorem

Let f be a continuous function on $[a, b]$. Then f is integrable over $[a, b]$.

In other words, $\int_a^b f(x)dx$ exists for any continuous (over $[a, b]$) function f .


Theorem (The Evaluation Theorem (FTC part 2))

If f is continuous on $[a, b]$, then

$$\int_a^b f(x)dx = F(b) - F(a),$$


where F is any antiderivative of f .

Example



Evaluate the integral $\int_{-2}^1 x^3 dx$.


Example



Evaluate the integral $\int_{-2}^1 x^3 \, dx$.

- x^3 is continuous on $[-2, 1]$ (in fact, it's continuous everywhere).


Example



Evaluate the integral $\int_{-2}^1 x^3 dx$.

- x^3 is continuous on $[-2, 1]$ (in fact, it's continuous everywhere).
- An antiderivative is $F(x) = ?$


Example



Evaluate the integral $\int_{-2}^1 x^3 dx$.

- x^3 is continuous on $[-2, 1]$ (in fact, it's continuous everywhere).
- An antiderivative is $F(x) = \frac{1}{4}x^4$.

Example




Evaluate the integral $\int_{-2}^1 x^3 dx$.

- x^3 is continuous on $[-2, 1]$ (in fact, it's continuous everywhere).
- An antiderivative is $F(x) = \frac{1}{4}x^4$.

$$\int_{-2}^1 x^3 dx = F(1) - F(-2)$$

Example




Evaluate the integral $\int_{-2}^1 x^3 dx$.

- x^3 is continuous on $[-2, 1]$ (in fact, it's continuous everywhere).
- An antiderivative is $F(x) = \frac{1}{4}x^4$.

$$\int_{-2}^1 x^3 dx = F(1) - F(-2) = \frac{1}{4}(1)^4 - \frac{1}{4}(-2)^4$$

Example



Evaluate the integral $\int_{-2}^1 x^3 dx$.

- x^3 is continuous on $[-2, 1]$ (in fact, it's continuous everywhere).
- An antiderivative is $F(x) = \frac{1}{4}x^4$.

$$\int_{-2}^1 x^3 dx = F(1) - F(-2) = \frac{1}{4}(1)^4 - \frac{1}{4}(-2)^4 = \frac{1}{4} - \frac{16}{4} = -\frac{15}{4}$$

We often use the notation

$$F(x)]_a^b = F(b) - F(a)$$

or

$$[F(x)]_a^b = F(b) - F(a)$$

Therefore we can write


$$\int_a^b f(x)dx = F(x)]_a^b$$

or

$$\int_a^b f(x)dx = [F(x)]_a^b$$

Example


Find the area under the parabola $y = x^2$ from 0 to 1.



Example

Find the area under the parabola $y = x^2$ from 0 to 1.


- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).



Example

Find the area under the parabola $y = x^2$ from 0 to 1.


- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is ?



Example

Find the area under the parabola $y = x^2$ from 0 to 1.


- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is $\frac{1}{3}x^3$.



Example

Find the area under the parabola $y = x^2$ from 0 to 1.

- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is $\frac{1}{3}x^3$.




$$\int_0^1 x^2 \, dx = \left[\frac{1}{3}x^3 \right]_0^1$$

Example

Find the area under the parabola $y = x^2$ from 0 to 1.

- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is $\frac{1}{3}x^3$.




$$\int_0^1 x^2 \, dx = \left[\frac{1}{3}x^3 \right]_0^1$$

Example

Find the area under the parabola $y = x^2$ from 0 to 1.

- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is $\frac{1}{3}x^3$.




$$\int_0^1 x^2 dx = \left[\frac{1}{3}x^3 \right]_0^1 = \frac{1}{3}(1)^3 - \frac{1}{3}(0)^3$$

Example

Find the area under the parabola $y = x^2$ from 0 to 1.

- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is $\frac{1}{3}x^3$.




$$\int_0^1 x^2 \, dx = \left[\frac{1}{3}x^3 \right]_0^1 = \frac{1}{3}(1)^3 - \frac{1}{3}(0)^3$$

Example

Find the area under the parabola $y = x^2$ from 0 to 1.

- x^2 is continuous on $[0, 1]$ (in fact, it's continuous everywhere).
- An antiderivative of x^2 is $\frac{1}{3}x^3$.



$$\int_0^1 x^2 \, dx = \left[\frac{1}{3}x^3 \right]_0^1 = \frac{1}{3}(1)^3 - \frac{1}{3}(0)^3 = \frac{1}{3}$$

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- **An antiderivative of $\cos x$ is ?**

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- **An antiderivative of $\cos x$ is $\sin x$.**

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- An antiderivative of $\cos x$ is $\sin x$.

$$\int_0^b \cos x \, dx = [\sin x]_0^b$$

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- **An antiderivative of $\cos x$ is $\sin x$.**

$$\int_0^b \cos x \, dx = [\sin x]_0^b$$

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- An antiderivative of $\cos x$ is $\sin x$.

$$\int_0^b \cos x \, dx = [\sin x]_0^b = \sin(b) - \sin(0)$$

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- An antiderivative of $\cos x$ is $\sin x$.

$$\int_0^b \cos x \, dx = [\sin x]_0^b = \sin(b) - \sin(0)$$

Example



Find the area under the cosine curve from 0 to b , where $0 \leq b \leq \frac{\pi}{2}$.

- $\cos x$ is continuous on $[0, \frac{\pi}{2}]$ (in fact, it's continuous everywhere).
- An antiderivative of $\cos x$ is $\sin x$.

$$\int_0^b \cos x \, dx = [\sin x]_0^b = \sin(b) - \sin(0) = \sin b$$

Indefinite Integrals

- The Evaluation Theorem establishes a connection between antiderivatives and definite integrals.
- It says that $\int_a^b f(x)dx$ equals $F(b) - F(a)$, where F is an antiderivative of f .
- We need convenient notation for writing antiderivatives.
- This is what the indefinite integral is.

Definition (Indefinite Integral)

The indefinite integral of f is another way of saying the antiderivative of f , and is written $\int f(x)dx$. In other words,

$$\int f(x)dx = F(x) \quad \text{means} \quad F'(x) = f(x).$$

Example

$$\int x^4 dx = ?$$

Example

$$\int x^4 dx = \frac{x^5}{5}$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of $\frac{1}{x}$ is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of $\frac{1}{x}$ is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

- We adopt the convention that the constant participating in an indefinite integral is only valid on one interval.

Example

$$\int x^4 dx = \frac{x^5}{5} + C$$

because

$$\frac{d}{dx} \left(\frac{x^5}{5} + C \right) = x^4.$$

- The indefinite integral represents a whole family of functions.
- Example: the general antiderivative of $\frac{1}{x}$ is

$$F(x) = \begin{cases} \ln|x| + C_1 & \text{if } x > 0 \\ \ln|x| + C_2 & \text{if } x < 0 \end{cases}$$

- We adopt the convention that the constant participating in an indefinite integral is only valid on one interval.
- $\int \frac{1}{x} dx = \ln|x| + C$, and this is valid either on $(-\infty, 0)$ or $(0, \infty)$.

Example

Find the indefinite integral.

$$\int (8x^3 - 3 \sec^2 x) dx$$

Example

Find the indefinite integral.

$$\int (8x^3 - 3 \sec^2 x) dx = 8 \int x^3 dx - 3 \int \sec^2 x dx$$

Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8? - 3?\end{aligned}$$

Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3?\end{aligned}$$

Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \text{?}\end{aligned}$$

Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \tan x\end{aligned}$$

Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \tan x + C\end{aligned}$$

Example

Find the indefinite integral.

$$\begin{aligned}\int (8x^3 - 3 \sec^2 x) dx &= 8 \int x^3 dx - 3 \int \sec^2 x dx \\ &= 8 \frac{x^4}{4} - 3 \tan x + C \\ &= 2x^4 - 3 \tan x + C\end{aligned}$$

Example

Find the general indefinite integral.

$$\int \frac{\cos \theta}{\sin^2 \theta} d\theta$$

Example

Find the general indefinite integral.

$$\int \frac{\cos \theta}{\sin^2 \theta} d\theta = \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \text{?} \quad \text{?} \quad d\theta\end{aligned}$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta? \quad d\theta\end{aligned}$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \text{?} \quad d\theta\end{aligned}$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta\end{aligned}$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta \\ &= ?\end{aligned}$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta \\ &= -\csc \theta\end{aligned}$$

Example

Find the general indefinite integral.

$$\begin{aligned}\int \frac{\cos \theta}{\sin^2 \theta} d\theta &= \int \left(\frac{1}{\sin \theta} \right) \left(\frac{\cos \theta}{\sin \theta} \right) d\theta \\ &= \int \csc \theta \cot \theta d\theta \\ &= -\csc \theta + C\end{aligned}$$

Example

$$\int_0^3 (x^3 - 6x) dx$$

Example

$$\int_0^3 (x^3 - 6x) dx = \left[\int (x^3 - 6x) dx \right]_0^3$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[\int (x^3 - 6x)dx \right]_0^3 \\ &= \left[\int x^3 dx - 6 \int x dx \right]_0^3\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[\int (x^3 - 6x)dx \right]_0^3 \\ &= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\ &= \left[? - 6? \right]_0^3\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[\int (x^3 - 6x)dx \right]_0^3 \\ &= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\ &= \left[\frac{x^4}{4} - 6x \right]_0^3\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[\int (x^3 - 6x) dx \right]_0^3 \\ &= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\ &= \left[\frac{x^4}{4} - 6? \right]_0^3\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x)dx &= \left[\int (x^3 - 6x)dx \right]_0^3 \\&= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[\frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[\int (x^3 - 6x) dx \right]_0^3 \\&= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[\frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left(\frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left(\frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right)\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[\int (x^3 - 6x) dx \right]_0^3 \\&= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[\frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left(\frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left(\frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right)\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[\int (x^3 - 6x) dx \right]_0^3 \\&= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[\frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left(\frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left(\frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right) \\&= \frac{81}{4} - 27 - 0 + 0\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[\int (x^3 - 6x) dx \right]_0^3 \\&= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[\frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left(\frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left(\frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right) \\&= \frac{81}{4} - 27 - 0 + 0\end{aligned}$$

Example

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \left[\int (x^3 - 6x) dx \right]_0^3 \\&= \left[\int x^3 dx - 6 \int x dx \right]_0^3 \\&= \left[\frac{x^4}{4} - 6 \frac{x^2}{2} \right]_0^3 \\&= \left(\frac{1}{4} \cdot 3^4 - 3 \cdot 3^2 \right) - \left(\frac{1}{4} \cdot 0^4 - 3 \cdot 0^2 \right) \\&= \frac{81}{4} - 27 - 0 + 0 = -\frac{27}{4}.\end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

Example

Evaluate:
$$\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$$

$$= \int_1^9 \left(2t + t^{\frac{1}{2}} - t^{-2} \right) dt$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$= \int_1^9 \left(2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9$$
$$= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9$$
$$= \left[? + ? - ? \right]_1^9$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left(2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + ? - ? \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left(2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + ? - ? \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left(2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - ? \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - ? \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[t^2 + \frac{2}{3}t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left(9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left(1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left(9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left(1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 \left(2t + t^{\frac{1}{2}} - t^{-2} \right) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left(9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left(1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \\ &= 81 + 18 + \frac{1}{9} - 1 - \frac{2}{3} - 1 \end{aligned}$$

Example

Evaluate: $\int_1^9 \frac{2t^3 + t^2\sqrt{t} - 1}{t^2} dt$

$$\begin{aligned} &= \int_1^9 (2t + t^{\frac{1}{2}} - t^{-2}) dt = \left[\int (2t + t^{\frac{1}{2}} - t^{-2}) dt \right]_1^9 \\ &= \left[\int 2t dt + \int t^{\frac{1}{2}} dt - \int t^{-2} dt \right]_1^9 \\ &= \left[t^2 + \frac{t^{\frac{3}{2}}}{\frac{3}{2}} - \frac{t^{-1}}{-1} \right]_1^9 = \left[t^2 + \frac{2}{3} t^{\frac{3}{2}} + \frac{1}{t} \right]_1^9 \\ &= \left(9^2 + \frac{2}{3} \cdot 9^{\frac{3}{2}} + \frac{1}{9} \right) - \left(1^2 + \frac{2}{3} \cdot 1^{\frac{3}{2}} + \frac{1}{1} \right) \\ &= 81 + 18 + \frac{1}{9} - 1 - \frac{2}{3} - 1 = \frac{868}{9}. \end{aligned}$$