

# Calculus III

## Lecture 6

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Curves in space

# Outline

- 1 Curves in space
- 2 Tangent vectors, tangents

# Outline

- 1 Curves in space
- 2 Tangent vectors, tangents
- 3 Line integrals

# Outline

- 1 Curves in space
- 2 Tangent vectors, tangents
- 3 Line integrals
- 4 Curvature

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
`https://creativecommons.org/licenses/by/3.0/us/`  
and the links therein.

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in ?$$

$$(0, 5) \in$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$



$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$

$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$ 

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in ?$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in$$

$$(0, 1, 2, 3, \dots, n) \in$$

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in ?$$

$$(0, 1, 2, 3, \dots, n) \in$$

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in \mathbb{R}^5$$

$$(0, 1, 2, 3, \dots, n) \in$$

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in \mathbb{R}^5$$

$$(0, 1, 2, 3, \dots, n) \in ?$$

$$\mathbb{R}^2, \mathbb{R}^3, \mathbb{R}^n$$

### Definition ( $\mathbb{R}^2$ )

The set of ordered pairs of real numbers is denoted by  $\mathbb{R}^2$ .

### Definition ( $\mathbb{R}^3$ )

The set of ordered triples of real numbers is denoted by  $\mathbb{R}^3$ .

### Definition ( $\mathbb{R}^n$ )

The set of ordered  $n$ -tuples of real numbers is denoted by  $\mathbb{R}^n$ .

### Example

$$(1, -2, 3) \in \mathbb{R}^3$$

$$(0, 5) \in \mathbb{R}^2$$

$$(0, 5, -2, 4, 0) \in \mathbb{R}^5$$

$$(0, 1, 2, 3, \dots, n) \in \mathbb{R}^{n+1}$$

# Parametric Equations of a Line Segment

- Recall parametric vector equation of line  $L$ :

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{u}, \quad t \text{ real number.}$$

$$\mathbf{r} = \mathbf{r}_0 + t(\mathbf{r}_1 - \mathbf{r}_0), \quad t \text{ real number.}$$

$$\mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1, \quad t \text{ real number.}$$



# Parametric Equations of a Line Segment

- Recall parametric vector equation of line  $L$ :

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{u}, \quad t \text{ real number.}$$

$$\mathbf{r} = \mathbf{r}_0 + t(\mathbf{r}_1 - \mathbf{r}_0), \quad t \text{ real number.}$$

$$\mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1, \quad t \text{ real number.}$$

- Parametric scalar equations:

$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} \Leftrightarrow \begin{cases} x = x_0 + t(x_1 - x_0) \\ y = y_0 + t(y_1 - y_0) \\ z = z_0 + t(z_1 - z_0) \end{cases} \Leftrightarrow \begin{cases} x = (1 - t)x_0 + tx_1 \\ y = (1 - t)y_0 + ty_1 \\ z = (1 - t)z_0 + tz_1 \end{cases}$$

# Parametric Equations of a Line Segment

- Recall parametric vector equation of line  $L$ :

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{u}, \quad t \text{ real number.}$$

$$\mathbf{r} = \mathbf{r}_0 + t(\mathbf{r}_1 - \mathbf{r}_0), \quad t \text{ real number.}$$

$$\mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1, \quad t \text{ real number.}$$

- Parametric scalar equations:

$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} \Leftrightarrow \begin{cases} x = x_0 + t(x_1 - x_0) \\ y = y_0 + t(y_1 - y_0) \\ z = z_0 + t(z_1 - z_0) \end{cases} \Leftrightarrow \begin{cases} x = (1 - t)x_0 + tx_1 \\ y = (1 - t)y_0 + ty_1 \\ z = (1 - t)z_0 + tz_1 \end{cases}$$

- Segment with endpoints  $P_0(\mathbf{r}_0)$  and  $P_1(\mathbf{r}_1)$ :

$$\mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1, \quad 0 \leq t \leq 1$$

## Example

### Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

## Example

Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

Direction of  $L$ :

## Example

Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

Direction of  $L$ :  $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$ .

## Example

Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

Direction of  $L$ :  $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$ . Parametric vectorial equations of  $L$ :

## Example

Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

Direction of  $L$ :  $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$ . Parametric vectorial equations of  $L$ :

$$\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t) .$$

## Example

### Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

Direction of  $L$ :  $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$ . Parametric vectorial equations of  $L$ :

$$\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t) .$$

Parametric scalar equations of line  $L$ :

$$\begin{cases} x = 1 + 4t \\ y = 2 \\ z = 3 - 2t \end{cases}, \quad t \text{ real number.}$$



## Example

### Parametrize

- the line  $L$  passing through  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ ;
- the line segment  $S$  connecting  $P_0(1, 2, 3)$  and  $P_1(5, 2, 1)$ .

Direction of  $L$ :  $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$ . Parametric vectorial equations of  $L$ :

$$\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t) .$$

Parametric scalar equations of line  $L$ :

$$\begin{cases} x = 1 + 4t \\ y = 2 \\ z = 3 - 2t \end{cases}, \quad t \text{ real number.}$$

Parametric vectorial equation of segment  $S$ :

$$\mathbf{r} = t(1, 2, 3) + (1 - t)(5, 2, 1) \quad t \in [0, 1] .$$

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  plane curve.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  plane curve.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$  space curve.



# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  plane curve.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$  space curve.
- Function  $\mathbf{r}$ : parametrization of the curve image.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  plane curve.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$  space curve.
- Function  $\mathbf{r}$ : parametrization of the curve image.
- $t$ : parameter of the curve parametrization.

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  plane curve.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$  space curve.
- Function  $\mathbf{r}$ : parametrization of the curve image.
- $t$ : parameter of the curve parametrization.
- $\mathbf{r}(t) = (x(t), y(t), z(t))$ ,  $a \leq t \leq b$ .

# Parametrized Curves

- A curve parametrization is a function  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$ , or  $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^n$  in general.
- Input is scalar (parameter).
- Output is (position) vector.
- The image of  $\mathbf{r}$  is a set of points in  $\mathbb{R}^n$ ; we call that set curve image.
- The term “curve” is ambiguous and either means a curve parametrization or a curve image.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^2$  plane curve.
- $\mathbf{r} : [a, b] \rightarrow \mathbb{R}^3$  space curve.
- Function  $\mathbf{r}$ : parametrization of the curve image.
- $t$ : parameter of the curve parametrization.
- $\mathbf{r}(t) = (x(t), y(t), z(t))$ ,  $a \leq t \leq b$ .
- $x, y, z : [a, b] \rightarrow \mathbb{R}$ , coordinate functions.

# Example

- $\mathbf{r}: \mathbb{R} \rightarrow \mathbb{R}^3,$

$$\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{u}$$

# Example

- $\mathbf{r}: \mathbb{R} \rightarrow \mathbb{R}^3$ ,

$$\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{u}$$

- Line through  $P(1, 4, 3)$  direction  $\mathbf{u} = (-1, 2, 0)$ :

$$\mathbf{r}(t) = (1 - t, 4 + 2t, 3)$$

# Example

- $\mathbf{r}: \mathbb{R} \rightarrow \mathbb{R}^3$ ,

$$\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{u}$$

- Line through  $P(1, 4, 3)$  direction  $\mathbf{u} = (-1, 2, 0)$ :

$$\mathbf{r}(t) = (1 - t, 4 + 2t, 3)$$

- Curve parametrization is not unique!

# Example

- $\mathbf{r}: \mathbb{R} \rightarrow \mathbb{R}^3$ ,

$$\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{u}$$

- Line through  $P(1, 4, 3)$  direction  $\mathbf{u} = (-1, 2, 0)$ :

$$\mathbf{r}(t) = (1 - t, 4 + 2t, 3)$$

$$\mathbf{p}(s) =$$

- Curve parametrization is not unique!
- For example  $t = 1 - s = \varphi(s)$  is a reparametrization.



# Example

- $\mathbf{r}: \mathbb{R} \rightarrow \mathbb{R}^3$ ,

$$\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{u}$$

- Line through  $P(1, 4, 3)$  direction  $\mathbf{u} = (-1, 2, 0)$ :

$$\mathbf{r}(t) = (1 - t, 4 + 2t, 3)$$

$$\mathbf{p}(s) = (s, 6 - s, 3)$$

- Curve parametrization is not unique!
- For example  $t = 1 - s = \varphi(s)$  is a reparametrization.


## Example

Describe the curve:

$$x(t) = 3t \cos(2t)$$

$$y(t) = 3t \sin(2t)$$

$$z(t) = t^2$$




## Example

Describe the curve:

$$x(t) = 3t \cos(2t)$$

$$y(t) = 3t \sin(2t)$$

$$z(t) = t^2$$



● Cylindrical coordinates:

$$r(t) = 3t$$

$$\theta(t) = 2t$$

$$z(t) = t^2$$


## Example

Describe the curve:

$$x(t) = 3t \cos(2t)$$

$$y(t) = 3t \sin(2t)$$

$$z(t) = t^2$$



- Cylindrical coordinates:

$$r(t) = 3t$$

$$\theta(t) = 2t$$

$$z(t) = t^2$$

- “Tornado”.

# Limits

## Definition

We say that

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$$

if by selecting that  $t \neq a$  be close enough to  $a$  we can guarantee that  $\mathbf{r}(t)$  is as close to  $\mathbf{u}$  as we want.

# Limits

## Definition

We say that

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$$

if by selecting that  $t \neq a$  be close enough to  $a$  we can guarantee that  $\mathbf{r}(t)$  is as close to  $\mathbf{u}$  as we want.

In strict mathematical language:  $\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $t$  with  $0 < |t - a| < \delta$  we have that  $|\mathbf{r}(t) - \mathbf{u}| < \varepsilon$ .

# Limits

## Definition

We say that

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$$

if by selecting that  $t \neq a$  be close enough to  $a$  we can guarantee that  $\mathbf{r}(t)$  is as close to  $\mathbf{u}$  as we want.

In strict mathematical language:  $\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $t$  with  $0 < |t - a| < \delta$  we have that  $|\mathbf{r}(t) - \mathbf{u}| < \varepsilon$ .

# Limits

## Definition

We say that

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$$

if **by selecting that**  $t \neq a$  be close enough to  $a$  we can guarantee that  $\mathbf{r}(t)$  is as close to  $\mathbf{u}$  as we want.

In strict mathematical language:  $\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$  if for every  $\varepsilon > 0$  **there exists**  $\delta > 0$  such that for all  $t$  with  $0 < |t - a| < \delta$  we have that  $|\mathbf{r}(t) - \mathbf{u}| < \varepsilon$ .



# Limits

## Definition

We say that

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$$

if by selecting that  $t \neq a$  be close enough to  $a$  we can guarantee that  $\mathbf{r}(t)$  is as close to  $\mathbf{u}$  as we want.

In strict mathematical language:  $\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u}$  if for every  $\varepsilon > 0$  there exists  $\delta > 0$  such that for all  $t$  with  $0 < |t - a| < \delta$  we have that  $|\mathbf{r}(t) - \mathbf{u}| < \varepsilon$ .

- We define the “postman distance” between  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  to be the number  $\max(|x_1 - x_2|, |y_1 - y_2|, |z_1 - z_2|)$ .

- We define the “postman distance” between  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  to be the number  $\max(|x_1 - x_2|, |y_1 - y_2|, |z_1 - z_2|)$ .
- Two points in Euclidean distance are close if and only if they are close in “postman distance”.

- We define the “postman distance” between  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  to be the number  $\max(|x_1 - x_2|, |y_1 - y_2|, |z_1 - z_2|)$ .
- Two points in Euclidean distance are close if and only if they are close in “postman distance”.
- Unlike higher dimensions, in dimension 1 postman distance coincides with Euclidean distance.

- We define the “postman distance” between  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  to be the number  $\max(|x_1 - x_2|, |y_1 - y_2|, |z_1 - z_2|)$ .
- Two points in Euclidean distance are close if and only if they are close in “postman distance”.
- Unlike higher dimensions, in dimension 1 postman distance coincides with Euclidean distance.
- Let  $\mathbf{r}(t) = (x(t), y(t), z(t))$  and  $\mathbf{u} = (u_1, u_2, u_3)$ .

- We define the “postman distance” between  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  to be the number  $\max(|x_1 - x_2|, |y_1 - y_2|, |z_1 - z_2|)$ .
- Two points in Euclidean distance are close if and only if they are close in “postman distance”.
- Unlike higher dimensions, in dimension 1 postman distance coincides with Euclidean distance.
- Let  $\mathbf{r}(t) = (x(t), y(t), z(t))$  and  $\mathbf{u} = (u_1, u_2, u_3)$ .
- Then

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \mathbf{u} \iff \begin{cases} \lim_{t \rightarrow a} x(t) = u_1 \\ \lim_{t \rightarrow a} y(t) = u_2 \\ \lim_{t \rightarrow a} z(t) = u_3 \end{cases} .$$

# Continuity

## Definition

Suppose

- $\mathbf{r}$  is defined at  $t_0$

## Observation

$\mathbf{r}(t) = (x(t), y(t), z(t))$  is continuous at  $t_0 \iff x(t), y(t), z(t)$  are all continuous at  $t_0$ .

# Continuity

## Definition

Suppose

- $\mathbf{r}$  is defined at  $t_0$
- $\lim_{t \rightarrow t_0} \mathbf{r}(t)$  exists.

## Observation

$\mathbf{r}(t) = (x(t), y(t), z(t))$  is continuous at  $t_0 \iff x(t), y(t), z(t)$  are all continuous at  $t_0$ .



# Continuity

## Definition

Suppose

- $\mathbf{r}$  is defined at  $t_0$
- $\lim_{t \rightarrow t_0} \mathbf{r}(t)$  exists.


Then we say that  $\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$  is continuous at  $t_0$  if

$$\lim_{t \rightarrow t_0} \mathbf{r}(t) = \mathbf{r}(t_0) \quad .$$

## Observation

$\mathbf{r}(t) = (x(t), y(t), z(t))$  is continuous at  $t_0 \iff x(t), y(t), z(t)$  are all continuous at  $t_0$ .


# Derivatives



$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t))\end{aligned}$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}.$


# Derivatives



$$\begin{aligned} \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t)) \end{aligned}$$

- Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .


# Derivatives



$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t))\end{aligned}$$

- Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}.$


# Derivatives



$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t))\end{aligned}$$

- Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}.$


# Derivatives



$$\begin{aligned} \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t)) \end{aligned}$$

- Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}.$

# Derivatives




$$\begin{aligned} \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t)) \end{aligned}$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .
- Instantaneous rate of change:

$$\mathbf{f}'(t_0) = \lim_{t \rightarrow t_0} \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$$

# Derivatives




$$\begin{aligned} \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t)) \end{aligned}$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .
- Instantaneous rate of change:

$$\mathbf{f}'(t_0) = \lim_{t \rightarrow t_0} \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0} = \lim_{h \rightarrow 0} \frac{\mathbf{f}(t_0 + h) - \mathbf{f}(t_0)}{h}.$$



# Derivatives




$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t))\end{aligned}$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .
- Instantaneous rate of change:

$$\mathbf{f}'(t_0) = \lim_{t \rightarrow t_0} \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0} = \lim_{h \rightarrow 0} \frac{\mathbf{f}(t_0 + h) - \mathbf{f}(t_0)}{h}.$$

# Derivatives




$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t))\end{aligned}$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .
- Instantaneous rate of change:

$$\mathbf{f}'(t_0) = \lim_{t \rightarrow t_0} \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0} = \lim_{h \rightarrow 0} \frac{\mathbf{f}(t_0 + h) - \mathbf{f}(t_0)}{h}.$$

- $\mathbf{f}(t)$  vector  $\implies \mathbf{f}'(t)$  vector.

# Derivatives



$$\mathbf{f}: [a, b] \rightarrow \mathbb{R}^3$$

$$\mathbf{f}(t) = (x(t), y(t), z(t))$$


$$\mathbf{f}'(t) = (x'(t), y'(t), z'(t))$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .
- Instantaneous rate of change:

$$\mathbf{f}'(t_0) = \lim_{t \rightarrow t_0} \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0} = \lim_{h \rightarrow 0} \frac{\mathbf{f}(t_0 + h) - \mathbf{f}(t_0)}{h}.$$

- $\mathbf{f}(t)$  vector  $\implies \mathbf{f}'(t)$  vector.

# Derivatives



$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t))\end{aligned}$$


$$\begin{aligned}\mathbf{f}'(t) &= (x'(t), y'(t), z'(t)) \\ \mathbf{f}''(t) &= (x''(t), y''(t), z''(t)) \\ &\vdots\end{aligned}$$

- **Average Velocity** =  $\frac{\text{change in position}}{\text{change in time}} = \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0}$ .
- Instantaneous rate of change:

$$\mathbf{f}'(t_0) = \lim_{t \rightarrow t_0} \frac{\mathbf{f}(t) - \mathbf{f}(t_0)}{t - t_0} = \lim_{h \rightarrow 0} \frac{\mathbf{f}(t_0 + h) - \mathbf{f}(t_0)}{h}.$$

- $\mathbf{f}(t)$  vector  $\implies \mathbf{f}'(t)$  vector.
- Higher order derivatives:  $\mathbf{f}'(t), \mathbf{f}''(t) = (\mathbf{f}'(t))'$  (acceleration), ...


# Tangent Lines



$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .


# Tangent Lines



$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .


# Tangent Lines



$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .

# Tangent Lines




$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .




# Tangent Lines



$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .


# Tangent Lines



$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .

# Tangent Lines




$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .
- Tangent equation at  $\mathbf{f}(t_0)$ :

$$\mathbf{r}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$

# Tangent Lines



$$\begin{aligned}
 \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\
 \mathbf{f}(t) &= (x(t), y(t), z(t)) \\
 \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\
 &= (x'(t), y'(t), z'(t))
 \end{aligned}$$


- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .
- Tangent equation at  $\mathbf{f}(t_0)$ :

$$\mathbf{r}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$

- Linear approximation:

$$\mathbf{f}(t) \approx L_{\mathbf{f}, t_0}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$

# Tangent Lines



$$\begin{aligned} \mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t)) \\ \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\ &= (x'(t), y'(t), z'(t)) \end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .

- Tangent equation at  $\mathbf{f}(t_0)$ :

$$\mathbf{r}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$


- Linear approximation:

$$\mathbf{f}(t) \approx L_{\mathbf{f}, t_0}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$

- A linear approximation is good if:

$$\lim_{t \rightarrow 0} \left| \frac{\mathbf{f}(t) - L_{\mathbf{f}, t_0}(t)}{t} \right| = 0.$$

# Tangent Lines



$$\begin{aligned}\mathbf{f}: [a, b] &\rightarrow \mathbb{R}^3 \\ \mathbf{f}(t) &= (x(t), y(t), z(t)) \\ \mathbf{f}'(t) &= \lim_{h \rightarrow 0} \frac{\mathbf{f}(t+h) - \mathbf{f}(t)}{h} \\ &= (x'(t), y'(t), z'(t))\end{aligned}$$

- $\mathbf{f}'(t_0)$ : direction of tangent line through  $\mathbf{f}(t_0)$ .

- Tangent equation at  $\mathbf{f}(t_0)$ :

$$\mathbf{r}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$

- Linear approximation:

$$\mathbf{f}(t) \approx L_{\mathbf{f}, t_0}(t) = \mathbf{f}(t_0) + t\mathbf{f}'(t_0).$$

- A linear approximation is good if:

$$\lim_{t \rightarrow 0} \left| \frac{\mathbf{f}(t) - L_{\mathbf{f}, t_0}(t)}{t} \right| = 0.$$

- Differentials:

$$d\mathbf{f} = \mathbf{f}'dt = (x', y', z')dt.$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) =$$



## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t)$ ,  $\mathbf{e}'_\phi(t)$ ,  $\mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) =$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t)$ ,  $\mathbf{e}'_\phi(t)$ ,  $\mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

$$\mathbf{e}'_\theta(t) =$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

$$\mathbf{e}'_\theta(t) = (-\rho \sin \phi \sin t, \rho \sin \phi \cos t, 0)$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter.

Find  $\mathbf{e}'_\rho(t)$ ,  $\mathbf{e}'_\phi(t)$ ,  $\mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

$$\mathbf{e}'_\theta(t) = (-\rho \sin \phi \sin t, \rho \sin \phi \cos t, 0)$$

$$(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\phi(\phi)) \cdot \mathbf{e}'_\theta(\theta) = \begin{vmatrix} \sin \phi \cos \theta & \sin \phi \sin \theta & \cos \phi \\ \rho \cos \phi \cos \theta & \rho \cos \phi \sin \theta & -\rho \sin \phi \\ -\rho \sin \phi \sin \theta & \rho \sin \phi \cos \theta & 0 \end{vmatrix}$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter.

Find  $\mathbf{e}'_\rho(t)$ ,  $\mathbf{e}'_\phi(t)$ ,  $\mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

$$\mathbf{e}'_\theta(t) = (-\rho \sin \phi \sin t, \rho \sin \phi \cos t, 0)$$

$$(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\phi(\phi)) \cdot \mathbf{e}'_\theta(\theta) = \begin{vmatrix} \sin \phi \cos \theta & \sin \phi \sin \theta & \cos \phi \\ \rho \cos \phi \cos \theta & \rho \cos \phi \sin \theta & -\rho \sin \phi \\ -\rho \sin \phi \sin \theta & \rho \sin \phi \cos \theta & 0 \end{vmatrix}$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter.

Find  $\mathbf{e}'_\rho(t)$ ,  $\mathbf{e}'_\phi(t)$ ,  $\mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

$$\mathbf{e}'_\theta(t) = (-\rho \sin \phi \sin t, \rho \sin \phi \cos t, 0)$$

$$(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\phi(\phi)) \cdot \mathbf{e}'_\theta(\theta) = \begin{vmatrix} \sin \phi \cos \theta & \sin \phi \sin \theta & \cos \phi \\ \rho \cos \phi \cos \theta & \rho \cos \phi \sin \theta & -\rho \sin \phi \\ -\rho \sin \phi \sin \theta & \rho \sin \phi \cos \theta & 0 \end{vmatrix}$$



## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\begin{aligned}\mathbf{e}_\rho(t) &= (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi) \\ \mathbf{e}_\phi(t) &= (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t) \\ \mathbf{e}_\theta(t) &= (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)\end{aligned}$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\begin{aligned}\mathbf{e}'_\rho(t) &= (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi) \\ \mathbf{e}'_\phi(t) &= (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t) \\ \mathbf{e}'_\theta(t) &= (-\rho \sin \phi \sin t, \rho \sin \phi \cos t, 0)\end{aligned}$$

$$\begin{aligned}(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\phi(\phi)) \cdot \mathbf{e}'_\theta(\theta) &= \begin{vmatrix} \sin \phi \cos \theta & \sin \phi \sin \theta & \cos \phi \\ \rho \cos \phi \cos \theta & \rho \cos \phi \sin \theta & -\rho \sin \phi \\ -\rho \sin \phi \sin \theta & \rho \sin \phi \cos \theta & 0 \end{vmatrix} \\ &= \end{aligned}$$

## Example

Let  $\mathbf{r}(t)$  be the coordinate curves for the spherical coordinates, i.e., let

$$\mathbf{e}_\rho(t) = (t \sin \phi \cos \theta, t \sin \phi \sin \theta, t \cos \phi)$$

$$\mathbf{e}_\phi(t) = (\rho \sin t \cos \theta, \rho \sin t \sin \theta, \rho \cos t)$$

$$\mathbf{e}_\theta(t) = (\rho \sin \phi \cos t, \rho \sin \phi \sin t, \rho \cos \phi)$$

where  $\rho, \phi, \theta$  are regarded as constants and  $t$  as the curve parameter. Find  $\mathbf{e}'_\rho(t), \mathbf{e}'_\phi(t), \mathbf{e}'_\theta(t)$ . Compute  $(\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\theta(\theta)) \cdot \mathbf{e}'_\phi(\phi)$ .

$$\mathbf{e}'_\rho(t) = (\sin \phi \cos \theta, \sin \phi \sin \theta, \cos \phi)$$

$$\mathbf{e}'_\phi(t) = (\rho \cos t \cos \theta, \rho \cos t \sin \theta, -\rho \sin t)$$

$$\mathbf{e}'_\theta(t) = (-\rho \sin \phi \sin t, \rho \sin \phi \cos t, 0)$$

$$\begin{aligned} (\mathbf{e}'_\rho(\rho) \times \mathbf{e}'_\phi(\phi)) \cdot \mathbf{e}'_\theta(\theta) &= \begin{vmatrix} \sin \phi \cos \theta & \sin \phi \sin \theta & \cos \phi \\ \rho \cos \phi \cos \theta & \rho \cos \phi \sin \theta & -\rho \sin \phi \\ -\rho \sin \phi \sin \theta & \rho \sin \phi \cos \theta & 0 \end{vmatrix} \\ &= \rho^2 \sin \phi \end{aligned}$$

# Differentiation Rules

Component-wise operation  $\implies$  same rules as for scalar output

# Differentiation Rules

Component-wise operation  $\implies$  same rules as for scalar output

Product Rules:

$$[f(t)\mathbf{r}(t)]' = f'(t)\mathbf{r}(t) + f(t)\mathbf{r}'(t)$$

$$[\mathbf{u}(t) \cdot \mathbf{v}(t)]' = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

$$[\mathbf{u}(t) \times \mathbf{v}(t)]' = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

# Differentiation Rules

Component-wise operation  $\implies$  same rules as for scalar output

Product Rules:

$$[f(t)\mathbf{r}(t)]' = f'(t)\mathbf{r}(t) + f(t)\mathbf{r}'(t)$$

$$[\mathbf{u}(t) \cdot \mathbf{v}(t)]' = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

$$[\mathbf{u}(t) \times \mathbf{v}(t)]' = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

Chain Rule:

$$[\mathbf{r}(f(t))]' = f'(t)\mathbf{r}'(f(t))$$

# Differentiation Rules

Component-wise operation  $\implies$  same rules as for scalar output

Product Rules:

$$[f(t)\mathbf{r}(t)]' = f'(t)\mathbf{r}(t) + f(t)\mathbf{r}'(t)$$

$$[\mathbf{u}(t) \cdot \mathbf{v}(t)]' = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

$$[\mathbf{u}(t) \times \mathbf{v}(t)]' = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

Chain Rule:

$$[\mathbf{r}(f(t))]' = f'(t)\mathbf{r}'(f(t))$$

Example:

$$\begin{aligned} \frac{d|\mathbf{r}(t)|}{dt} &= [\sqrt{\mathbf{r}(t) \cdot \mathbf{r}(t)}]' = [\sqrt{\square}]' = \frac{1}{2\sqrt{\square}}\square' = \frac{1}{2\sqrt{\mathbf{r}(t) \cdot \mathbf{r}(t)}}[\mathbf{r}(t) \cdot \mathbf{r}(t)]' = \\ &= \frac{1}{2|\mathbf{r}(t)|}[\mathbf{r}'(t) \cdot \mathbf{r}(t) + \mathbf{r}(t) \cdot \mathbf{r}'(t)] = \frac{\mathbf{r}(t) \cdot \mathbf{r}'(t)}{|\mathbf{r}(t)|} \end{aligned}$$

# Application

$$|\mathbf{r}'| = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:

# Application

$$|\mathbf{r}'| = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:

$$|\mathbf{r}(t)| = \text{constant}$$



# Application

$$|\mathbf{r}'| = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:

$$\begin{aligned} |\mathbf{r}(t)| &= \text{constant} \\ |\mathbf{r}(t)|' &= 0 \end{aligned}$$

# Application

$$|\mathbf{r}'| = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:

$$\begin{aligned} |\mathbf{r}(t)| &= \text{constant} \\ |\mathbf{r}(t)|' &= 0 \\ \mathbf{r}(t) \cdot \mathbf{r}'(t) &= 0 \end{aligned}$$

# Application

$$|\mathbf{r}'| = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:

$$\begin{aligned} |\mathbf{r}(t)| &= \text{constant} \\ |\mathbf{r}(t)|' &= 0 \\ \mathbf{r}(t) \cdot \mathbf{r}'(t) &= 0 \end{aligned}$$

- In other words,  $\mathbf{r}(t) = \text{constant}$  implies  $\mathbf{r}(t) \perp \mathbf{r}'(t)$ .

# Application

$$|\mathbf{r}|' = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:

$$\begin{aligned} |\mathbf{r}(t)| &= \text{constant} \\ |\mathbf{r}(t)|' &= 0 \\ \mathbf{r}(t) \cdot \mathbf{r}'(t) &= 0 \end{aligned}$$

- In other words,  $\mathbf{r}(t) = \text{constant}$  implies  $\mathbf{r}(t) \perp \mathbf{r}'(t)$ .
- Velocity  $\perp$  Position  $\iff$  Velocity vector  $\mathbf{r}'(t)$  tangent to sphere.

# Application

$$|\mathbf{r}|' = \frac{\mathbf{r} \cdot \mathbf{r}'}{|\mathbf{r}|}$$

- Suppose a point has a trajectory on a sphere with center at origin.  
Therefore:


$$\begin{aligned} |\mathbf{r}(t)| &= \text{constant} \\ |\mathbf{r}(t)|' &= 0 \\ \mathbf{r}(t) \cdot \mathbf{r}'(t) &= 0 \end{aligned}$$

- In other words,  $|\mathbf{r}(t)| = \text{constant}$  implies  $\mathbf{r}(t) \perp \mathbf{r}'(t)$ .
- Velocity  $\perp$  Position  $\iff$  Velocity vector  $\mathbf{r}'(t)$  tangent to sphere.
- What can we say about constant acceleration?

$$\mathbf{r} \cdot \mathbf{r}' \equiv 0 \implies [\mathbf{r} \cdot \mathbf{r}']' = 0 \iff \mathbf{r}' \cdot \mathbf{r}' + \mathbf{r} \cdot \mathbf{r}'' = 0 \implies \mathbf{r} \cdot \mathbf{r}'' = -|\mathbf{r}'|^2 \leq 0$$

Acceleration vector  $\mathbf{r}''$  points inside the sphere.

## Example




Compute the acceleration vector when traversing the loxodrome curve below.

$$x = \rho \sin(at) \cos(bt)$$

$$y = \rho \sin(at) \sin(bt)$$

$$z = \rho \cos(at)$$

Spherical coordinates:



$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

$$z = \rho \cos \phi$$

# Line Integrals

$$\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$$

- Division  $a = t_0 < t_1 < t_2 < \cdots < t_k < t_{k+1} < \cdots < t_n = b$

# Line Integrals

$$\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$$

- Division  $a = t_0 < t_1 < t_2 < \cdots < t_k < t_{k+1} < \cdots < t_n = b$
- Sample points  $t_k \leq s_k \leq t_{k+1}$



# Line Integrals

$$\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$$

- Division  $a = t_0 < t_1 < t_2 < \cdots < t_k < t_{k+1} < \cdots < t_n = b$
- Sample points  $t_k \leq s_k \leq t_{k+1}$
- Riemann sum:

$$\sum_{k=0}^{n-1} (t_{k+1} - t_k) \mathbf{r}(s_k)$$

# Line Integrals

$$\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$$

- Division  $a = t_0 < t_1 < t_2 < \cdots < t_k < t_{k+1} < \cdots < t_n = b$
- Sample points  $t_k \leq s_k \leq t_{k+1}$
- Riemann sum:

$$\sum_{k=0}^{n-1} (t_{k+1} - t_k) \mathbf{r}(s_k)$$

- Definite integral:

$$\int_{t=a}^{t=b} \mathbf{r}(t) dt = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} (t_{k+1} - t_k) \mathbf{r}(s_k)$$

# Line Integrals

$$\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$$

- Division  $a = t_0 < t_1 < t_2 < \cdots < t_k < t_{k+1} < \cdots < t_n = b$
- Sample points  $t_k \leq s_k \leq t_{k+1}$
- Riemann sum:

$$\sum_{k=0}^{n-1} (t_{k+1} - t_k) \mathbf{r}(s_k)$$

- Definite integral:

$$\int_{t=a}^{t=b} \mathbf{r}(t) dt = \lim_{n \rightarrow \infty} \sum_{k=0}^{n-1} (t_{k+1} - t_k) \mathbf{r}(s_k)$$

- Result: a vector.

# Line Integral Properties

- Component-wise: if  $\mathbf{r}(t) = (x(t), y(t), z(t))$ , then

$$\int_a^b \mathbf{r}(t) \, dt = \left( \int_a^b x(t) \, dt, \int_a^b y(t) \, dt, \int_a^b z(t) \, dt \right)$$

# Line Integral Properties

- Component-wise: if  $\mathbf{r}(t) = (x(t), y(t), z(t))$ , then

$$\int_a^b \mathbf{r}(t) \, dt = \left( \int_a^b x(t) \, dt, \int_a^b y(t) \, dt, \int_a^b z(t) \, dt \right)$$

- Substitution Rule:

$$\int_a^b f'(t) \mathbf{r}(f(t)) \, dt = \int_{f(a)}^{f(b)} \mathbf{r}(\tau) \, d\tau$$

# Line Integral Properties

- Component-wise: if  $\mathbf{r}(t) = (x(t), y(t), z(t))$ , then

$$\int_a^b \mathbf{r}(t) dt = \left( \int_a^b x(t) dt, \int_a^b y(t) dt, \int_a^b z(t) dt \right)$$

- Substitution Rule:

$$\int_a^b f'(t) \mathbf{r}(f(t)) dt = \int_{f(a)}^{f(b)} \mathbf{r}(\tau) d\tau$$

- Fundamental Theorem of Calculus: If

$$\mathbf{u}'(t) \equiv \mathbf{r}(t), \text{ then } \int_{t=a}^{t=b} \mathbf{r}(t) dt = \mathbf{u}(b) - \mathbf{u}(a)$$

# Line Integral Properties

- Component-wise: if  $\mathbf{r}(t) = (x(t), y(t), z(t))$ , then

$$\int_a^b \mathbf{r}(t) dt = \left( \int_a^b x(t) dt, \int_a^b y(t) dt, \int_a^b z(t) dt \right)$$

- Substitution Rule:

$$\int_a^b f'(t) \mathbf{r}(f(t)) dt = \int_{f(a)}^{f(b)} \mathbf{r}(\tau) d\tau$$

- Fundamental Theorem of Calculus: If

$$\mathbf{u}'(t) \equiv \mathbf{r}(t), \text{ then } \int_{t=a}^{t=b} \mathbf{r}(t) dt = \mathbf{u}(b) - \mathbf{u}(a)$$

- Derivative  $\implies$  Total change

$$\mathbf{r}(t) = \mathbf{r}(t_0) + \int_{t_0}^t \mathbf{r}'(\tau) d\tau$$

## Example

An object thrown from initial position  $\mathbf{r}_0$  with initial velocity  $\mathbf{v}_0$ . Describe the trajectory of the object (ignore air resistance).



## Example

An object thrown from initial position  $\mathbf{r}_0$  with initial velocity  $\mathbf{v}_0$ . Describe the trajectory of the object (ignore air resistance).

Total force: gravity:  $m\mathbf{a} = \mathbf{F} = -mg\mathbf{k} \implies \mathbf{a} = -g\mathbf{k} \implies \mathbf{r}''(t) = -g\mathbf{k}$

## Example

An object thrown from initial position  $\mathbf{r}_0$  with initial velocity  $\mathbf{v}_0$ . Describe the trajectory of the object (ignore air resistance).

Total force: gravity:  $m\mathbf{a} = \mathbf{F} = -mg\mathbf{k} \implies \mathbf{a} = -g\mathbf{k} \implies \mathbf{r}''(t) = -g\mathbf{k}$

$$\mathbf{r}'(t) = \mathbf{r}'(0) + \int_0^t -g\mathbf{k} \, d\tau = \mathbf{v}_0 + \left( \int_0^t -g \, d\tau \right) \mathbf{k} = \mathbf{v}_0 - g t \mathbf{k}$$

## Example

An object thrown from initial position  $\mathbf{r}_0$  with initial velocity  $\mathbf{v}_0$ . Describe the trajectory of the object (ignore air resistance).

Total force: gravity:  $m\mathbf{a} = \mathbf{F} = -mg\mathbf{k} \implies \mathbf{a} = -g\mathbf{k} \implies \mathbf{r}''(t) = -g\mathbf{k}$

$$\mathbf{r}'(t) = \mathbf{r}'(0) + \int_0^t -g\mathbf{k} \, d\tau = \mathbf{v}_0 + \left( \int_0^t -g \, d\tau \right) \mathbf{k} = \mathbf{v}_0 - g t \mathbf{k}$$

$$\mathbf{r}(t) = \mathbf{r}(0) + \int_0^t (\mathbf{v}_0 - g\tau\mathbf{k}) \, d\tau = \mathbf{r}_0 + \left( \int_0^t d\tau \right) \mathbf{v}_0 - \left( \int_0^t g\tau \, d\tau \right) \mathbf{k} \implies$$

$$\boxed{\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{v}_0 - \frac{1}{2}gt^2\mathbf{k}}$$

## Example

An object thrown from initial position  $\mathbf{r}_0$  with initial velocity  $\mathbf{v}_0$ . Describe the trajectory of the object (ignore air resistance).

Total force: gravity:  $m\mathbf{a} = \mathbf{F} = -mg\mathbf{k} \implies \mathbf{a} = -g\mathbf{k} \implies \mathbf{r}''(t) = -g\mathbf{k}$

$$\mathbf{r}'(t) = \mathbf{r}'(0) + \int_0^t -g\mathbf{k} \, d\tau = \mathbf{v}_0 + \left( \int_0^t -g \, d\tau \right) \mathbf{k} = \mathbf{v}_0 - g t \mathbf{k}$$

$$\mathbf{r}(t) = \mathbf{r}(0) + \int_0^t (\mathbf{v}_0 - g\tau\mathbf{k}) \, d\tau = \mathbf{r}_0 + \left( \int_0^t d\tau \right) \mathbf{v}_0 - \left( \int_0^t g\tau \, d\tau \right) \mathbf{k} \implies$$

$$\boxed{\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{v}_0 - \frac{1}{2}gt^2\mathbf{k}}$$

Parabola in the plane determined by  $\mathbf{v}_0$  and  $\mathbf{k}$ .

# Arclength

- $\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$ : piecewise smooth function
- Distance traveled = Speed  $\cdot$  Time

$$dL = |\mathbf{r}'(t)| dt \implies L = \int_{t=a}^{t=b} |\mathbf{r}'(t)| dt$$

# Arclength

- $\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$ : piecewise smooth function
- Distance traveled = Speed  $\cdot$  Time

$$dL = |\mathbf{r}'(t)| dt \implies L = \int_{t=a}^{t=b} |\mathbf{r}'(t)| dt$$

- Let  $\Delta s$  = distance traveled along curve in short time  $\Delta t$ :

$$\Delta s \approx \sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2} = \sqrt{\left(\frac{\Delta x}{\Delta t}\right)^2 + \left(\frac{\Delta y}{\Delta t}\right)^2 + \left(\frac{\Delta z}{\Delta t}\right)^2} \Delta t.$$

# Arclength

- $\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$ : piecewise smooth function
- Distance traveled = Speed  $\cdot$  Time

$$dL = |\mathbf{r}'(t)| dt \implies L = \int_{t=a}^{t=b} |\mathbf{r}'(t)| dt$$

- Let  $\Delta s$  = distance traveled along curve in short time  $\Delta t$ :

$$\Delta s \approx \sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2} = \sqrt{\left(\frac{\Delta x}{\Delta t}\right)^2 + \left(\frac{\Delta y}{\Delta t}\right)^2 + \left(\frac{\Delta z}{\Delta t}\right)^2} \Delta t.$$

- Infinitesimal element of arclength ( $\Delta t \rightarrow 0$ ):

$$ds = \sqrt{(x'(t))^2 + (y'(t))^2 + (z'(t))^2} dt = |\mathbf{r}'(t)| dt$$

# Arclength

- $\mathbf{r}: [a, b] \rightarrow \mathbb{R}^3$ : piecewise smooth function
- Distance traveled = Speed  $\cdot$  Time

$$dL = |\mathbf{r}'(t)| dt \implies L = \int_{t=a}^{t=b} |\mathbf{r}'(t)| dt$$

- Let  $\Delta s$  = distance traveled along curve in short time  $\Delta t$ :

$$\Delta s \approx \sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2} = \sqrt{\left(\frac{\Delta x}{\Delta t}\right)^2 + \left(\frac{\Delta y}{\Delta t}\right)^2 + \left(\frac{\Delta z}{\Delta t}\right)^2} \Delta t.$$

- Infinitesimal element of arclength ( $\Delta t \rightarrow 0$ ):

$$ds = \sqrt{(x'(t))^2 + (y'(t))^2 + (z'(t))^2} dt = |\mathbf{r}'(t)| dt$$

- Length of parametrized curve:

$$L = \int_{t=a}^{t=b} ds = \int_a^b |\mathbf{r}'(t)| dt.$$



# Arclength Function

- Fix  $t = a$  as starting point. For  $t \geq a$ , let
- $L(t)$  = distance traveled between  $a$  and  $t$  = length of the piece of the curve corresponding to values of the parameter between  $a$  and  $t$ .

$$L(t) = \int_a^t |\mathbf{r}'(\tau)| \, d\tau$$

# Arclength Function

- Fix  $t = a$  as starting point. For  $t \geq a$ , let
- $L(t)$  = distance traveled between  $a$  and  $t$  = length of the piece of the curve corresponding to values of the parameter between  $a$  and  $t$ .

$$L(t) = \int_a^t |\mathbf{r}'(\tau)| \, d\tau$$


- The function  $L: [a, b] \rightarrow \mathbb{R}$  is called the *arclength function*.

## Example

Let  $\mathbf{r}(t) = (\cos t, \sin t, t)$

- Do you know the name of this curve?
- Find the arclength function.
- Find the length of the segment of the curve given by  $t \in [0, 2\pi]$ .


## Example



Let  $\mathbf{r}(t) = (\cos t, \sin t, t)$  (the curve is called a **helix** - (not a spiral) ).

- Do you know the name of this curve?
- Find the arclength function.
- Find the length of the segment of the curve given by  $t \in [0, 2\pi]$ .

## Example




Let  $\mathbf{r}(t) = (\cos t, \sin t, t)$  (the curve is called a helix - (not a spiral) ).

- Do you know the name of this curve?
- Find the arclength function.
- Find the length of the segment of the curve given by  $t \in [0, 2\pi]$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

## Example




Let  $\mathbf{r}(t) = (\cos t, \sin t, t)$  (the curve is called a helix - (not a spiral) ).

- Do you know the name of this curve?
- Find the arclength function.
- Find the length of the segment of the curve given by  $t \in [0, 2\pi]$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) \\ |\mathbf{r}'(t)| &= \sqrt{2} \ .\end{aligned}$$

## Example




Let  $\mathbf{r}(t) = (\cos t, \sin t, t)$  (the curve is called a helix - (not a spiral) ).

- Do you know the name of this curve?
- Find the arclength function.
- Find the length of the segment of the curve given by  $t \in [0, 2\pi]$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) \\ |\mathbf{r}'(t)| &= \sqrt{2} \\ L(t) &= \int_0^t |\mathbf{r}'(\tau)| d\tau\end{aligned}$$

## Example



Let  $\mathbf{r}(t) = (\cos t, \sin t, t)$  (the curve is called a helix - (not a spiral) ).

- Do you know the name of this curve?
- Find the arclength function.
- Find the length of the segment of the curve given by  $t \in [0, 2\pi]$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) \\ |\mathbf{r}'(t)| &= \sqrt{2} \quad . \\ L(t) &= \int_0^t |\mathbf{r}'(\tau)| d\tau \\ &= t\sqrt{2} \quad .\end{aligned}$$



# Parametrization by Arclength

- $C$ : piecewise smooth parametrized curve joining points  $A$  and  $B$ ;
- $\mathbf{r}: I \rightarrow C$ : parametrization of  $C$ ,

$\mathbf{r}(t)$  = position at time  $t$

Not canonically defined: “depends who is driving”.

# Parametrization by Arclength

- $C$ : piecewise smooth parametrized curve joining points  $A$  and  $B$ ;
- $\mathbf{r}: I \rightarrow C$ : parametrization of  $C$ ,

$\mathbf{r}(t)$  = position at time  $t$

Not canonically defined: “depends who is driving”.

- $\mathbf{p}: [0, L] \rightarrow C$ :

$\mathbf{p}(s)$  = position at distance  $s$  from  $A$  along  $C$

Canonically defined: “distance markers along the road”.

# Parametrization by Arclength

- $C$ : piecewise smooth parametrized curve joining points  $A$  and  $B$ ;
- $\mathbf{r}: I \rightarrow C$ : parametrization of  $C$ ,

$\mathbf{r}(t)$  = position at time  $t$

Not canonically defined: “depends who is driving”.

- $\mathbf{p}: [0, L] \rightarrow C$ :

$\mathbf{p}(s)$  = position at distance  $s$  from  $A$  along  $C$

Canonically defined: “distance markers along the road”.

- $\mathbf{p}$ : parametrization by arclength

$$s = L(s) = \int_{\sigma=a}^{\sigma=s} |\mathbf{p}'(\sigma)| \, d\sigma \implies 1 = L'(s) = |\mathbf{p}'(s)| .$$

# Curve Reparametrizations

- Let  $C$  be a piecewise smooth curve joining pts  $A$  and  $B$ .
- Let  $\mathbf{r}: [a, b] \rightarrow C$  be parametrization of  $C$ .
- Let  $\mathbf{p}: [0, L] \rightarrow C$  be arclength parametrization.

# Curve Reparametrizations

- Let  $C$  be a piecewise smooth curve joining pts  $A$  and  $B$ .
- Let  $\mathbf{r}: [a, b] \rightarrow C$  be parametrization of  $C$ .
- Let  $\mathbf{p}: [0, L] \rightarrow C$  be arclength parametrization.
- Question: How do we get  $\mathbf{p}$  from  $\mathbf{r}$ ?

# Curve Reparametrizations

- Let  $C$  be a piecewise smooth curve joining pts  $A$  and  $B$ .
- Let  $\mathbf{r}: [a, b] \rightarrow C$  be parametrization of  $C$ .
- Let  $\mathbf{p}: [0, L] \rightarrow C$  be arclength parametrization.
- Question: How do we get  $\mathbf{p}$  from  $\mathbf{r}$ ?
- $Q = \mathbf{r}(t) = \mathbf{p}(s)$ : point on curve  $C$ .

$$s = \text{distance from } A \text{ to } Q \text{ along } C = \int_{\tau=a}^{\tau=t} |\mathbf{r}'(\tau)| d\tau = \varphi(t)$$

# Curve Reparametrizations

- Let  $C$  be a piecewise smooth curve joining pts  $A$  and  $B$ .
- Let  $\mathbf{r}: [a, b] \rightarrow C$  be parametrization of  $C$ .
- Let  $\mathbf{p}: [0, L] \rightarrow C$  be arclength parametrization.
- Question: How do we get  $\mathbf{p}$  from  $\mathbf{r}$ ?
- $Q = \mathbf{r}(t) = \mathbf{p}(s)$ : point on curve  $C$ .

$$s = \text{distance from } A \text{ to } Q \text{ along } C = \int_{\tau=a}^{\tau=t} |\mathbf{r}'(\tau)| d\tau = \varphi(t)$$

- $\varphi$  invertible and  $\varphi^{-1}$  smooth  $\iff \varphi'(t) \neq 0 \iff \mathbf{r}'(t) \neq \mathbf{0}$ .

# Curve Reparametrizations

- Let  $C$  be a piecewise smooth curve joining pts  $A$  and  $B$ .
- Let  $\mathbf{r}: [a, b] \rightarrow C$  be parametrization of  $C$ .
- Let  $\mathbf{p}: [0, L] \rightarrow C$  be arclength parametrization.
- Question: How do we get  $\mathbf{p}$  from  $\mathbf{r}$ ?
- $Q = \mathbf{r}(t) = \mathbf{p}(s)$ : point on curve  $C$ .

$$s = \text{distance from } A \text{ to } Q \text{ along } C = \int_{\tau=a}^{\tau=t} |\mathbf{r}'(\tau)| d\tau = \varphi(t)$$

- $\varphi$  invertible and  $\varphi^{-1}$  smooth  $\iff \varphi'(t) \neq 0 \iff \mathbf{r}'(t) \neq \mathbf{0}$ .
- Regular parametrization:  $\mathbf{r}'(t) \neq \mathbf{0}$  for all  $t$



# Curve Reparametrizations

- Let  $C$  be a piecewise smooth curve joining pts  $A$  and  $B$ .
- Let  $\mathbf{r}: [a, b] \rightarrow C$  be parametrization of  $C$ .
- Let  $\mathbf{p}: [0, L] \rightarrow C$  be arclength parametrization.
- Question: How do we get  $\mathbf{p}$  from  $\mathbf{r}$ ?
- $Q = \mathbf{r}(t) = \mathbf{p}(s)$ : point on curve  $C$ .

$$s = \text{distance from } A \text{ to } Q \text{ along } C = \int_{\tau=a}^{\tau=t} |\mathbf{r}'(\tau)| d\tau = \varphi(t)$$

- $\varphi$  invertible and  $\varphi^{-1}$  smooth  $\iff \varphi'(t) \neq 0 \iff \mathbf{r}'(t) \neq \mathbf{0}$ .
- Regular parametrization:  $\mathbf{r}'(t) \neq \mathbf{0}$  for all  $t$
- $t = \varphi^{-1}(s) \implies \mathbf{p}(s) = \mathbf{r}(\varphi^{-1}(s))$ .

## Example

Reparametrize  $\mathbf{r}(t) = (\cos t, \sin t, t)$  via arclength.

## Example

Reparametrize  $\mathbf{r}(t) = (\cos t, \sin t, t)$  via arclength.

- $|\mathbf{r}'(t)| = \sqrt{2} \implies$

$$\varphi(t) = \int_{\tau=0}^{\tau=t} |\mathbf{r}'(\tau)| \, d\tau = t\sqrt{2}$$

## Example

Reparametrize  $\mathbf{r}(t) = (\cos t, \sin t, t)$  via arclength.

- $|\mathbf{r}'(t)| = \sqrt{2} \implies$

$$\varphi(t) = \int_{\tau=0}^{\tau=t} |\mathbf{r}'(\tau)| d\tau = t\sqrt{2}$$

- $\varphi^{-1}(s) = \frac{s}{\sqrt{2}}$

$$\mathbf{p}(s) = \left( \cos \left( \frac{s}{\sqrt{2}} \right), \sin \left( \frac{s}{\sqrt{2}} \right), \left( \frac{s}{\sqrt{2}} \right) \right)$$

is a parametrization by arclength.

# Curvature

- Question: How can we *measure* how bent a curve is?

# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.

# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.
- $\mathbf{p} = \mathbf{p}(s)$ : parametrization by arclength of smooth curve  $C$ ;

# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.
- $\mathbf{p} = \mathbf{p}(s)$ : parametrization by arclength of smooth curve  $C$ ;
- $\mathbf{T}(s)$ : unit tangent vector.



# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.
- $\mathbf{p} = \mathbf{p}(s)$ : parametrization by arclength of smooth curve  $C$ ;
- $\mathbf{T}(s)$ : unit tangent vector.
- $\mathbf{v}$ : fixed direction (unit vector);  $\alpha(s)$ : angle between  $\mathbf{T}(s)$  and  $\mathbf{v}$ .

# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.
- $\mathbf{p} = \mathbf{p}(s)$ : parametrization by arclength of smooth curve  $C$ ;
- $\mathbf{T}(s)$ : unit tangent vector.
- $\mathbf{v}$ : fixed direction (unit vector);  $\alpha(s)$ : angle between  $\mathbf{T}(s)$  and  $\mathbf{v}$ .
- Define the *curvature* of  $C$  at  $\mathbf{p}(s)$  to be

$$\kappa(s) = |\alpha'(s)| .$$

# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.
- $\mathbf{p} = \mathbf{p}(s)$ : parametrization by arclength of smooth curve  $C$ ;
- $\mathbf{T}(s)$ : unit tangent vector.
- $\mathbf{v}$ : fixed direction (unit vector);  $\alpha(s)$ : angle between  $\mathbf{T}(s)$  and  $\mathbf{v}$ .
- Define the *curvature* of  $C$  at  $\mathbf{p}(s)$  to be

$$\kappa(s) = |\alpha'(s)| .$$

- Alternative formulas:

$$\kappa(s) = |\mathbf{T}'(s)| = \left| \frac{d\mathbf{T}}{ds} \right| .$$

# Curvature

- Question: How can we *measure* how bent a curve is?
- Answer: Measure change in tangent direction with respect to arclength.
- $\mathbf{p} = \mathbf{p}(s)$ : parametrization by arclength of smooth curve  $C$ ;
- $\mathbf{T}(s)$ : unit tangent vector.
- $\mathbf{v}$ : fixed direction (unit vector);  $\alpha(s)$ : angle between  $\mathbf{T}(s)$  and  $\mathbf{v}$ .
- Define the *curvature* of  $C$  at  $\mathbf{p}(s)$  to be

$$\kappa(s) = |\alpha'(s)| .$$

- Alternative formulas:

$$\kappa(s) = |\mathbf{T}'(s)| = \left| \frac{d\mathbf{T}}{ds} \right| .$$

- $\mathbf{r} = \mathbf{r}(t)$ : smooth parametrization, not necessary by arclength.

$$\kappa(t) = \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3} .$$

$$\kappa = \left| \frac{d\mathbf{T}}{ds} \right| \Rightarrow \kappa = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3}$$

$s(t) = \int_{t_0}^t |\mathbf{r}'(x)| dx$  - curve (arc) length function.


$$|\mathbf{v}|' = (\sqrt{\mathbf{v} \cdot \mathbf{v}})' = \frac{(\mathbf{v} \cdot \mathbf{v})'}{2\sqrt{\mathbf{v} \cdot \mathbf{v}}} = \frac{2\mathbf{v}' \cdot \mathbf{v}}{2|\mathbf{v}|} = \frac{\mathbf{v}' \cdot \mathbf{v}}{|\mathbf{v}|} \cdot \frac{ds}{dt} = |\mathbf{r}'(t)| \frac{d\mathbf{T}}{ds} \frac{ds}{dt} = \frac{d\mathbf{T}}{dt}$$

$$\begin{aligned} \kappa &= \left| \frac{d\mathbf{T}}{ds} \right| = \left| \frac{\frac{d\mathbf{T}}{dt}}{\frac{ds}{dt}} \right| = \left| \frac{\mathbf{T}'(t)}{|\mathbf{r}'(t)|} \right| = \frac{\left| \left( \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|} \right)' \right|}{|\mathbf{r}'(t)|} = \frac{\frac{|\mathbf{r}''|\mathbf{r}' - \mathbf{r}'(|\mathbf{r}'|)'}{|\mathbf{r}'|^2}}{|\mathbf{r}'|} \\ &= \frac{|\mathbf{r}''|\mathbf{r}' - \mathbf{r}' \frac{\mathbf{r}'' \cdot \mathbf{r}'}{|\mathbf{r}'|}}{|\mathbf{r}'|^3} = \frac{\sqrt{(\mathbf{r}''|\mathbf{r}' - \mathbf{r}' \frac{\mathbf{r}'' \cdot \mathbf{r}'}{|\mathbf{r}'|}) \cdot (\mathbf{r}''|\mathbf{r}' - \mathbf{r}' \frac{\mathbf{r}'' \cdot \mathbf{r}'}{|\mathbf{r}'|})}}{|\mathbf{r}'|^3} \\ &= \frac{\sqrt{|\mathbf{r}''|^2 |\mathbf{r}'|^2 - 2\mathbf{r}'' \cdot \mathbf{r}' \frac{\mathbf{r}'' \cdot \mathbf{r}'}{|\mathbf{r}'|} + \frac{|\mathbf{r}'|^2 (\mathbf{r}'' \cdot \mathbf{r}')^2}{|\mathbf{r}'|^2}}{|\mathbf{r}'|^3} = \frac{\sqrt{|\mathbf{r}''|^2 |\mathbf{r}'|^2 - (\mathbf{r}'' \cdot \mathbf{r}')^2}}{|\mathbf{r}'|^3} \\ &= \frac{\sqrt{|\mathbf{r}''|^2 |\mathbf{r}'|^2 - |\mathbf{r}''|^2 |\mathbf{r}'|^2 \cos^2 \alpha}}{|\mathbf{r}'|^3} = \frac{\sqrt{|\mathbf{r}''|^2 |\mathbf{r}'|^2 \sin^2 \alpha}}{|\mathbf{r}'|^3} = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3} \end{aligned}$$

Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .




Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) =$$




Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$






Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$|\mathbf{r}'(t)| =$$




Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.


## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$\mathbf{T}(t) =$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.


## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.


## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1) \quad \mathbf{T}'(t) =$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example


Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$\mathbf{T}'(t) = \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .


$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$\mathbf{T}'(t) = \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$|\mathbf{T}'| =$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .


$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$

$$\mathbf{T}'(t) = \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0)$$

$$|\mathbf{T}'| = \frac{1}{\sqrt{2}}$$





Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$


$$\mathbf{T}'(t) = \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0)$$

$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$|\mathbf{T}'| = \frac{1}{\sqrt{2}}$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$

$$\kappa(t) =$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\mathbf{r}'(t) = (-\sin t, \cos t, 1)$$


$$|\mathbf{r}'(t)| = \sqrt{2}$$

$$\mathbf{T}(t) = \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1)$$

$$\mathbf{T}'(t) = \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0)$$

$$|\mathbf{T}'| = \frac{1}{\sqrt{2}}$$

$$\kappa(t) = \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2}.$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.


## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned} \mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2} \end{aligned}$$

Alternatively:

$$\mathbf{r}'' =$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.


## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2}.\end{aligned}$$

Alternatively:

$$\mathbf{r}'' = (-\cos t, -\sin t, 0)$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example


Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned} \mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2} \end{aligned}$$

Alternatively:

$$\mathbf{r}'' = (-\cos t, -\sin t, 0)$$

$$\mathbf{r}'' \times \mathbf{r}' =$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.


## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2} .\end{aligned}$$

Alternatively:

$$\begin{aligned}\mathbf{r}'' &= (-\cos t, -\sin t, 0) \\ \mathbf{r}'' \times \mathbf{r}' &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\cos t & -\sin t & 0 \\ -\sin t & \cos t & 1 \end{vmatrix} \\ &= -\sin t \mathbf{i} + \cos t \mathbf{j} - \mathbf{k}\end{aligned}$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example


Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned} \mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2} \end{aligned}$$

Alternatively:

$$\begin{aligned} \mathbf{r}'' &= (-\cos t, -\sin t, 0) \\ \mathbf{r}'' \times \mathbf{r}' &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\cos t & -\sin t & 0 \\ -\sin t & \cos t & 1 \end{vmatrix} \\ &= -\sin t \mathbf{i} + \cos t \mathbf{j} - \mathbf{k} \end{aligned}$$

$$\kappa(t) =$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example


Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2}.\end{aligned}$$

Alternatively:

$$\begin{aligned}\mathbf{r}'' &= (-\cos t, -\sin t, 0) \\ \mathbf{r}'' \times \mathbf{r}' &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\cos t & -\sin t & 0 \\ -\sin t & \cos t & 1 \end{vmatrix} \\ &= -\sin t \mathbf{i} + \cos t \mathbf{j} - \mathbf{k}\end{aligned}$$

$$\kappa(t) = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3} = \frac{\sqrt{2}}{(\sqrt{2})^3} = \frac{1}{2}.$$





Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example


Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{\sqrt{2}}{1} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2} .\end{aligned}$$

Alternatively:

$$\begin{aligned}\mathbf{r}'' &= (-\cos t, -\sin t, 0) \\ \mathbf{r}'' \times \mathbf{r}' &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\cos t & -\sin t & 0 \\ -\sin t & \cos t & 1 \end{vmatrix} \\ &= -\sin t \mathbf{i} + \cos t \mathbf{j} - \mathbf{k} \end{aligned}$$

$$\kappa(t) = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3} = \frac{\sqrt{2}}{(\sqrt{2})^3} = \frac{1}{2} .$$



Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example


Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{array}{ll} \mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{1}{\sqrt{2}} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2} \end{array}$$

Alternatively:

$$\begin{aligned}\mathbf{r}'' &= (-\cos t, -\sin t, 0) \\ \mathbf{r}'' \times \mathbf{r}' &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\cos t & -\sin t & 0 \\ -\sin t & \cos t & 1 \end{vmatrix} \\ &= -\sin t \mathbf{i} + \cos t \mathbf{j} - \mathbf{k}\end{aligned}$$

$$\kappa(t) = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3} = \frac{\sqrt{2}}{(\sqrt{2})^3} = \frac{1}{2}.$$




Notation:  $\mathbf{T}$ - unit tangent vector,  $\mathbf{r}$ - position vector,  $\kappa$ -curvature.

## Example

Compute the curvature of  $\mathbf{r}(t) = (\cos t, \sin t, t)$ .

$$\begin{aligned}\mathbf{r}'(t) &= (-\sin t, \cos t, 1) & \mathbf{T}'(t) &= \frac{1}{\sqrt{2}}(-\cos t, -\sin t, 0) \\ |\mathbf{r}'(t)| &= \sqrt{2} & |\mathbf{T}'| &= \frac{\sqrt{2}}{1} \\ \mathbf{T}(t) &= \frac{1}{\sqrt{2}}(-\sin t, \cos t, 1) & \kappa(t) &= \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{1}{2}.\end{aligned}$$




Alternatively:

$$\begin{aligned}\mathbf{r}'' &= (-\cos t, -\sin t, 0) \\ \mathbf{r}'' \times \mathbf{r}' &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -\cos t & -\sin t & 0 \\ -\sin t & \cos t & 1 \end{vmatrix} \\ &= -\sin t \mathbf{i} + \cos t \mathbf{j} - \mathbf{k}\end{aligned}$$

$$\kappa(t) = \frac{|\mathbf{r}''(t) \times \mathbf{r}'(t)|}{|\mathbf{r}'(t)|^3} = \frac{\sqrt{2}}{(\sqrt{2})^3} = \frac{1}{2}.$$

## Example




Compute the curvature of the (torus) trefoil curve

$$x = (R + r \sin(3t)) \cos(2t)$$

$$y = (R + r \sin(3t)) \sin(2t)$$

$$z = r \cos(3t)$$

## Example




Compute the curvature of the (torus)  
trefoil curve

$$x = (R + r \sin(3t)) \cos(2t)$$

$$y = (R + r \sin(3t)) \sin(2t)$$

$$z = r \cos(3t)$$

# Example




Compute the curvature of the loxodrome

$$x = \rho \sin(at) \cos(bt)$$

$$y = \rho \sin(at) \sin(bt)$$

$$z = \rho \cos(at).$$

Spherical coordinates:



$$x = \rho \sin \phi \cos \theta$$

$$y = \rho \sin \phi \sin \theta$$

$$z = \rho \cos \phi$$

# Components of Acceleration

- Object moves through space,  $\mathbf{r} = \mathbf{r}(t)$  position vector at time  $t$ ;
- Velocity vector  $\mathbf{v}(t) = \mathbf{r}'(t)$ ;
- Tangent direction:  $\mathbf{T}(t) = \frac{\mathbf{v}(t)}{|\mathbf{v}(t)|}$ ;
- Speed is  $v(t) = |\mathbf{v}(t)|$ ;
- Acceleration  $\mathbf{a}(t) = \mathbf{r}''(t)$ ;
- Tangential component  $\mathbf{a}_T(t)$ :

$$\mathbf{a}_T(t) = \text{proj}_{\mathbf{T}(t)} \mathbf{a}(t) = \frac{\mathbf{a} \cdot \mathbf{T}}{|\mathbf{T}|} \mathbf{T} = \frac{\mathbf{v}' \cdot \mathbf{v}}{|\mathbf{v}|} \mathbf{T} = |\mathbf{v}'| \mathbf{T} = v' \mathbf{T},$$

$$a_T(t) = |\mathbf{a}_T(t)| = |v'(t)|.$$

- Normal component  $\mathbf{a}_N(t) = \text{orth}_{\mathbf{T}(t)} \mathbf{a}(t)$ ,

$$a_N(t) = |\mathbf{a}_N(t)| = |\text{orth}_{\mathbf{T}} \mathbf{a}| = |\mathbf{a} \times \mathbf{T}| = \frac{|\mathbf{r}'' \times \mathbf{r}'|}{|\mathbf{r}'|} = \kappa |\mathbf{r}'|^2 = \kappa(t) v^2(t).$$