

Calculus I

Lecture 20

Areas, Integration

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

- 1 Areas and Distances
 - The Area Problem

Outline

- 1 Areas and Distances
 - The Area Problem

- 2 The Definite Integral
 - Review of the \sum notation
 - Riemann sums, areas and integrals

Outline

1 Areas and Distances

- The Area Problem

2 The Definite Integral

- Review of the \sum notation
- Riemann sums, areas and integrals

3 The Definite Integral

- Evaluating Integrals with Riemann Sums
- Properties of the Definite Integral

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.


The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.





The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.





The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.





The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.





The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.





The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.




The Area Problem

- How can we find the area under $y = x^2$ between $x = 0$ and $x = 1$?
- We can approximate it using rectangles.
- Divide $[0, 1]$ into three strips of width $\frac{1}{3}$, and draw rectangles in those strips, the heights of which are the same as the height of the function at the right end of that strip.
- Four strips gives a better approximation. Five is even better.
- We could use the left endpoints to find the heights instead.



Example


Find the sum of the areas of the four approximating rectangles obtained using right endpoints.



Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.


- Let R_4 denote the sum of the areas of the rectangles.



Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.


- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.



Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.


- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.



Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.


- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are
 $?$, $?$, $?$, and $?$.



Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.


- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are $\left(\frac{1}{4}\right)^2$, $\left(\frac{1}{2}\right)^2$, $\left(\frac{3}{4}\right)^2$, and 1^2 .



Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.

- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are $\left(\frac{1}{4}\right)^2$, $\left(\frac{1}{2}\right)^2$, $\left(\frac{3}{4}\right)^2$, and 1^2 .




$$R_4 = \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 + \frac{1}{4} \cdot (1)^2$$

Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.

- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are $\left(\frac{1}{4}\right)^2$, $\left(\frac{1}{2}\right)^2$, $\left(\frac{3}{4}\right)^2$, and 1^2 .




$$R_4 = \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 + \frac{1}{4} \cdot (1)^2 = \frac{15}{32} = 0.46875$$

Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.

- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are $\left(\frac{1}{4}\right)^2$, $\left(\frac{1}{2}\right)^2$, $\left(\frac{3}{4}\right)^2$, and 1^2 .
- A similar calculation works for L_4 , the sum of the areas of the left endpoint rectangles.




$$R_4 = \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 + \frac{1}{4} \cdot (1)^2 = \frac{15}{32} = 0.46875$$

$$L_4 = \frac{1}{4} \cdot (0)^2 + \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2$$

Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.

- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are $\left(\frac{1}{4}\right)^2$, $\left(\frac{1}{2}\right)^2$, $\left(\frac{3}{4}\right)^2$, and 1^2 .
- A similar calculation works for L_4 , the sum of the areas of the left endpoint rectangles.




$$R_4 = \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 + \frac{1}{4} \cdot (1)^2 = \frac{15}{32} = 0.46875$$

$$L_4 = \frac{1}{4} \cdot (0)^2 + \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2$$

Example

Find the sum of the areas of the four approximating rectangles obtained using right endpoints.

- Let R_4 denote the sum of the areas of the rectangles.
- Each rectangle has width $\frac{1}{4}$.
- The heights are $\left(\frac{1}{4}\right)^2$, $\left(\frac{1}{2}\right)^2$, $\left(\frac{3}{4}\right)^2$, and 1^2 .
- A similar calculation works for L_4 , the sum of the areas of the left endpoint rectangles.




$$R_4 = \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 + \frac{1}{4} \cdot (1)^2 = \frac{15}{32} = 0.46875$$

$$L_4 = \frac{1}{4} \cdot (0)^2 + \frac{1}{4} \cdot \left(\frac{1}{4}\right)^2 + \frac{1}{4} \cdot \left(\frac{1}{2}\right)^2 + \frac{1}{4} \cdot \left(\frac{3}{4}\right)^2 = \frac{7}{32} = 0.21875$$

Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,


$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$



Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$




Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width ?.
- The heights are ? , ? , ..., ? .




Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $?$, $?$, \dots , $?$.




Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $?$, $?$, \dots , $?$.




Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.




Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.




$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \cdots + \frac{1}{n} \left(\frac{n}{n}\right)^2$$

Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.



$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \cdots + \frac{1}{n} \left(\frac{n}{n}\right)^2 = \frac{1}{n^3} (1^2 + 2^2 + \cdots + n^2)$$

Example


For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.
- New formula:

$$\bullet \quad 1^2 + 2^2 + 3^2 + \dots + n^2 = ?$$

$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \dots + \frac{1}{n} \left(\frac{n}{n}\right)^2 = \frac{1}{n^3} (1^2 + 2^2 + \dots + n^2)$$




Example

For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.
- New formula:
- $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}.$

$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \dots + \frac{1}{n} \left(\frac{n}{n}\right)^2 = \frac{1}{n^3} \left(1^2 + 2^2 + \dots + n^2\right)$$



Example


For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.
- New formula:
- $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$.

$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \dots + \frac{1}{n} \left(\frac{n}{n}\right)^2 = \frac{1}{n^3} \left(1^2 + 2^2 + \dots + n^2\right)$$

$$\lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6}$$



Example


For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.
- New formula:
- $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$.

$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \dots + \frac{1}{n} \left(\frac{n}{n}\right)^2 = \frac{1}{n^3} (1^2 + 2^2 + \dots + n^2)$$

$$\lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \lim_{n \rightarrow \infty} \frac{1}{6} \left(1 + \frac{1}{n}\right) \left(2 + \frac{1}{n}\right)$$



Example


For the region S underneath the parabola $y = x^2$ from 0 to 1, show that the area under the approximating rectangles approaches $\frac{1}{3}$, that is,

$$\lim_{n \rightarrow \infty} R_n = \frac{1}{3}.$$

- Each rectangle has width $\frac{1}{n}$.
- The heights are $\left(\frac{1}{n}\right)^2, \left(\frac{2}{n}\right)^2, \dots, \left(\frac{n}{n}\right)^2$.
- New formula:
- $1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}$.

$$R_n = \frac{1}{n} \left(\frac{1}{n}\right)^2 + \frac{1}{n} \left(\frac{2}{n}\right)^2 + \dots + \frac{1}{n} \left(\frac{n}{n}\right)^2 = \frac{1}{n^3} (1^2 + 2^2 + \dots + n^2)$$

$$\lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} \frac{1}{n^3} \frac{n(n+1)(2n+1)}{6} = \lim_{n \rightarrow \infty} \frac{1}{6} \left(1 + \frac{1}{n}\right) \left(2 + \frac{1}{n}\right) = \frac{1}{3}$$



Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the \dots notation and using the \sum notation.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the \dots notation and using the \sum notation.

$$A = ?$$

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the \dots notation and using the \sum notation.

$$A = 2 + 4 + 6 + \dots + 124$$

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$A = 2 + 4 + 6 + \dots + 124$$

- We aim to introduce the \sum notation for series via this example.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$A = 2 + 4 + 6 + \dots + 124$$

- We aim to introduce the \sum notation for series via this example.
- The \dots notation is informal but easier to read.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The \dots notation is informal but easier to read.
- If the \dots are too ambiguous, we should include the general term.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \end{aligned}$$

$2n$.

- We aim to introduce the \sum notation for series via this example.
- The \dots notation is informal but easier to read.
- If the \dots are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the **pattern of the general term**.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned}
 A &= 2 + 4 + 6 + \cdots + 124 \\
 &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\
 &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\
 &\quad 2n \quad .
 \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The ... notation is informal but easier to read.
- If the ... are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the **pattern of the general term**.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &\quad 2n \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The ... notation is informal but easier to read.
- If the ... are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the **pattern of the general term**.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned}
 A &= 2 + 4 + 6 + \cdots + 124 \\
 &= 2 + 4 + \textcolor{red}{6} + \cdots + 2n + \cdots + 124 \\
 &= 2 \cdot 1 + 2 \cdot 2 + \textcolor{red}{2} \cdot \textcolor{red}{3} + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\
 &\quad \textcolor{red}{2n} \quad .
 \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The ... notation is informal but easier to read.
- If the ... are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the **pattern of the general term**.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &\quad 2n \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The ... notation is informal but easier to read.
- If the ... are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the **pattern of the general term**.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot n + \dots + 2 \cdot 62 \\ &\quad 2n \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The \dots notation is informal but easier to read.
- If the \dots are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the pattern of the general term.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- We aim to introduce the \sum notation for series via this example.
- The ... notation is informal but easier to read.
- If the ... are too ambiguous, we should include the general term.
- To make it clearer we should rewrite all elements in the pattern of the general term.
- If that is still ambiguous we should switch to the completely unambiguous \sum notation.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot n + \dots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot n + \dots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add **several copies of the summed term**, where in each term the index is replaced by a concrete value.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the **boundaries of summation**.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the **boundaries of summation**.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the boundaries of summation.
- The index varies over all integers starting with the lower boundary and ending with upper boundary.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot n + \dots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the boundaries of summation.
- The index varies over all integers starting with the lower boundary and ending with upper boundary.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the boundaries of summation.
- The index varies over all integers starting with the lower boundary and ending with upper boundary.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the boundaries of summation.
- The index varies over all integers starting with the lower boundary and ending with upper boundary.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot n + \dots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the boundaries of summation.
- The index varies over all integers starting with the lower boundary and ending with upper boundary.

Example (The \dots and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the \dots notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \dots + 124 \\ &= 2 + 4 + 6 + \dots + 2n + \dots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \dots + 2 \cdot n + \dots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- The number n is the index (counter) of the sum.
- \sum tells us to add several copies of the summed term, where in each term the index is replaced by a concrete value.
- The values taken by the index are determined by the boundaries of summation.
- The index varies over all integers starting with the lower boundary and ending with upper boundary.
- In programming, what objects are similar to \sum ?

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- To go from \sum to ... notation: substitute few values for the index.
Make sure to include the last value.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- To go from \sum to ... notation: substitute few values for the index.
Make sure to include the last value.
- To go from ... to \sum notation:
 - figure out a pattern for the general term just as with sequences;

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- To go from \sum to ... notation: substitute few values for the index.
Make sure to include the last value.
- To go from ... to \sum notation:
 - figure out a pattern for the general term just as with sequences;
 - select first and last index so that your general term formula reproduces the first and last terms of the sequence.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- Bear in mind the ... notation is informal.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- Bear in mind the ... notation is informal.
 - There are infinitely many formulas that fit any single pattern.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- Bear in mind the ... notation is informal.
 - There are infinitely many formulas that fit any single pattern.
 - Thus it is acceptable to use the ... notation only when we believe there is a single completely obvious pattern that will be recognized by every one.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124. Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- Bear in mind the ... notation is informal.
 - There are infinitely many formulas that fit any single pattern.
 - Thus it is acceptable to use the ... notation only when we believe there is a single completely obvious pattern that will be recognized by every one.
 - The pattern should be obvious not only to us, but also to our potential readers.

Example (The ... and \sum notations for series)

Let A be the sum of the positive even integers between 2 and 124.
Write A using the ... notation and using the \sum notation.

$$\begin{aligned} A &= 2 + 4 + 6 + \cdots + 124 \\ &= 2 + 4 + 6 + \cdots + 2n + \cdots + 124 \\ &= 2 \cdot 1 + 2 \cdot 2 + 2 \cdot 3 + \cdots + 2 \cdot n + \cdots + 2 \cdot 62 \\ &= \sum_{n=1}^{62} 2n . \end{aligned}$$

- Bear in mind the ... notation is informal.
 - There are infinitely many formulas that fit any single pattern.
 - Thus it is acceptable to use the ... notation only when we believe there is a single completely obvious pattern that will be recognized by every one.
 - The pattern should be obvious not only to us, but also to our potential readers.
 - If in doubt or seeking complete rigor we should use the \sum notation.

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.

NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.

NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Example

$$\sum_{j=3}^7 j^2 = ? + ? + ?$$

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.

NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Example

$$\sum_{j=3}^7 j^2 = 9 + ? + ?$$

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.

NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Example

$$\sum_{j=3}^7 j^2 = 9 + ? + ?$$

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.

NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Example

$$\sum_{j=3}^7 j^2 = 9 + 16 + ?$$

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.

NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Example

$$\sum_{j=3}^7 j^2 = 9 + 16 + ?$$

Definition

Sigma Notation: The sum of n terms a_1, a_2, \dots, a_n is written as

$$\sum_{i=1}^n a_i = a_1 + a_2 + \cdots + a_n$$

where i is the *index of summation*, a_i is the i 'th term, and the *upper and lower bounds of summation* are n and 1 respectively.


NOTE: The lower bound doesn't have to be 1. Any integer less than or equal to the upper bound is legitimate.

The index i may be replaced with another symbol, often j or k .

Example

$$\sum_{j=3}^7 j^2 = 9 + 16 + 25 + 36 + 49$$

Estimate the area under $y = f(x)$ between a and b using n strips.



- The right endpoints of the subintervals are

$$x_1 = a + \Delta x$$

$$x_2 = a + 2\Delta x$$


$$x_3 = a + 3\Delta x$$

$$\vdots$$

- The width of the interval is $b - a$.
- The width one strip is $\Delta x = \frac{b-a}{n}$.
- $[a, b]$ is divided into n subintervals: $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$, where $x_0 = a$ and $x_n = b$.
- The height of the i th rectangle is $f(x_i)$.
- The area of the i th rectangle is $f(x_i)\Delta x$.

$$R_n = f(x_1)\Delta x + f(x_2)\Delta x + f(x_3)\Delta x + \dots + f(x_n)\Delta x$$

Estimate the area under $y = f(x)$ between a and b using n strips.



- The **left** endpoints of the subintervals are

$$x_0 = a$$

$$x_1 = a + \Delta x$$

$$x_2 = a + 2\Delta x$$

$$\vdots$$

- The width of the interval is $b - a$.
- The width one strip is $\Delta x = \frac{b-a}{n}$.
- $[a, b]$ is divided into n subintervals: $[x_0, x_1], [x_1, x_2], \dots, [x_{n-1}, x_n]$, where $x_0 = a$ and $x_n = b$.
- The height of the i th rectangle is $f(x_{i-1})$.
- The area of the i th rectangle is $f(x_{i-1})\Delta x$.

$$L_n = f(x_0)\Delta x + f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_{n-1})\Delta x$$

Definition (Area Under a Curve)

Let $f(x) > 0$. The area of the region S that lies under $y = f(x)$ is the limit of the sum of the areas of the approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1)\Delta x + f(x_2)\Delta x + \cdots + f(x_n)\Delta x]$$

Definition (Area Under a Curve)

Let $f(x) > 0$. The area of the region S that lies under $y = f(x)$ is the limit of the sum of the areas of the approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1)\Delta x + f(x_2)\Delta x + \cdots + f(x_n)\Delta x]$$

- This limit always exists if f is continuous.

Definition (Area Under a Curve)

Let $f(x) > 0$. The area of the region S that lies under $y = f(x)$ is the limit of the sum of the areas of the approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1)\Delta x + f(x_2)\Delta x + \cdots + f(x_n)\Delta x]$$

- This limit always exists if f is continuous.
- If f is continuous, we get the same limit if we use left endpoints:

$$A = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} [f(x_0)\Delta x + f(x_1)\Delta x + \cdots + f(x_{n-1})\Delta x]$$

Definition (Area Under a Curve)

Let $f(x) > 0$. The area of the region S that lies under $y = f(x)$ is the limit of the sum of the areas of the approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1)\Delta x + f(x_2)\Delta x + \cdots + f(x_n)\Delta x]$$

- This limit always exists if f is continuous.
- If f is continuous, we get the same limit if we use left endpoints:

$$A = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} [f(x_0)\Delta x + f(x_1)\Delta x + \cdots + f(x_{n-1})\Delta x]$$

- If f is continuous, we get the same limit if we use any number x_i^* in the interval $[x_{i-1}, x_i]$. x_i^* is called a sample point.

$$A = \lim_{n \rightarrow \infty} [f(x_1^*)\Delta x + f(x_2^*)\Delta x + \cdots + f(x_n^*)\Delta x]$$

Definition (Area Under a Curve)

Let $f(x) > 0$. The area of the region S that lies under $y = f(x)$ is the limit of the sum of the areas of the approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} [f(x_1)\Delta x + f(x_2)\Delta x + \cdots + f(x_n)\Delta x]$$

- This limit always exists if f is continuous.
- If f is continuous, we get the same limit if we use left endpoints:

$$A = \lim_{n \rightarrow \infty} L_n = \lim_{n \rightarrow \infty} [f(x_0)\Delta x + f(x_1)\Delta x + \cdots + f(x_{n-1})\Delta x]$$

- If f is continuous, we get the same limit if we use any number x_i^* in the interval $[x_{i-1}, x_i]$. x_i^* is called a sample point.

$$A = \lim_{n \rightarrow \infty} [f(x_1^*)\Delta x + f(x_2^*)\Delta x + \cdots + f(x_n^*)\Delta x]$$

Definition (Riemann Sum)

A Riemann sum is any sum of the form

$$f(x_1^*)\Delta x + f(x_2^*)\Delta x + \cdots + f(x_n^*)\Delta x.$$

The Definite Integral

Definition (Definite Integral)

- Let f be a function defined for $a \leq x \leq b$.
- Divide the interval $[a, b]$ into n subintervals of equal width $\Delta x = (b - a)/n$ and set $x_0 = a$, $x_n = b$.
- Let x_0, x_1, \dots, x_n be the endpoints of the subintervals.
- Let $x_1^*, x_2^*, \dots, x_n^*$ be any sample points in these subintervals; that is, x_i^* is in $[x_{i-1}, x_i]$.

Suppose the limit $\lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x$ exists and is independent of the choice of sample points x_i^* . Then we say that f is an integrable function. If f is integrable we call the limit the integral of f over $[a, b]$ and write

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x \quad .$$

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*)\Delta x,$$

- \int is called the integration sign.
- $f(x)$ is called the integrand.
- a and b are called the limits of integration.

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*)\Delta x,$$

- \int is called the integration sign.
- $f(x)$ is called the integrand.
- a and b are called the limits of integration.

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x,$$

- \int is called the integration sign.
- $f(x)$ is called the integrand.
- a and b are called the limits of integration.

$$\int\limits_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x,$$


- \int is called the integration sign.
- $f(x)$ is called the integrand.
- a and b are called the limits of integration.

$$\int_a^b f(x)dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i^*) \Delta x,$$


- \int is called the integration sign.
- $f(x)$ is called the integrand.
- a and b are called the limits of integration.
- The definite integral is a number. It does not depend on x . We could use any variable instead of x .

$$\int_a^b f(x)dx = \int_a^b f(t)dt = \int_a^b f(r)dr = \int_a^b f(\theta)d\theta$$


- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.




- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.




- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.




- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.




- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.




- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.




- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.



- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.





- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.



- What if $f(x)$ is sometimes negative?

- We know already that if $f(x)$ is always positive, then $\int_a^b f(x)dx$ is the area under the curve.



- What if $f(x)$ is sometimes negative?
- Then $\int_a^b f(x)dx = A_1 - A_2$.
- A_1 is the area of the region above the x-axis and below the graph of f .
- A_2 is the area of the region below the x-axis and above the graph of f .

Theorem

Let f be a continuous function on $[a, b]$. Then f is integrable over $[a, b]$.

- In particular the integral does not depend the choice of sampling points so long as the intervals containing them shrink.
- The proof of this theorem is not difficult, but is outside of the scope of Calculus I and II.
- The only “difficulty” in the proof stems from the fact that we have not rigorously constructed the real numbers.
- We already (silently) assumed a construction of the real numbers when we defined limits.
- Such a construction is also (silently) assumed in most regular high school mathematics courses.
- A student interested in a proof of the theorem should google “Darboux integral”.

The following **power sums** will be useful in what follows:

$$① \sum_{i=1}^n i = \frac{n(n+1)}{2}$$

The following **power sums** will be useful in what follows:

$$① \sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$② \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

The following **power sums** will be useful in what follows:


$$① \sum_{i=1}^n i = \frac{n(n+1)}{2}$$

$$② \sum_{i=1}^n i^2 = \frac{n(n+1)(2n+1)}{6}$$

$$③ \sum_{i=1}^n i^3 = \left[\frac{n(n+1)}{2} \right]^2$$


Example

Evaluate $\int_0^3 (x^3 - 6x) dx$.




Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} =$



Example


Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f \left(\quad \right)$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\right) \frac{3}{n}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\right) \frac{3}{n}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.

$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(\underline{x}_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n}$$




Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.

$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n}$$


$$= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6\left(\frac{3i}{n}\right) \right]$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\ &= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right]\end{aligned}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.

$$\begin{aligned}\int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\&= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right] \\&= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right]\end{aligned}$$



Example


Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.

$$\int_0^3 (x^3 - 6x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n}$$

$$= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right]$$

$$= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right]$$


$$= \lim_{n \rightarrow \infty} \left(\frac{81}{n^4} - \frac{54}{n^2} \right)$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\begin{aligned}
 \int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\
 &= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right] \\
 &= \lim_{n \rightarrow \infty} \left(\frac{81}{n^4} \left[\frac{n(n+1)}{2} \right]^2 - \frac{54}{n^2} \right)
 \end{aligned}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\begin{aligned}
 \int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\
 &= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right] \\
 &= \lim_{n \rightarrow \infty} \left(\frac{81}{n^4} \left[\frac{n(n+1)}{2} \right]^2 - \frac{54}{n^2} \right)
 \end{aligned}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\begin{aligned}
 \int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\
 &= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right] \\
 &= \lim_{n \rightarrow \infty} \left(\frac{81}{n^4} \left[\frac{n(n+1)}{2} \right]^2 - \frac{54}{n^2} \frac{n(n+1)}{2} \right)
 \end{aligned}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.


$$\begin{aligned}
 \int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\
 &= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right] \\
 &= \lim_{n \rightarrow \infty} \left(\frac{81}{n^4} \left[\frac{n(n+1)}{2} \right]^2 - \frac{54}{n^2} \frac{n(n+1)}{2} \right) \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{4} \left(1 + \frac{1}{n} \right)^2 - 27 \left(1 + \frac{1}{n} \right) \right]
 \end{aligned}$$



Example

Evaluate $\int_0^3 (x^3 - 6x) dx$. $\Delta x = \frac{b-a}{n} = \frac{3}{n}$.

$$\begin{aligned}
 \int_0^3 (x^3 - 6x) dx &= \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x = \lim_{n \rightarrow \infty} \sum_{i=1}^n f\left(\frac{3i}{n}\right) \frac{3}{n} \\
 &= \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\left(\frac{3i}{n}\right)^3 - 6 \left(\frac{3i}{n}\right) \right] = \lim_{n \rightarrow \infty} \frac{3}{n} \sum_{i=1}^n \left[\frac{27}{n^3} i^3 - \frac{18}{n} i \right] \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{n^4} \sum_{i=1}^n i^3 - \frac{54}{n^2} \sum_{i=1}^n i \right] \\
 &= \lim_{n \rightarrow \infty} \left(\frac{81}{n^4} \left[\frac{n(n+1)}{2} \right]^2 - \frac{54}{n^2} \frac{n(n+1)}{2} \right) \\
 &= \lim_{n \rightarrow \infty} \left[\frac{81}{4} \left(1 + \frac{1}{n} \right)^2 - 27 \left(1 + \frac{1}{n} \right) \right] = \frac{81}{4} - 27 = -\frac{27}{4}
 \end{aligned}$$



Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

$$\int_b^a f(x)dx =$$

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

$$\int_b^a f(x)dx = - \int_a^b f(x)dx$$

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

$$\int_b^a f(x)dx = - \int_a^b f(x)dx$$

- If $a = b$, then $\Delta x =$

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

$$\int_b^a f(x)dx = - \int_a^b f(x)dx$$

- If $a = b$, then $\Delta x = 0$.

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

$$\int_b^a f(x)dx = - \int_a^b f(x)dx$$

- If $a = b$, then $\Delta x = 0$.

$$\int_a^a f(x)dx =$$

Properties of the Definite Integral

- So far when we have calculated $\int_a^b f(x)dx$, we have assumed that $a < b$.
- The definition as a limit of Riemann sums will still work even if we don't assume this.
- If we reverse a and b , then Δx changes from $\frac{b-a}{n}$ to $\frac{a-b}{n}$.

$$\int_b^a f(x)dx = - \int_a^b f(x)dx$$

- If $a = b$, then $\Delta x = 0$.

$$\int_a^a f(x)dx = 0$$

Properties of the Integral

- ① $\int_a^b c dx = c(b - a)$, where c is any constant.
- ② $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$.
- ③ $\int_a^b c f(x) dx = c \int_a^b f(x) dx$, where c is any constant.
- ④ $\int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$.

Properties of the Integral

- 1 $\int_a^b c dx = c(b - a)$, where c is any constant.
- 2 $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$.
- 3 $\int_a^b c f(x) dx = c \int_a^b f(x) dx$, where c is any constant.
- 4 $\int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$.

Properties of the Integral

- 1 $\int_a^b c dx = c(b - a)$, where c is any constant.
- 2 $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx.$
- 3 $\int_a^b c f(x) dx = c \int_a^b f(x) dx$, where c is any constant.
- 4 $\int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx.$

Properties of the Integral

- 1 $\int_a^b c dx = c(b - a)$, where c is any constant.
- 2 $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$.
- 3 $\int_a^b c f(x) dx = c \int_a^b f(x) dx$, where c is any constant.
- 4 $\int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$.

Properties of the Integral

- ① $\int_a^b c dx = c(b - a)$, where c is any constant.
- ② $\int_a^b [f(x) + g(x)] dx = \int_a^b f(x) dx + \int_a^b g(x) dx$.
- ③ $\int_a^b c f(x) dx = c \int_a^b f(x) dx$, where c is any constant.
- ④ $\int_a^b [f(x) - g(x)] dx = \int_a^b f(x) dx - \int_a^b g(x) dx$.

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

$$= \int_0^1 4 dx + 3 \int_0^1 x^2 dx \quad \text{Property}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

$$= \int_0^1 4 dx + 3 \int_0^1 x^2 dx \quad \text{Property 3}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

$$= \int_0^1 4 dx + 3 \int_0^1 x^2 dx \quad \text{Property 3}$$

$$= \quad + 3 \int_0^1 x^2 dx \quad \text{Property}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

$$= \int_0^1 4 dx + 3 \int_0^1 x^2 dx \quad \text{Property 3}$$

$$= 4(1 - 0) + 3 \int_0^1 x^2 dx \quad \text{Property}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2)dx = \int_0^1 4dx + \int_0^1 3x^2dx \quad \text{Property 2}$$

$$= \int_0^1 4dx + 3 \int_0^1 x^2dx \quad \text{Property 3}$$

$$= 4(1 - 0) + 3 \int_0^1 x^2dx \quad \text{Property 1}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

$$= \int_0^1 4 dx + 3 \int_0^1 x^2 dx \quad \text{Property 3}$$

$$= 4(1 - 0) + 3 \int_0^1 x^2 dx \quad \text{Property 1}$$

$$= 4 + 3 \cdot$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2) dx = \int_0^1 4 dx + \int_0^1 3x^2 dx \quad \text{Property 2}$$

$$= \int_0^1 4 dx + 3 \int_0^1 x^2 dx \quad \text{Property 3}$$

$$= 4(1 - 0) + 3 \int_0^1 x^2 dx \quad \text{Property 1}$$

$$= 4 + 3 \cdot \frac{1}{3} \quad \text{From preceding lectures/slides}$$

Example

Use the properties of integrals to evaluate

$$\int_0^1 (4 + 3x^2)dx = \int_0^1 4dx + \int_0^1 3x^2dx \quad \text{Property 2}$$

$$= \int_0^1 4dx + 3 \int_0^1 x^2dx \quad \text{Property 3}$$


$$= 4(1 - 0) + 3 \int_0^1 x^2dx \quad \text{Property 1}$$

$$= 4 + 3 \cdot \frac{1}{3} \quad \text{From preceding lectures/slides}$$

$$= 5$$


Properties of the Integral

$$⑤ \int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$




Properties of the Integral

$$⑤ \int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$



Properties of the Integral

$$⑤ \int_a^b f(x)dx = \int_a^c f(x)dx + \int_c^b f(x)dx$$



Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\int_0^8 f(x)dx + \int_8^{10} f(x)dx$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\int_0^8 f(x)dx + \int_8^{10} f(x)dx = \int_0^{10} f(x)dx$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\int_0^8 f(x)dx + \int_8^{10} f(x)dx = \int_0^{10} f(x)dx$$

$$\int_8^{10} f(x)dx = \int_0^{10} f(x)dx - \int_0^8 f(x)dx$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\begin{aligned}\int_0^8 f(x)dx + \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx \\ \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx - \int_0^8 f(x)dx \\ &= 17 - 12\end{aligned}$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\begin{aligned}\int_0^8 f(x)dx + \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx \\ \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx - \int_0^8 f(x)dx \\ &= 17 - 12\end{aligned}$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\begin{aligned}\int_0^8 f(x)dx + \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx \\ \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx - \int_0^8 f(x)dx \\ &= 17 - 12\end{aligned}$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.

$$\begin{aligned}\int_0^8 f(x)dx + \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx \\ \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx - \int_0^8 f(x)dx \\ &= 17 - 12\end{aligned}$$

Example

If it is known that $\int_0^{10} f(x)dx = 17$ and $\int_0^8 f(x)dx = 12$, then find $\int_8^{10} f(x)dx$.


$$\begin{aligned}\int_0^8 f(x)dx + \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx \\ \int_8^{10} f(x)dx &= \int_0^{10} f(x)dx - \int_0^8 f(x)dx \\ &= 17 - 12 \\ &= 5\end{aligned}$$

Comparison Properties of the Integral

⑥ If $f(x) \geq 0$ for all $a \leq x \leq b$, then $\int_a^b f(x)dx \geq 0$.

Comparison Properties of the Integral


7 If $f(x) \leq g(x)$ for all $a \leq x \leq b$, then $\int_a^b f(x)dx \leq \int_a^b g(x)dx$.



$$\int_a^b f(x)dx \leq \int_a^b g(x)dx$$

Comparison Properties of the Integral


7 If $f(x) \leq g(x)$ for all $a \leq x \leq b$, then $\int_a^b f(x)dx \leq \int_a^b g(x)dx$.



$$\int_a^b f(x)dx \leq \int_a^b g(x)dx$$

Comparison Properties of the Integral

7 If $f(x) \leq g(x)$ for all $a \leq x \leq b$, then $\int_a^b f(x)dx \leq \int_a^b g(x)dx$.




$$\int_a^b f(x)dx \leq \int_a^b g(x)dx$$

Comparison Properties of the Integral

8 If $m \leq f(x) \leq M$ for all $a \leq x \leq b$, then


$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$



Comparison Properties of the Integral

8 If $m \leq f(x) \leq M$ for all $a \leq x \leq b$, then


$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$



Comparison Properties of the Integral

8 If $m \leq f(x) \leq M$ for all $a \leq x \leq b$, then

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$



Comparison Properties of the Integral

8 If $m \leq f(x) \leq M$ for all $a \leq x \leq b$, then

$$m(b-a) \leq \int_a^b f(x) dx \leq M(b-a)$$

