

Calculus II

Lecture 4

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

- 1 Integration of Rational Functions
 - Building block integrals

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:

<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

- Yes. We will learn how in what follows.

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - We use algebra to split $\frac{P(x)}{Q(x)}$ into smaller pieces (“partial fractions”).

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - We use algebra to split $\frac{P(x)}{Q(x)}$ into smaller pieces (“partial fractions”).
 - We use linear substitutions to transform each piece to one of 3 pairs of basic building block integrals.

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - We use algebra to split $\frac{P(x)}{Q(x)}$ into smaller pieces (“partial fractions”).
 - We use linear substitutions to transform each piece to one of 3 pairs of basic building block integrals.
 - We solve each building block integral and collect the terms.

Integrating arbitrary rational functions

Let $\frac{P(x)}{Q(x)}$ be an arbitrary rational function, i.e., a quotient of polynomials.

Question

Can we integrate $\int \frac{P(x)}{Q(x)} dx$?

- Yes. We will learn how in what follows.
- The algorithm for integration is roughly:
 - We use algebra to split $\frac{P(x)}{Q(x)}$ into smaller pieces (“partial fractions”).
 - We use linear substitutions to transform each piece to one of 3 pairs of basic building block integrals.
 - We solve each building block integral and collect the terms.
- We study the algorithm “from the ground up”: we start with the building blocks.

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx.$$

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1 + x^2, x dx = \frac{1}{2} du \text{ transforms II to I}).$$

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1 + x^2, \quad x dx = \frac{1}{2} du \text{ transforms II to I}).$$

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1 + x^2, xdx = \frac{1}{2}du \text{ transforms II to I}).$$

- (Building block III) The third building block integral is:

$$\int \frac{1}{(1+x^2)^n} dx \quad .$$

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1+x^2, xdx = \frac{1}{2}du \text{ transforms II to I}).$$

- (Building block III) The third building block integral is:

$$\int \frac{1}{(1+x^2)^n} dx \quad .$$

- The case $n = 1$ is special for each of the building blocks:

$$\int \frac{1}{x} dx, \int \frac{x}{1+x^2} dx \text{ and } \int \frac{1}{1+x^2} dx.$$

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1+x^2, xdx = \frac{1}{2}du \text{ transforms II to I}).$$

- (Building block III) The third building block integral is:

$$\int \frac{1}{(1+x^2)^n} dx \quad .$$

- The case $n = 1$ is special for each of the building blocks:

$$\int \frac{1}{x} dx, \int \frac{x}{1+x^2} dx \text{ and } \int \frac{1}{1+x^2} dx.$$

- The case $n = 1$ we call respectively building block Ia, IIa and IIIa.

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1+x^2, xdx = \frac{1}{2}du \text{ transforms II to I}).$$

- (Building block III) The third building block integral is:

$$\int \frac{1}{(1+x^2)^n} dx \quad .$$

- The case $n = 1$ is special for each of the building blocks:

$$\int \frac{1}{x} dx, \int \frac{x}{1+x^2} dx \text{ and } \int \frac{1}{1+x^2} dx.$$

- The case $n = 1$ we call respectively building block Ia, IIa and IIIa.
The case $n > 1$ we call respectively building block Ib, IIb and IIIb.

The building blocks

Let n be a positive integer.

- (Building block I) The first building block integral is:

$$\int \frac{1}{x^n} dx \quad .$$

- (Building block II) The second building block integral is:

$$\int \frac{x}{(1+x^2)^n} dx. \quad (\text{Note: } u = 1+x^2, xdx = \frac{1}{2}du \text{ transforms II to I}).$$

- (Building block III) The third building block integral is:

$$\int \frac{1}{(1+x^2)^n} dx \quad .$$

- The case $n = 1$ is special for each of the building blocks:

$$\int \frac{1}{x} dx, \int \frac{x}{1+x^2} dx \text{ and } \int \frac{1}{1+x^2} dx.$$

- The case $n = 1$ we call respectively building block Ia, IIa and IIIa.
The case $n > 1$ we call respectively building block Ib, IIb and IIIb.
This “building block” terminology is for our convenience, and is not a part of standard mathematical terminology.

Building block Ia

Building block Ia: $\int \frac{1}{x} dx$.

Example

Integrate building block Ia

$$\int \frac{1}{x} dx$$

Building block Ia

Building block Ia: $\int \frac{1}{x} dx$.

Example

Integrate building block Ia

$$\int \frac{1}{x} dx = ?$$

Building block Ia

Building block Ia: $\int \frac{1}{x} dx$.

Example

Integrate building block Ia

$$\int \frac{1}{x} dx = \ln |x| + C$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\int \frac{1}{-4x + 5} dx$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\int \frac{1}{-4x+5} dx = \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)}$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}\int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\ &= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)}\end{aligned}$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}\int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\ &= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)} \\ &= \int \frac{1}{u} \frac{du}{(-4)}\end{aligned}$$

Set $u = -4x + 5$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}\int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\&= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)} && \left| \text{Set } u = -4x+5 \right. \\&= \int \frac{1}{u} \frac{du}{(-4)} \\&= -\frac{1}{4} \int u^{-1} du\end{aligned}$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}\int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\&= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)} && \left| \text{Set } u = -4x+5 \right. \\&= \int \frac{1}{u} \frac{du}{(-4)} \\&= -\frac{1}{4} \int u^{-1} du\end{aligned}$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}\int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\&= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)} && \left| \text{Set } u = -4x+5 \right. \\&= \int \frac{1}{u} \frac{du}{(-4)} \\&= -\frac{1}{4} \int u^{-1} du = -\frac{1}{4} \ln |u| + C\end{aligned}$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}\int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\&= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)} && \left| \text{Set } u = -4x+5 \right. \\&= \int \frac{1}{u} \frac{du}{(-4)} \\&= -\frac{1}{4} \int u^{-1} du = -\frac{1}{4} \ln |u| + C \\&= -\frac{1}{4} \ln |-4x+5| + C.\end{aligned}$$

Linear substitutions leading to building block Ia

Building block Ia: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{-4x+5} dx &= \int \frac{1}{(-4x+5)} \frac{d(-4x)}{(-4)} \\ &= \int \frac{1}{(-4x+5)} \frac{d(-4x+5)}{(-4)} && \left| \text{Set } u = -4x+5 \right. \\ &= \int \frac{1}{u} \frac{du}{(-4)} \\ &= -\frac{1}{4} \int u^{-1} du = -\frac{1}{4} \ln |u| + C \\ &= -\frac{1}{4} \ln |-4x+5| + C. \end{aligned}$$

Lin. subst. leading to building block 1a: general case

Building block 1a: $\int \frac{1}{x} dx = \ln |x| + C.$

Example

Integrate

$$\begin{aligned}
 \int \frac{1}{ax+b} dx &= \int \frac{1}{(ax+b)} \frac{d(ax)}{a} \\
 &= \int \frac{1}{(ax+b)} \frac{d(ax+b)}{a} && \left| \text{Set } u = ax+b \right. \\
 &= \int \frac{1}{u} \frac{du}{a} \\
 &= \frac{1}{a} \int u^{-1} du = \frac{1}{a} \ln |u| + C \\
 &= \frac{1}{a} \ln |ax+b| + C.
 \end{aligned}$$

Building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx, n \neq 1.$

Example (Block Ib)

$$\int \frac{1}{x^n} dx$$

Building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx, n \neq 1.$

Example (Block Ib)

$$\int \frac{1}{x^n} dx = \int x^{-n} dx$$

Building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx, n \neq 1.$

Example (Block Ib)

$$\int \frac{1}{x^n} dx = \int x^{-n} dx = ?$$

Building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx, n \neq 1.$

Example (Block Ib)

$$\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\int \frac{1}{(3x+5)^3} dx$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\int \frac{1}{(3x+5)^3} dx = \int \frac{1}{(3x+5)^3} \frac{d(\textcolor{red}{3}x)}{\textcolor{red}{3}}$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{(3x+5)^3} dx &= \int \frac{1}{(3x+5)^3} \frac{\overset{\text{red}}{d(3x)}}{3} \\ &= \int \frac{1}{(3x+5)^3} \frac{\overset{\text{red}}{d(3x+5)}}{3} \end{aligned}$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{(3x+5)^3} dx &= \int \frac{1}{(3x+5)^3} \frac{d(3x)}{3} \\ &= \int \frac{1}{(\textcolor{red}{3x+5})^3} \frac{d(\textcolor{red}{3x+5})}{3} && \left| \text{Set } \textcolor{red}{u} = \textcolor{red}{3x+5} \right. \\ &= \int \frac{1}{\textcolor{red}{u}^3} \frac{d\textcolor{red}{u}}{3} \end{aligned}$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{(3x+5)^3} dx &= \int \frac{1}{(3x+5)^3} \frac{d(3x)}{3} \\ &= \int \frac{1}{(3x+5)^3} \frac{d(3x+5)}{3} && \left| \text{Set } u = 3x+5 \right. \\ &= \int \frac{1}{u^3} \frac{du}{3} \\ &= \frac{1}{3} \int u^{-3} du \end{aligned}$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{(3x+5)^3} dx &= \int \frac{1}{(3x+5)^3} \frac{d(3x)}{3} \\ &= \int \frac{1}{(3x+5)^3} \frac{d(3x+5)}{3} && \left| \text{Set } u = 3x+5 \right. \\ &= \int \frac{1}{u^3} \frac{du}{3} \\ &= \frac{1}{3} \int u^{-3} du = \frac{1}{3} \frac{u^{-2}}{(-2)} + C \end{aligned}$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{(3x+5)^3} dx &= \int \frac{1}{(3x+5)^3} \frac{d(3x)}{3} \\ &= \int \frac{1}{(3x+5)^3} \frac{d(3x+5)}{3} && \left| \text{Set } u = 3x+5 \right. \\ &= \int \frac{1}{u^3} \frac{du}{3} \\ &= \frac{1}{3} \int u^{-3} du = \frac{1}{3} \frac{u^{-2}}{(-2)} + C \\ &= -\frac{1}{6(3x+5)^2} + C. \end{aligned}$$

Linear substitutions leading to building block Ib

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Integrate

$$\begin{aligned} \int \frac{1}{(3x+5)^3} dx &= \int \frac{1}{(3x+5)^3} \frac{d(3x)}{3} \\ &= \int \frac{1}{(3x+5)^3} \frac{d(3x+5)}{3} && \left| \text{Set } u = 3x+5 \right. \\ &= \int \frac{1}{u^3} \frac{du}{3} \\ &= \frac{1}{3} \int u^{-3} du = \frac{1}{3} \frac{u^{-2}}{(-2)} + C \\ &= -\frac{1}{6(3x+5)^2} + C. \end{aligned}$$

Lin. subst. leading to building block Ib: general case

Building block Ib: $\int \frac{1}{x^n} dx = \int x^{-n} dx = \frac{x^{-n+1}}{-n+1} + C, n \neq 1.$

Example

Let $n \neq 1$. Integrate

$$\begin{aligned}
 \int \frac{1}{(ax+b)^n} dx &= \int \frac{1}{(ax+b)^n} \frac{d(ax)}{a} \\
 &= \int \frac{1}{(ax+b)^n} \frac{d(ax+b)}{a} && \left| \text{Set } u = ax+b \right. \\
 &= \int \frac{1}{u^n} \frac{du}{a} \\
 &= \frac{1}{a} \int u^{-n} du = -\frac{1}{a} \frac{u^{-n+1}}{n-1} + C \\
 &= -\frac{1}{a(n-1)(ax+b)^{n-1}} + C.
 \end{aligned}$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\int \frac{x}{1+x^2} dx$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\int \frac{x}{1+x^2} dx = \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2}$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\ &= \int \frac{1}{1+x^2} \frac{d(1+x^2)}{2}\end{aligned}$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\ &= \int \frac{1}{\overset{1+x^2}{\textcolor{red}{1+x^2}}} \frac{\textcolor{red}{d(1+x^2)}}{2} \\ &= \int \frac{\textcolor{red}{1} \textcolor{red}{du}}{\textcolor{red}{u}} \frac{1}{2}\end{aligned}$$

Set $\textcolor{red}{u} = 1 + x^2$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\ &= \int \frac{1}{1+x^2} \frac{d(1+x^2)}{2} && \left| \text{Set } u = 1+x^2 \right. \\ &= \int \frac{1}{u} \frac{du}{2} \\ &= \frac{1}{2} \ln |u| + C\end{aligned}$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\&= \int \frac{1}{1+x^2} \frac{d(1+x^2)}{2} \\&= \int \frac{1}{u} \frac{du}{2} \\&= \frac{1}{2} \ln |u| + C = \frac{1}{2} \ln (1+x^2) + C .\end{aligned}$$

Set $u = 1 + x^2$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\&= \int \frac{1}{1+x^2} \frac{d(1+x^2)}{2} && \left| \text{Set } u = 1+x^2 \right. \\&= \int \frac{1}{u} \frac{du}{2} \\&= \frac{1}{2} \ln |u| + C = \frac{1}{2} \ln (1+x^2) + C .\end{aligned}$$

Example (Block IIIa)

$$\int \frac{1}{1+x^2} dx$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\ &= \int \frac{1}{1+x^2} \frac{d(1+x^2)}{2} \\ &= \int \frac{1}{u} \frac{du}{2} \\ &= \frac{1}{2} \ln |u| + C = \frac{1}{2} \ln (1+x^2) + C \quad .\end{aligned} \quad \left| \quad \text{Set } u = 1+x^2 \right.$$

Example (Block IIIa)

$$\int \frac{1}{1+x^2} dx = ?$$

Building blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx$. Building block IIIa: $\int \frac{1}{1+x^2} dx$.

Example (Block IIa)

$$\begin{aligned}\int \frac{x}{1+x^2} dx &= \int \frac{1}{(1+x^2)} \frac{d(x^2)}{2} \\ &= \int \frac{1}{1+x^2} \frac{d(1+x^2)}{2} \\ &= \int \frac{1}{u} \frac{du}{2} \\ &= \frac{1}{2} \ln |u| + C = \frac{1}{2} \ln (1+x^2) + C \quad .\end{aligned} \quad \left| \text{Set } u = 1 + x^2 \right.$$

Example (Block IIIa)

$$\int \frac{1}{1+x^2} dx = \arctan x + C$$

Linear substitutions leading to block IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Example

$$\int \frac{x}{2x^2+3} dx$$

Linear substitutions leading to block IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

“Theoretical way” to solve example below: transform to IIa; this is slow.

Example

$$\int \frac{x}{2x^2+3} dx$$

Linear substitutions leading to block IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

“Theoretical way” to solve example below: transform to IIa; this is slow.

Feel free to skip slide, we will redo in next slide with a shortcut.

Example

$$\int \frac{x}{2x^2+3} dx$$

Linear substitutions leading to block IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

“Theoretical way” to solve example below: transform to IIa; this is slow.

Feel free to skip slide, we will redo in next slide with a shortcut.

Example

$$\int \frac{x}{2x^2+3} dx = \int \frac{x}{3(\frac{2}{3}x^2+1)} dx = \int \frac{x}{3\left(\left(\sqrt{\frac{2}{3}}x\right)^2+1\right)} dx$$

$$= \frac{3}{2} \int \frac{\sqrt{\frac{2}{3}}x}{3\left(\left(\sqrt{\frac{2}{3}}x\right)^2+1\right)} d\left(\sqrt{\frac{2}{3}}x\right)$$

$$\left| \text{Set } u = \sqrt{\frac{2}{3}}x \right.$$

$$= \frac{1}{2} \int \frac{u}{u^2+1} du = \frac{1}{4} \ln(1+u^2) + C$$

$$= \frac{1}{4} \ln\left(\frac{1}{3}(2x^2+3)\right) + C$$

$$= \frac{1}{4} \ln(2x^2+3) + \frac{\ln(\frac{1}{3})}{4} + C$$

$$= \frac{1}{4} \ln(2x^2+3) + K.$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\int \frac{x}{2x^2 + 3} dx$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\int \frac{x}{2x^2+3} dx = \int \frac{1}{2x^2+3} d\left(\frac{x^2}{2}\right)$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\begin{aligned} \int \frac{x}{2x^2+3} dx &= \int \frac{1}{2x^2+3} d\left(\frac{x^2}{2}\right) \\ &= \int \frac{1}{2x^2+3} d\left(\frac{2x^2}{4}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\begin{aligned} \int \frac{x}{2x^2+3} dx &= \int \frac{1}{2x^2+3} d\left(\frac{x^2}{2}\right) \\ &= \int \frac{1}{2x^2+3} d\left(\frac{2x^2+3}{4}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\begin{aligned} \int \frac{x}{2x^2+3} dx &= \int \frac{1}{2x^2+3} d\left(\frac{x^2}{2}\right) \\ &= \int \frac{1}{\color{red}{2x^2+3}} d\left(\frac{\color{red}{2x^2+3}}{4}\right) \quad \Bigg| \quad \text{Set } \color{red}{u} = \color{red}{2x^2+3} \\ &= \frac{1}{4} \int \frac{1}{\color{red}{u}} d\color{red}{u} \end{aligned}$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\begin{aligned} \int \frac{x}{2x^2+3} dx &= \int \frac{1}{2x^2+3} d\left(\frac{x^2}{2}\right) \\ &= \int \frac{1}{2x^2+3} d\left(\frac{2x^2+3}{4}\right) && \left| \text{Set } u = 2x^2+3 \right. \\ &= \frac{1}{4} \int \frac{1}{u} du \\ &= \frac{1}{4} \ln |u| + C \end{aligned}$$

Linear substitutions leading to blocks IIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C$.

The example below can be done directly, without transforming to block IIa.

Example

$$\begin{aligned} \int \frac{x}{2x^2+3} dx &= \int \frac{1}{2x^2+3} d\left(\frac{x^2}{2}\right) \\ &= \int \frac{1}{2x^2+3} d\left(\frac{2x^2+3}{4}\right) && \left| \text{Set } u = 2x^2 + 3 \right. \\ &= \frac{1}{4} \int \frac{1}{u} du \\ &= \frac{1}{4} \ln |u| + C \\ &= \frac{1}{4} \ln(2x^2 + 3) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\int \frac{1}{x^2 + 2} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\int \frac{1}{x^2 + 2} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2 + 1} du = ? + C.$

Example

$$\int \frac{1}{x^2 + 2} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\int \frac{1}{x^2 + 2} dx = \int \frac{1}{2 \left(\frac{1}{2}x^2 + 1 \right)} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\int \frac{1}{x^2 + 2} dx = \int \frac{1}{2 \left(\frac{1}{2} x^2 + 1 \right)} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\int \frac{1}{x^2+2} dx = \int \frac{1}{2(\frac{1}{2}x^2+1)} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned}\int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} d\left(\frac{x}{\sqrt{2}}\right)\end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned} \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx && \left| \text{Use } 2 = (\sqrt{2})^2 \right. \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} d\left(\frac{x}{\sqrt{2}}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned}
 \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx & \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\
 &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} d\left(\frac{x}{\sqrt{2}}\right)
 \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned} \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \end{aligned} \quad \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right.$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned} \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx && \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \\ &= \frac{1}{\sqrt{2}} \int \frac{1}{u^2+1} du \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned}\int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx && \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \\ &= \frac{1}{\sqrt{2}} \int \frac{1}{u^2+1} du\end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = ? + C.$

Example

$$\begin{aligned}
 \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx && \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\
 &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \\
 &= \frac{1}{\sqrt{2}} \int \frac{1}{u^2+1} du \\
 &= \frac{1}{\sqrt{2}} ? + C
 \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = \arctan u + C$.

Example

$$\begin{aligned} \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx && \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \\ &= \frac{1}{\sqrt{2}} \int \frac{1}{u^2+1} du \\ &= \frac{1}{\sqrt{2}} \arctan(u) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = \arctan u + C$.

Example

$$\begin{aligned}\int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx \\&= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \\&= \frac{1}{\sqrt{2}} \int \frac{1}{u^2+1} du \\&= \frac{1}{\sqrt{2}} \arctan(\textcolor{red}{u}) + C \\&= \frac{1}{\sqrt{2}} \arctan\left(\frac{\textcolor{red}{x}}{\sqrt{2}}\right) + C\end{aligned}$$

$$\text{Use } 2 = (\sqrt{2})^2$$

$$\text{Set } \frac{\textcolor{red}{x}}{\sqrt{2}} = \textcolor{red}{u}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = \arctan u + C$.

Example

$$\begin{aligned} \int \frac{1}{x^2+2} dx &= \int \frac{1}{2\left(\frac{1}{2}x^2+1\right)} dx && \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\ &= \int \frac{1}{2\left(\left(\frac{x}{\sqrt{2}}\right)^2+1\right)} \sqrt{2} d\left(\frac{x}{\sqrt{2}}\right) \\ &= \frac{1}{\sqrt{2}} \int \frac{1}{u^2+1} du \\ &= \frac{1}{\sqrt{2}} \arctan(u) + C \\ &= \frac{1}{\sqrt{2}} \arctan\left(\frac{x}{\sqrt{2}}\right) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{u^2+1} du = \arctan u + C.$

Example

$$\begin{aligned}
 \int \frac{1}{x^2 + 2} dx &= \int \frac{1}{2 \left(\frac{1}{2} x^2 + 1 \right)} dx && \left| \begin{array}{l} \text{Use } 2 = (\sqrt{2})^2 \\ \text{Set } \frac{x}{\sqrt{2}} = u \end{array} \right. \\
 &= \int \frac{1}{2 \left(\left(\frac{x}{\sqrt{2}} \right)^2 + 1 \right)} \sqrt{2} d \left(\frac{x}{\sqrt{2}} \right) \\
 &= \frac{1}{\sqrt{2}} \int \frac{1}{u^2 + 1} du \\
 &= \frac{1}{\sqrt{2}} \arctan(u) + C \\
 &= \frac{1}{\sqrt{2}} \arctan \left(\frac{x}{\sqrt{2}} \right) + C
 \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{x^2+1} dx = \arctan x + C$. **Let $a > 0$.**

Example

$$\begin{aligned} \int \frac{1}{x^2 + a} dx &= \int \frac{1}{a \left(\frac{1}{a} x^2 + 1 \right)} dx \\ &= \int \frac{1}{a \left(\left(\frac{x}{\sqrt{a}} \right)^2 + 1 \right)} \sqrt{a} d \left(\frac{x}{\sqrt{a}} \right) \\ &= \frac{1}{\sqrt{a}} \int \frac{1}{u^2 + 1} du \\ &= \frac{1}{\sqrt{a}} \arctan(u) + C \\ &= \frac{1}{\sqrt{a}} \arctan \left(\frac{x}{\sqrt{a}} \right) + C \end{aligned} \quad \left| \begin{array}{l} \text{Use } a = (\sqrt{a})^2 \\ \text{Set } u = \frac{x}{\sqrt{a}} \end{array} \right.$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{x^2+1} dx = \arctan x + C$. **Let $a > 0$.**

Example

$$\begin{aligned}\int \frac{1}{x^2+a} dx &= \int \frac{1}{a\left(\frac{1}{a}x^2+1\right)} dx \\ &= \int \frac{1}{a\left(\left(\frac{x}{\sqrt{a}}\right)^2+1\right)} \sqrt{a} d\left(\frac{x}{\sqrt{a}}\right) \\ &= \frac{1}{\sqrt{a}} \int \frac{1}{u^2+1} du \\ &= \frac{1}{\sqrt{a}} \arctan(u) + C \\ &= \frac{1}{\sqrt{a}} \arctan\left(\frac{x}{\sqrt{a}}\right) + C\end{aligned}$$

Use $a = (\sqrt{a})^2$

Set $u = \frac{x}{\sqrt{a}}$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{x^2+1} dx = \arctan x + C$. **Let** $a > 0$.

Example

$$\begin{aligned} \int \frac{1}{x^2+a} dx &= \int \frac{1}{a\left(\frac{1}{a}x^2+1\right)} dx \\ &= \int \frac{1}{a\left(\left(\frac{x}{\sqrt{a}}\right)^2+1\right)} \sqrt{a} d\left(\frac{x}{\sqrt{a}}\right) \\ &= \frac{1}{\sqrt{a}} \int \frac{1}{u^2+1} du \\ &= \frac{1}{\sqrt{a}} \arctan(u) + C \\ &= \frac{1}{\sqrt{a}} \arctan\left(\frac{x}{\sqrt{a}}\right) + C \end{aligned} \quad \left| \begin{array}{l} \text{Use } a = (\sqrt{a})^2 \\ \text{Set } u = \frac{x}{\sqrt{a}} \end{array} \right.$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIIa: $\int \frac{1}{x^2+1} dx = \arctan x + C$. **Let** $a > 0$.

Example

$$\begin{aligned}\int \frac{1}{x^2+a} dx &= \int \frac{1}{a\left(\frac{1}{a}x^2+1\right)} dx \\ &= \int \frac{1}{a\left(\left(\frac{x}{\sqrt{a}}\right)^2+1\right)} \sqrt{a} d\left(\frac{x}{\sqrt{a}}\right) \\ &= \frac{1}{\sqrt{a}} \int \frac{1}{u^2+1} du \\ &= \frac{1}{\sqrt{a}} \arctan(u) + C \\ &= \frac{1}{\sqrt{a}} \arctan\left(\frac{x}{\sqrt{a}}\right) + C\end{aligned}$$

Use $a = (\sqrt{a})^2$
Set $u = \frac{x}{\sqrt{a}}$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

- Let $ax^2 + bx + c$ have no real roots.

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

- Let $ax^2 + bx + c$ have no real roots.
- We can find p, q so that the linear substitution $u = px + q$ transforms the quadratic to:

$$ax^2 + bx + c = r(u^2 + 1)$$

(where r is some number to be determined).

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

- Let $ax^2 + bx + c$ have no real roots.
- We can find p, q so that the linear substitution $u = px + q$ transforms the quadratic to:

$$ax^2 + bx + c = r(u^2 + 1)$$

(where r is some number to be determined).

- To find p, q , we **complete the square**.

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

- Let $ax^2 + bx + c$ have no real roots.
- We can find p, q so that the linear substitution $u = px + q$ transforms the quadratic to:

$$ax^2 + bx + c = r(u^2 + 1)$$

(where r is some number to be determined).

- To find p, q , we complete the square.
- In this way, integrals of the form $\int \frac{Ax + B}{ax^2 + bx + c} dx$ are transformed to combinations of building blocks IIa and IIIa.

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

- Let $ax^2 + bx + c$ have no real roots.
- We can find p, q so that the linear substitution $u = px + q$ transforms the quadratic to:

$$ax^2 + bx + c = r(u^2 + 1)$$

(where r is some number to be determined).

- To find p, q , we complete the square.
- In this way, integrals of the form $\int \frac{Ax + B}{ax^2 + bx + c} dx$ are transformed to combinations of building blocks IIa and IIIa.
- We show examples; the general case is analogous and we leave it to the student.

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

$$\int \frac{x}{x^2 + x + 1} dx =$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square.

$$\int \frac{x}{x^2 + x + 1} dx = \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square.

$$\int \frac{x}{x^2 + x + 1} dx = \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \\ &= \int \frac{u - \frac{1}{2}}{u^2 + \frac{3}{4}} du \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \\ &= \int \frac{\overset{\text{red}}{u} - \frac{1}{2}}{u^2 + \frac{3}{4}} du \\ &= \int \frac{\overset{\text{red}}{u}}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \\ &= \int \frac{u - \frac{1}{2}}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{x}{x^2 + 2 \cdot \frac{1}{2}x + \frac{1}{4} - \frac{1}{4} + 1} dx \\ &= \int \frac{x + \frac{1}{2} - \frac{1}{2}}{\left(x + \frac{1}{2}\right)^2 + \frac{3}{4}} d\left(x + \frac{1}{2}\right) \\ &= \int \frac{u - \frac{1}{2}}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\int \frac{x}{x^2 + x + 1} dx = \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\int \frac{x}{x^2 + x + 1} dx = \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du$$
$$\int \frac{1}{u^2 + \frac{3}{4}} du$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left(\frac{4}{3} u^2 + 1 \right)} du \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left(\frac{4}{3} u^2 + 1 \right)} du \\ &= \int \frac{1}{\frac{3}{4} \left(\left(\frac{2u}{\sqrt{3}} \right)^2 + 1 \right)} \frac{\sqrt{3}}{2} d\left(\frac{2u}{\sqrt{3}} \right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left(\frac{4}{3} u^2 + 1 \right)} du \\ &= \int \frac{1}{\frac{3}{4} \left(\left(\frac{2u}{\sqrt{3}} \right)^2 + 1 \right)} \frac{\sqrt{3}}{2} d\left(\frac{2u}{\sqrt{3}} \right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left(\frac{4}{3} u^2 + 1 \right)} du \\ &= \int \frac{1}{\frac{3}{4} \left(\left(\frac{2u}{\sqrt{3}} \right)^2 + 1 \right)} \frac{\sqrt{3}}{2} d\left(\frac{2u}{\sqrt{3}} \right) \\ &= \frac{2\sqrt{3}}{3} \int \frac{1}{z^2 + 1} dz \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left(\frac{4}{3} u^2 + 1 \right)} du \\ &= \int \frac{1}{\frac{3}{4} \left(\left(\frac{2u}{\sqrt{3}} \right)^2 + 1 \right)} \frac{\sqrt{3}}{2} d\left(\frac{2u}{\sqrt{3}} \right) \\ &= \frac{2\sqrt{3}}{3} \int \frac{1}{z^2 + 1} dz = \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned}
 \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\
 \int \frac{1}{u^2 + \frac{3}{4}} du &= \int \frac{1}{\frac{3}{4} \left(\frac{4}{3} u^2 + 1 \right)} du \\
 &= \int \frac{1}{\frac{3}{4} \left(\left(\frac{2u}{\sqrt{3}} \right)^2 + 1 \right)} \frac{\sqrt{3}}{2} d\left(\frac{2u}{\sqrt{3}} \right) \\
 &= \frac{2\sqrt{3}}{3} \int \frac{1}{z^2 + 1} dz = \frac{2\sqrt{3}}{3} \arctan z + C
 \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\int \frac{u}{u^2 + \frac{3}{4}} du$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\int \frac{\overset{\text{red}}{u}}{u^2 + \frac{3}{4}} \overset{\text{red}}{du} = \int \frac{1}{u^2 + \frac{3}{4}} d\left(\frac{\overset{\text{red}}{u^2}}{2}\right)$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\begin{aligned} \int \frac{u}{u^2 + \frac{3}{4}} du &= \int \frac{1}{u^2 + \frac{3}{4}} d\left(\frac{u^2}{2}\right) \\ &= \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} d\left(u^2\right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\begin{aligned} \int \frac{u}{u^2 + \frac{3}{4}} du &= \int \frac{1}{u^2 + \frac{3}{4}} d\left(\frac{u^2}{2}\right) \\ &= \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} d\left(u^2 + \frac{3}{4}\right) \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}.$

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\begin{aligned} \int \frac{u}{u^2 + \frac{3}{4}} du &= \int \frac{1}{u^2 + \frac{3}{4}} d\left(\frac{u^2}{2}\right) \\ &= \frac{1}{2} \int \frac{1}{\textcolor{red}{u}^2 + \frac{3}{4}} d\left(\textcolor{red}{u}^2 + \frac{3}{4}\right) = \frac{1}{2} \ln\left(\textcolor{red}{u}^2 + \frac{3}{4}\right) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

$$\begin{aligned} \int \frac{u}{u^2 + \frac{3}{4}} du &= \int \frac{1}{u^2 + \frac{3}{4}} d\left(\frac{u^2}{2}\right) \\ &= \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} d\left(u^2 + \frac{3}{4}\right) = \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \\ &= \frac{1}{2} \ln \left(\left(x + \frac{1}{2} \right)^2 + \frac{3}{4} \right) - \frac{\sqrt{3}}{3} \arctan \left(\frac{2u}{\sqrt{3}} \right) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \\ &= \frac{1}{2} \ln \left(\left(x + \frac{1}{2} \right)^2 + \frac{3}{4} \right) - \frac{\sqrt{3}}{3} \arctan \left(\frac{2u}{\sqrt{3}} \right) + C \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned}
 \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\
 &= \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \\
 &= \frac{1}{2} \ln \left(\left(x + \frac{1}{2} \right)^2 + \frac{3}{4} \right) - \frac{\sqrt{3}}{3} \arctan \left(\frac{2u}{\sqrt{3}} \right) + C \\
 &= \frac{1}{2} \ln \left(x^2 + x + 1 \right) - \frac{\sqrt{3}}{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) + C
 \end{aligned}$$

Linear substitutions leading to blocks IIa and IIIa

Building block IIa: $\int \frac{x}{1+x^2} dx = \frac{1}{2} \ln(1+x^2) + C.$

Building block IIIa: $\int \frac{1}{1+x^2} dx = \arctan x + C.$

Example

No real roots \Rightarrow complete the square. Let $u = x + \frac{1}{2}$, let $z = \frac{2u}{\sqrt{3}}$.

$$\begin{aligned} \int \frac{x}{x^2 + x + 1} dx &= \int \frac{u}{u^2 + \frac{3}{4}} du - \frac{1}{2} \int \frac{1}{u^2 + \frac{3}{4}} du \\ &= \frac{1}{2} \ln \left(u^2 + \frac{3}{4} \right) - \frac{1}{2} \frac{2\sqrt{3}}{3} \arctan z + C \\ &= \frac{1}{2} \ln \left(\left(x + \frac{1}{2} \right)^2 + \frac{3}{4} \right) - \frac{\sqrt{3}}{3} \arctan \left(\frac{2u}{\sqrt{3}} \right) + C \\ &= \frac{1}{2} \ln (x^2 + x + 1) - \frac{\sqrt{3}}{3} \arctan \left(\frac{2x+1}{\sqrt{3}} \right) + C \end{aligned}$$

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\int \frac{x}{(x^2 + 1)^n} dx$$

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\int \frac{x}{(x^2 + 1)^n} dx = \int \frac{1}{(x^2 + 1)^n} \frac{d(x^2 + 1)}{2}$$

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\begin{aligned}\int \frac{x}{(x^2 + 1)^n} dx &= \int \frac{1}{(x^2 + 1)^n} \frac{d(x^2 + 1)}{2} \\ &= \frac{1}{2} \int u^{-n} du\end{aligned}$$

where we used the substitution $u = x^2 + 1$.

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\begin{aligned}\int \frac{x}{(x^2 + 1)^n} dx &= \int \frac{1}{(x^2 + 1)^n} \frac{d(x^2 + 1)}{2} \\ &= \frac{1}{2} \int u^{-n} du \\ &= \begin{cases} ? & \text{if } n = 1 \\ & \text{if } n \neq 1 \end{cases},\end{aligned}$$

where we used the substitution $u = x^2 + 1$.

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\begin{aligned}\int \frac{x}{(x^2 + 1)^n} dx &= \int \frac{1}{(x^2 + 1)^n} \frac{d(x^2 + 1)}{2} \\ &= \frac{1}{2} \int u^{-n} du \\ &= \begin{cases} \frac{1}{2} \ln(x^2 + 1) + C & \text{if } n = 1 \\ & \text{if } n \neq 1 \end{cases},\end{aligned}$$

where we used the substitution $u = x^2 + 1$.

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\begin{aligned}\int \frac{x}{(x^2 + 1)^n} dx &= \int \frac{1}{(x^2 + 1)^n} \frac{d(x^2 + 1)}{2} \\ &= \frac{1}{2} \int u^{-n} du \\ &= \begin{cases} \frac{1}{2} \ln(x^2 + 1) + C & \text{if } n = 1 \\ ? & \text{if } n \neq 1 \end{cases},\end{aligned}$$

where we used the substitution $u = x^2 + 1$.

Building blocks IIa and IIb

We solve building block IIb. For completeness, we solve block IIa again as well.

Example

$$\begin{aligned}\int \frac{x}{(x^2 + 1)^n} dx &= \int \frac{1}{(x^2 + 1)^n} \frac{d(x^2 + 1)}{2} \\ &= \frac{1}{2} \int u^{-n} du \\ &= \begin{cases} \frac{1}{2} \ln(x^2 + 1) + C & \text{if } n = 1 \\ \frac{1}{2} \frac{(x^2 + 1)^{-n+1}}{(-n+1)} + C & \text{if } n \neq 1 \end{cases},\end{aligned}$$

where we used the substitution $u = x^2 + 1$.

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\arctan x + C = \int \frac{1}{x^2 + 1} dx$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2 + 1} dx \\ &= \frac{1}{x^2 + 1} x - \int x d\left(\frac{1}{x^2 + 1}\right)\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\ &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right)\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\ &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\ &= \frac{x}{x^2+1} - \int x \left(\text{?} \right) dx\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\ &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\ &= \frac{x}{x^2+1} - \int x \left(-\frac{2x}{(x^2+1)^2}\right) dx\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\ &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\ &= \frac{x}{x^2+1} - \int \overset{\text{red}}{x} \left(-\frac{\overset{\text{red}}{2x}}{(x^2+1)^2} \right) dx \\ &= \frac{x}{x^2+1} + \overset{\text{red}}{2} \int \frac{\overset{\text{red}}{x^2}}{(x^2+1)^2} dx\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\ &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\ &= \frac{x}{x^2+1} - \int x \left(-\frac{2x}{(x^2+1)^2}\right) dx \\ &= \frac{x}{x^2+1} + 2 \int \frac{-1 + x^2 + 1}{(x^2+1)^2} dx\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}
 \arctan x + C &= \int \frac{1}{x^2+1} dx \\
 &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\
 &= \frac{x}{x^2+1} - \int x \left(-\frac{2x}{(x^2+1)^2}\right) dx \\
 &= \frac{x}{x^2+1} + 2 \int \frac{-1 + x^2 + 1}{(x^2+1)^2} dx \\
 &= \frac{x}{x^2+1} + 2 \int \frac{1}{x^2+1} dx - 2 \int \frac{1}{(x^2+1)^2} dx
 \end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\&= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\&= \frac{x}{x^2+1} - \int x \left(-\frac{2x}{(x^2+1)^2}\right) dx \\&= \frac{x}{x^2+1} + 2 \int \frac{-1 + x^2 + 1}{(x^2+1)^2} dx \\&= \frac{x}{x^2+1} + 2 \int \frac{1}{x^2+1} dx - 2 \int \frac{1}{(x^2+1)^2} dx\end{aligned}$$

Building block IIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}
 \arctan x + C &= \int \frac{1}{x^2+1} dx \\
 &= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\
 &= \frac{x}{x^2+1} - \int x \left(-\frac{2x}{(x^2+1)^2}\right) dx \\
 &= \frac{x}{x^2+1} + 2 \int \frac{-1 + x^2 + 1}{(x^2+1)^2} dx \\
 &= \frac{x}{x^2+1} + 2 \int \frac{1}{x^2+1} dx - 2 \int \frac{1}{(x^2+1)^2} dx \\
 &= \frac{x}{x^2+1} + 2 \arctan x - 2 \int \frac{dx}{(x^2+1)^2}
 \end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\begin{aligned}\arctan x + C &= \int \frac{1}{x^2+1} dx \\&= \frac{1}{x^2+1} x - \int x d\left(\frac{1}{x^2+1}\right) \\&= \frac{x}{x^2+1} - \int x \left(-\frac{2x}{(x^2+1)^2}\right) dx \\&= \frac{x}{x^2+1} + 2 \int \frac{-1+x^2+1}{(x^2+1)^2} dx \\&= \frac{x}{x^2+1} + 2 \int \frac{1}{x^2+1} dx - 2 \int \frac{1}{(x^2+1)^2} dx \\&= \frac{x}{x^2+1} + 2 \arctan x - 2 \int \frac{dx}{(x^2+1)^2}\end{aligned}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\arctan x + C = \frac{x}{x^2 + 1} + 2 \arctan x - 2 \int \frac{dx}{(x^2 + 1)^2}$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\arctan x + C = \frac{x}{x^2+1} + 2 \arctan x - 2 \int \frac{dx}{(x^2+1)^2}$$

Rearrange terms

$$2 \int \frac{dx}{(1+x^2)^2} = \left(\frac{x}{x^2+1} + \arctan x \right) + L \quad .$$

Building block IIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\arctan x + C = \frac{x}{x^2+1} + 2 \arctan x - 2 \int \frac{dx}{(x^2+1)^2}$$

Rearrange terms

$$2 \int \frac{dx}{(1+x^2)^2} = \left(\frac{x}{x^2+1} + \arctan x \right) + L \quad .$$

Building block IIIb: example illustrating main idea

Example

Integrate $\int \frac{dx}{(x^2+1)^2}$. We start with an already known integral:

$$\arctan x + C = \frac{x}{x^2+1} + 2 \arctan x - 2 \int \frac{dx}{(x^2+1)^2}$$

Rearrange terms and divide by 2 to get the desired integral:

$$\int \frac{dx}{(1+x^2)^2} = \frac{1}{2} \left(\frac{x}{x^2+1} + \arctan x \right) + K .$$

Building block IIIb

- Building block IIIa:

$$\int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

Building block IIIb

- Building block IIIa:

$$\int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$\int \frac{1}{(x^2 + 1)^n} dx$$

Building block IIIb

- Building block IIIa:

$$\int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$\int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.

Building block IIIb

- Building block IIIa:

$$\int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$J(n) = \int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.
- Set $J(n) = \int \frac{1}{(x^2+1)^n} dx$.

Building block IIIb

- Building block IIIa:

$$\int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$J(n) = \int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.
- Set $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We are looking for a formula for $J(n)$.

Building block IIIb

- Building block IIIa:

$$J(1) = \int \frac{1}{(x^2 + 1)} dx = \arctan x + C .$$

- Block IIIb:

$$J(n) = \int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.
- Set $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We are looking for a formula for $J(n)$. We know $J(1) = \arctan x + C$ (this is block IIIa).

Building block IIIb

- Building block IIIa:

$$J(1) = \int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$J(n) = \int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.
- Set $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We are looking for a formula for $J(n)$. We know $J(1) = \arctan x + C$ (this is block IIIa).
- We start by $J(n-1) = \int \frac{1}{(x^2+1)^{n-1}} dx$ and integrate by parts.

Building block IIIb

- Building block IIIa:

$$J(1) = \int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$J(n) = \int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.
- Set $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We are looking for a formula for $J(n)$. We know $J(1) = \arctan x + C$ (this is block IIIa).
- We start by $J(n-1) = \int \frac{1}{(x^2+1)^{n-1}} dx$ and integrate by parts.
- In this way we end up expressing $J(n)$ via $J(n-1)$.

Building block IIIb

- Building block IIIa:

$$J(1) = \int \frac{1}{(x^2 + 1)} dx = \arctan x + C \quad .$$

- Block IIIb:

$$J(n) = \int \frac{1}{(x^2 + 1)^n} dx$$

- Unlike other cases, IIIb is much harder than IIIa.
- Set $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We are looking for a formula for $J(n)$. We know $J(1) = \arctan x + C$ (this is block IIIa).
- We start by $J(n-1) = \int \frac{1}{(x^2+1)^{n-1}} dx$ and integrate by parts.
- In this way we end up expressing $J(n)$ via $J(n-1)$.
- We work our way from $J(n)$ to $J(n-1)$, from $J(n-1)$ to $J(n-2)$, and so on, until we get to $J(1)$.

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$.

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \int \frac{1}{(x^2+1)^{n-1}} dx$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\ &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(? \right) dx \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\ &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\ &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n} \right) dx \\ &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{x^2}{(1+x^2)^n} dx \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\ &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \\ &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1+x^2-1}{(1+x^2)^n} dx \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\ &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \\ &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1+x^2-1}{(1+x^2)^n} dx \\ &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1}{(1+x^2)^{n-1}} dx \\ &\quad - 2(n-1) \int \frac{1}{(1+x^2)^n} dx \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned} J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\ &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\ &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \\ &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1+x^2-1}{(1+x^2)^n} dx \\ &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1}{(1+x^2)^{n-1}} dx \\ &\quad - 2(n-1) \int \frac{1}{(1+x^2)^n} dx \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned}
 J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\
 &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\
 &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1+x^2-1}{(1+x^2)^n} dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1}{(1+x^2)^{n-1}} dx \\
 &\quad - 2(n-1) \int \frac{1}{(1+x^2)^n} dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) .
 \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned}
 J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\
 &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\
 &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1+x^2-1}{(1+x^2)^n} dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1}{(1+x^2)^{n-1}} dx \\
 &\quad - 2(n-1) \int \frac{1}{(1+x^2)^n} dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) .
 \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$\begin{aligned}
 J(n-1) &= \int \frac{1}{(x^2+1)^{n-1}} dx \\
 &= \frac{1}{(x^2+1)^{n-1}} x - \int x d\left(\frac{1}{(1+x^2)^{n-1}}\right) \\
 &= \frac{x}{(x^2+1)^{n-1}} - \int x \left(\frac{(-n+1)2x}{(1+x^2)^n}\right) dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1+x^2-1}{(1+x^2)^n} dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1) \int \frac{1}{(1+x^2)^{n-1}} dx \\
 &\quad - 2(n-1) \int \frac{1}{(1+x^2)^n} dx \\
 &= \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) .
 \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) .$$

Rearrange to get:

$$2(n-1)J(n) = \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1)$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) .$$

Rearrange to get:

$$2(n-1)J(n) = \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1)$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) \quad . \end{aligned}$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) \quad . \end{aligned}$$

In this way we expressed $J(n)$ using $J(n-1)$.

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) \quad . \end{aligned}$$

In this way we expressed $J(n)$ using $J(n-1)$. We apply the above formula consecutively:

$$J(n) = \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1)$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) \quad . \end{aligned}$$

In this way we expressed $J(n)$ using $J(n-1)$. We apply the above formula consecutively:

$$J(n) = \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1)$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) . \end{aligned}$$

In this way we expressed $J(n)$ using $J(n-1)$. We apply the above formula consecutively:

$$J(n) = \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2} \left(\frac{x}{(2n-4)(x^2+1)^{n-2}} + \frac{2n-5}{2n-4}J(n-2) \right)$$

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) \quad . \end{aligned}$$

In this way we expressed $J(n)$ using $J(n-1)$. We apply the above formula consecutively:

$$J(n) = \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2} \left(\frac{x}{(2n-4)(x^2+1)^{n-2}} + \frac{2n-5}{2n-4} J(n-2) \right) = \dots$$

and so on.

Example

Recall that $J(n) = \int \frac{1}{(x^2+1)^n} dx$. We have that:

$$J(n-1) = \frac{x}{(x^2+1)^{n-1}} + 2(n-1)J(n-1) - 2(n-1)J(n) \quad .$$

Rearrange to get:

$$\begin{aligned} 2(n-1)J(n) &= \frac{x}{(x^2+1)^{n-1}} + (2n-3)J(n-1) \\ J(n) &= \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2}J(n-1) \quad . \end{aligned}$$

In this way we expressed $J(n)$ using $J(n-1)$. We apply the above formula consecutively:

$$J(n) = \frac{x}{(2n-2)(x^2+1)^{n-1}} + \frac{2n-3}{2n-2} \left(\frac{x}{(2n-4)(x^2+1)^{n-2}} + \frac{2n-5}{2n-4} J(n-2) \right) = \dots$$

and so on. The above can be used to write a formula for the final result, but that is as complicated as the process above.

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.
- The types in the remaining columns can be transformed to building block ones:

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.
- The types in the remaining columns can be transformed to building block ones:
 - Block I, linear substitutions: done in full detail.

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.
- The types in the remaining columns can be transformed to building block ones:
 - Block I, linear substitutions: done in full detail.
 - Block IIa, IIIa, linear substitutions: done in full detail,

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.
- The types in the remaining columns can be transformed to building block ones:
 - Block I, linear substitutions: done in full detail.
 - Block IIa, IIIa, linear substitutions: done in full detail, **by means of completing the square.**

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.
- The types in the remaining columns can be transformed to building block ones:
 - Block I, linear substitutions: done in full detail.
 - Block IIa, IIIa, linear substitutions: done in full detail, by means of completing the square.
 - Block IIb, IIIb, linear substitutions: **done by means of completing the square;**

Building block integral summary

Type	a	b	Type a, lin. sub.	Type b, lin. sub
I	$\int \frac{1}{x} dx$	$\int \frac{1}{x^n} dx$	$\int \frac{A}{ax+b} dx$	$\int \frac{A}{(ax+b)^n} dx$
II	$\int \frac{x}{x^2+1} dx$	$\int \frac{x}{(x^2+1)^n} dx$	$\int \frac{A(x+\frac{b}{2a})}{ax^2+bx+c} dx$	$\int \frac{A(x+\frac{b}{2a})}{(ax^2+bx+c)^n} dx$
III	$\int \frac{1}{x^2+1} dx$	$\int \frac{1}{(x^2+1)^n} dx$	$\int \frac{B}{ax^2+bx+c} dx$	$\int \frac{B}{(ax^2+bx+c)^n} dx$

where A, B are arbitrary constants and a, b, c are constants with $b^2 - 4ac < 0$. The quadratics in the denominators have no real roots.

- We solved building blocks I, II and III in almost complete detail.
- The types in the remaining columns can be transformed to building block ones:
 - Block I, linear substitutions: done in full detail.
 - Block IIa, IIIa, linear substitutions: done in full detail, by means of completing the square.
 - Block IIb, IIIb, linear substitutions: done by means of completing the square; **computations are analogous and we leave them for exercise.**