

# Precalculus

## Lecture 1

### Angles

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Angles

- The Unit circle
- Three Meanings of Angle
- Two Meanings of Rotation
- Angles and the Coordinate System
- Radians and Degrees
- Area cut off by an angle

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free


- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.


## Definition

The *unit circle* is the circle with radius 1 and center at the center of the coordinate system.




# Three Meanings of the Term Angle

- The term “angle” is used to denote three distinct mathematical objects:




# Three Meanings of the Term Angle

- The term “angle” is used to denote three distinct mathematical objects:
  - the (geometric) angle formed by two rays,




# Three Meanings of the Term Angle

- The term “angle” is used to denote three distinct mathematical objects:
  - the (geometric) angle formed by two rays,
  - the angle-measure of such a geometric angle



# Three Meanings of the Term Angle


- The term “angle” is used to denote three distinct mathematical objects:
  - the (geometric) angle formed by two rays,
  - the angle-measure of such a geometric angle
  - the angle-measure of a rotation.






# Three Meanings of the Term Angle

- The term “angle” is used to denote three distinct mathematical objects:
  - the (geometric) angle formed by two rays,
  - the angle-measure of such a geometric angle
  - the angle-measure of a rotation.
- **All three are referred to as “angle”**: use context to decide whether “angle” means “angle formed by two rays”, “angle measure” or “angle-measure of a rotation”.




# Three Meanings of the Term Angle

- The term “angle” is used to denote three distinct mathematical objects:
  - the (geometric) angle formed by two rays,
  - the angle-measure of such a geometric angle
  - the angle-measure of a rotation.
- All three are referred to as “angle”: use **context** to decide whether “angle” means **“angle formed by two rays”**, **“angle measure”** or **“angle-measure of a rotation”**.



# Three Meanings of the Term Angle




- The term “angle” is used to denote three distinct mathematical objects:
  - the (geometric) angle formed by two rays,
  - the angle-measure of such a geometric angle
  - the angle-measure of a rotation.
- All three are referred to as “angle”: use context to decide whether “angle” means “angle formed by two rays”, “angle measure” or “angle-measure of a rotation”.
- Except for a few introductory slides, we take full advantage of this convention.

# Geometric angle definition

## Definition (Geometric angle)


A *geometric angle* (*angle* for short) is the figure formed by two rays, called arms, sharing a common endpoint called the vertex of the angle.



# Geometric angle definition

## Definition (Geometric angle)


A *geometric angle* (*angle* for short) is the figure formed by **two rays, called arms**, sharing a common endpoint called the vertex of the angle.



# Geometric angle definition

## Definition (Geometric angle)


A *geometric angle* (*angle* for short) is the figure formed by two rays, called arms, sharing a common endpoint called the **vertex** of the angle.



# Geometric angle definition

## Definition (Geometric angle)

A *geometric angle* (*angle* for short) is the figure formed by two rays, called arms, sharing a common endpoint called the vertex of the angle. The rays are ordered.




- The ray that comes first is called the initial arm (side) of the angle.

# Geometric angle definition

## Definition (Geometric angle)

A *geometric angle* (*angle* for short) is the figure formed by two rays, called arms, sharing a common endpoint called the vertex of the angle. The rays are ordered.




- The ray that comes first is called the initial arm (side) of the angle.
- The ray that comes second is called the terminal arm (side) of the angle.



# Geometric angle definition

## Definition (Geometric angle)

A *geometric angle* (*angle* for short) is the figure formed by two rays, called arms, sharing a common endpoint called the vertex of the angle. The rays are ordered.




- The ray that comes first is called the initial arm (side) of the angle.
- The ray that comes second is called the terminal arm (side) of the angle.
- Angle measures are depicted as arcs pointing from the initial arm towards the terminal arm.

# Geometric angle definition

## Definition (Geometric angle)


A *geometric angle* (*angle* for short) is the figure formed by two rays, called arms, sharing a common endpoint called the vertex of the angle. The rays are ordered.



- The ray that comes first is called the initial arm (side) of the angle.
- The ray that comes second is called the terminal arm (side) of the angle.
- Angle measures are depicted as arcs pointing from the initial arm towards the terminal arm.
- By convention, the rays are allowed to coincide; the resulting angle is then called the *zero angle*.


# Geometric angle definition

- A ray can be identified by its starting point and any other point on the ray.




# Geometric angle definition

- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.




# Geometric angle definition

- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.




# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .

# Geometric angle definition



- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .


# Geometric angle definition



- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .




# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.
- In  $\angle AOB$  the ray  $OA$  is the initial arm and the ray  $OB$  is the terminal arm.

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.
- In  $\angle AOB$  the ray  $OA$  is the initial arm and the ray  $OB$  is the terminal arm.

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.
- In  $\angle AOB$  the ray  $OA$  is the initial arm and the ray  $OB$  is the terminal arm.
- In  $\angle BOA$  the ray  $OB$  is the initial arm, the ray  $OA$  is the terminal arm, and the angle measure points in the opposite direction.

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.
- In  $\angle AOB$  the ray  $OA$  is the initial arm and the ray  $OB$  is the terminal arm.
- In  $\angle BOA$  the ray  $OB$  is the initial arm, the ray  $OA$  is the terminal arm, and the angle measure points in the opposite direction.

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.
- In  $\angle AOB$  the ray  $OA$  is the initial arm and the ray  $OB$  is the terminal arm.
- In  $\angle BOA$  the ray  $OB$  is the initial arm, the ray  $OA$  is the terminal arm, and the angle measure **points in the opposite direction**.

# Geometric angle definition




- A ray can be identified by its starting point and any other point on the ray.
- Therefore an angle can be identified by its vertex and one point on each of its arms.
- If  $A$  is pt. on the first ray and  $B$  on the second and  $O$  is the vertex, we denote the angle by  $\angle AOB$ .
- The choice  $A$  and  $B$  is not unique - for example  $\angle AOB$  and  $\angle A'OB$  coincide.
- In  $\angle AOB$  the ray  $OA$  is the initial arm and the ray  $OB$  is the terminal arm.
- In  $\angle BOA$  the ray  $OB$  is the initial arm, the ray  $OA$  is the terminal arm, and the angle measure points in the opposite direction.
- In this way  $\angle AOB \neq \angle BOA$ .





## Definition (Radian measure of geometric angle)


The measure of a geometric angle is a number determined as follows.



## Definition (Radian measure of geometric angle)

The measure of a geometric angle is a number determined as follows.


- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.



## Definition (Radian measure of geometric angle)

The measure of a geometric angle is a number determined as follows.


- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.
- Whenever traversing the arc from initial arm to terminal results in clockwise motion, take measure with negative sign, else with positive.



## Definition (Radian measure of geometric angle)

The measure of a geometric angle is a number determined as follows.


- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.
- Whenever traversing the arc from initial arm to terminal results in clockwise motion, take measure with negative sign, else with positive.
- The unit of this angle measure is called radians, denoted by rad.



## Definition (Radian measure of geometric angle)

The measure of a geometric angle is a number determined as follows.


- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.
- Whenever traversing the arc from initial arm to terminal results in clockwise motion, take measure with negative sign, else with positive.
- The unit of this angle measure is called radians, denoted by rad.
- A circle of radius 1 has circumference  $2\pi$ .



## Definition (Radian measure of geometric angle)

The measure of a geometric angle is a number determined as follows.


- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.
- Whenever traversing the arc from initial arm to terminal results in clockwise motion, take measure with negative sign, else with positive.
- The unit of this angle measure is called radians, denoted by rad.
- A circle of radius 1 has circumference  $2\pi$ .
- Convention: half-turn angle is measured with  $\pi$  (rather than  $-\pi$ ).



## Definition (Radian measure of geometric angle)

The measure of a geometric angle is a number determined as follows.

- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.
- Whenever traversing the arc from initial arm to terminal results in clockwise motion, take measure with negative sign, else with positive.
- The unit of this angle measure is called radians, denoted by rad.
- A circle of radius 1 has circumference  $2\pi$ .
- Convention: half-turn angle is measured with  $\pi$  (rather than  $-\pi$ ).
- Therefore a geometric angle is measured with a number between  $(-\pi, \pi]$ .



## Definition (Radian measure of geometric angle)


The measure of a geometric angle is a number determined as follows.

- Its magnitude is the length of the short arc cut off by the angle from a radius 1 circle centered at the vertex.
- Whenever traversing the arc from initial arm to terminal results in clockwise motion, take measure with negative sign, else with positive.
- The unit of this angle measure is called radians, denoted by rad.
- A circle of radius 1 has circumference  $2\pi$ .
- Convention: half-turn angle is measured with  $\pi$  (rather than  $-\pi$ ).
- Therefore a geometric angle is measured with a number between  $(-\pi, \pi]$ .
- Angle measures are frequently denoted by greek letters such as  $\alpha, \beta, \gamma, \theta, \dots$




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the **limit of the lengths of ever finer straight line approximations**.




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the **limit of the lengths of ever finer straight line approximations**.




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the **limit of the lengths of ever finer straight line approximations**.




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the **limit of the lengths of ever finer straight line approximations**.




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the **limit of the lengths of ever finer straight line approximations.**




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the limit of the lengths of ever finer straight line approximations.
- The details of how this is done require integrals and we postpone this for later/another course.




# Arc-length of a circle arc

- There is a definition of arc-length of arbitrary smooth curve.
- The definition states that the arc-length of a smooth curve is the limit of the lengths of ever finer straight line approximations.
- The details of how this is done require integrals and we postpone this for later/another course.
- Until then we ask the reader to think of arc-length of a curve as the quantity obtained by “aligning a rope along the curve” and measuring the “length of this rope”.






# Arc-length of a circle arc



## Proposition

*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

# Arc-length of a circle arc




## Proposition

*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\frac{s}{r} = \frac{M}{L}$$

# Arc-length of a circle arc



## Proposition


*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\frac{s}{r} = \frac{M}{L}$$

$$\frac{1}{r} = \frac{\alpha}{L}$$

Choose  $s = 1$ , relabel  $M = \alpha$

# Arc-length of a circle arc




## Proposition

*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\begin{aligned} \frac{s}{r} &= \frac{M}{L} \\ \frac{1}{r} &= \frac{\alpha}{L} \end{aligned}$$

Choose  $s = 1$ , relabel  $M = \alpha$

# Arc-length of a circle arc




## Proposition

*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\begin{aligned} \frac{s}{r} &= \frac{M}{L} \\ \frac{1}{r} &= \frac{\alpha}{L} \\ \frac{\textcolor{red}{L}}{r} &= \alpha r \end{aligned}$$

Choose  $s = 1$ , relabel  $M = \alpha$

# Arc-length of a circle arc




## Proposition

*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\begin{aligned} \frac{s}{r} &= \frac{M}{L} \\ \frac{1}{r} &= \frac{\alpha}{L} \\ \frac{\textcolor{red}{r}}{L} &= \alpha \textcolor{red}{r} \end{aligned}$$

Choose  $s = 1$ , relabel  $M = \alpha$

# Arc-length of a circle arc



## Proposition


*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\begin{aligned}\frac{s}{r} &= \frac{M}{L} \\ \frac{1}{r} &= \frac{\alpha}{L} \\ L &= \alpha r\end{aligned}$$

Choose  $s = 1$ , relabel  $M = \alpha$

The angle-measure of a geometric angle is the arc-length cut off from a radius 1 circle, therefore we get the following.

# Arc-length of a circle arc




## Proposition

*Let two circles have common center and radii  $s$  and  $r$ . Suppose an arbitrary geometric angle with vertex at the common center of the circles cuts off short arcs of length  $M$  and  $L$ . Then  $\frac{s}{r} = \frac{M}{L}$ .*

$$\begin{aligned} \frac{s}{r} &= \frac{M}{L} \\ \frac{1}{r} &= \frac{\alpha}{L} \\ L &= \alpha r \end{aligned}$$

Choose  $s = 1$ , relabel  $M = \alpha$

The angle-measure of a geometric angle is the arc-length cut off from a radius 1 circle, therefore we get the following.




## Corollary

*The arc-length cut off by an angle with measure  $\alpha$  from a circle of radius  $r$  equals  $\alpha r$ .*




## Example



Find the length of an arc of a circle of radius 2 cut off by an angle of measure  $\frac{7\pi}{6}$  ( $= 210^\circ$ ).


## Example



Find the length of an arc of a circle of radius 2 cut off by an angle of measure  $\frac{7\pi}{6}$  ( $= 210^\circ$ ).

$$\text{arc-length} = \alpha r$$


## Example



Find the length of an arc of a circle of radius 2 cut off by an **angle of measure**  $\frac{7\pi}{6}$  ( $= 210^\circ$ ).

$$\text{arc-length} = \alpha r = \frac{7\pi}{6} \cdot 2$$


## Example



Find the length of an arc of a circle of **radius 2** cut off by an angle of measure  $\frac{7\pi}{6}$  ( $= 210^\circ$ ).

$$\text{arc-length} = \alpha r = \frac{7\pi}{6} \cdot 2$$

## Example




Find the length of an arc of a circle of radius 2 cut off by an angle of measure  $\frac{7\pi}{6}$  ( $= 210^\circ$ ).

$$\text{arc-length} = \alpha r = \frac{7\pi}{6} \cdot 2 = \frac{7\pi}{3} \approx 7.33038 \text{ (units)}$$


- The term rotation refers to two distinct objects:



- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and




- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and







- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and




- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and



- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and




- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and




- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and
  - *rotation* - an instantaneous transformation of space.




- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and
  - *rotation* - an instantaneous transformation of space. All points transition from their initial to their final positions instantaneously.



- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and
  - *rotation* - an instantaneous transformation of space. All points transition from their initial to their final positions instantaneously.
- In mathematics, the term rotation usually refers to “instantaneous” rotation.





- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and
  - *rotation* - an instantaneous transformation of space. All points transition from their initial to their final positions instantaneously.
- In mathematics, the term rotation usually refers to “instantaneous” rotation.
- In physics, the term rotation usually refers to continuous rotation (time is explicitly parametrized).






- The term rotation refers to two distinct objects:
  - *continuous rotation* (*rotation* for short) - a gradual with respect to time transformation of space and
  - *rotation* - an instantaneous transformation of space. All points transition from their initial to their final positions instantaneously.
- In mathematics, the term rotation usually refers to “instantaneous” rotation.
- In physics, the term rotation usually refers to continuous rotation (time is explicitly parametrized).
- Whether the term rotation refers to continuous rotation or to “instantaneous” rotation should be inferred from context.





## Definition (Continuous rotation)


A continuous rotation about a point (center of rotation), is a continuous motion of points for which:



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
- The distance between each rotated point and the center of rotation does not change.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
- The distance between each rotated point and the center of rotation does not change.
- The distance between each pair of rotated points is preserved.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
- The distance between each rotated point and the center of rotation does not change.
- The distance between each pair of rotated points is preserved.
- The position of a point under a continuous rotation is assumed to be a function of time.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:

- All points move in a circular fashion around the center of rotation.
- The distance between each rotated point and the center of rotation does not change.
- The distance between each pair of rotated points is preserved.
- The position of a point under a continuous rotation is assumed to be a function of time.
- The trajectory of a point is an arc of a circle.




## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:

- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.






## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.



## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:


- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.




## Definition (Continuous rotation)

A continuous rotation about a point (center of rotation), is a continuous motion of points for which:

- All points move in a circular fashion around the center of rotation.
  - The distance between each rotated point and the center of rotation does not change.
  - The distance between each pair of rotated points is preserved.
- 
- The position of a point under a continuous rotation is assumed to be a function of time.
  - The trajectory of a point is an arc of a circle.
  - A point can traverse a full circle - more than once. In this case the moving point passes through the same positions more than once.




- We say that a continuous rotation is proper if points either move **clockwise** or counter-clockwise relative to the center, without “changing direction”.




- We say that a continuous rotation is proper if points either move clockwise or **counter-clockwise** relative to the center, without “changing direction”.






- We say that a continuous rotation is **proper** if points either move clockwise or counter-clockwise relative to the center, **without “changing direction”**.



- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)


- The **radian measure of rotation is a number** whose magnitude equals the length of the arc traversed by a point divided by the distance of that point from the center of rotation.



- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)

- The radian measure of rotation is a number whose magnitude equals the **length of the arc traversed by a point** divided by the distance of that point from the center of rotation.




- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)


- The radian measure of rotation is a number whose magnitude equals the length of the arc traversed by a point divided by the **distance** of that point **from the center** of rotation.

- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.



### Definition (Radian measure of proper continuous rotation)


- The radian measure of rotation is a number whose magnitude equals the length of the arc traversed by a point divided by the distance of that point from the center of rotation.
- The sign of the radian measure is taken to be **negative** if the rotation is **clockwise**, else it is taken to be positive.



- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)


- The radian measure of rotation is a number whose magnitude equals the length of the arc traversed by a point divided by the distance of that point from the center of rotation.
- The sign of the radian measure is taken to be negative if the rotation is clockwise, else it is taken to be **positive**.



- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)

- The radian measure of rotation is a number whose magnitude equals the length of the arc traversed by a point divided by the distance of that point from the center of rotation.
- The sign of the radian measure is taken to be negative if the rotation is clockwise, else it is taken to be positive.
- The radian measure (radians for short) does not change when we change the point whose path length we are measuring.




- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)

- The radian measure of rotation is a number whose magnitude equals the length of the arc traversed by a point divided by the distance of that point from the center of rotation.
- The sign of the radian measure is taken to be negative if the rotation is clockwise, else it is taken to be positive.
- The radian measure (radians for short) does not change when we change the point whose path length we are measuring.
- The radian measure of rotation equals the signed arc-length traveled by point at **distance 1 from the center**.






- We say that a continuous rotation is proper if points either move clockwise or counter-clockwise relative to the center, without “changing direction”.

### Definition (Radian measure of proper continuous rotation)


- The radian measure of rotation is a number whose magnitude equals the length of the arc traversed by a point divided by the distance of that point from the center of rotation.
- The sign of the radian measure is taken to be negative if the rotation is clockwise, else it is taken to be positive.
- The radian measure (radians for short) does not change when we change the point whose path length we are measuring.
- The radian measure of rotation equals the signed arc-length traveled by point at distance 1 from the center.
- A circle of radius 1 has circumference  $2\pi$ , therefore a full counter-clockwise turn is measured by  $2\pi$  radians.

# Equivalence of angles




## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.




# Equivalence of angles



## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.


# Equivalence of angles



## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.


# Equivalence of angles



## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.


# Equivalence of angles



## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.


# Equivalence of angles



## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.

# Equivalence of angles




## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.



# Equivalence of angles




## Definition (Congruent angles)

Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.

## Proposition

*Two geometric angles are congruent if and only if they have equal angle measures.*

# Equivalence of angles



## Definition (Congruent angles)


Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.

## Proposition

*Two geometric angles are congruent if and only if they have equal angle measures.*

- Recall “angle” refers to both geometric angle and angle measure (depending on context).

# Equivalence of angles



## Definition (Congruent angles)


Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.

## Proposition

*Two geometric angles are congruent if and only if they have equal angle measures.*

- Recall “angle” refers to both geometric angle and angle measure (depending on context).
- The expression “the two angles are equal” is to be interpreted as “the angle measures are equal” and therefore “the geometric angles are congruent”.

# Equivalence of angles



## Definition (Congruent angles)


Two geometric angles are congruent (equivalent) if they one can be transformed onto the other with a sequence of translations and rotations.

## Proposition

*Two geometric angles are congruent if and only if they have equal angle measures.*


- Recall “angle” refers to both geometric angle and angle measure (depending on context).
- The expression “the two angles are equal” is to be interpreted as “the angle measures are equal” and therefore “the geometric angles are congruent”.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.

# Angles and the coordinate system



- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.


# Angles and the coordinate system



- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating **the point  $(1, 0)$**  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .




# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, **move it along the circle** with radius 1 for  $\alpha$  units of arc-length.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- **Select the terminal arm** by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/smaller than  $-\pi$  rad.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
  - Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
  - To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/smaller than  $-\pi$  rad.

# Angles and the coordinate system




- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
  - Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
  - To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/**smaller than  $-\pi$  rad**.

# Angles and the coordinate system



- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/smaller than  $-\pi$  rad.


# Angles and the coordinate system



- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/smaller than  $-\pi$  rad.
- In this way to every real  $\alpha$  we can assign a geometric angle.




# Angles and the coordinate system



- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/smaller than  $-\pi$  rad.
- In this way to every real  $\alpha$  we can assign a geometric angle.
- If  $\alpha$  is in the interval  $(-\pi, \pi]$  the so obtained geometric angle does have measure  $\alpha$ , else the measure of the geometric angle differs from  $\alpha$  by an even multiple of  $\pi$ .

# Angles and the coordinate system



- Given an angle measure  $\alpha$  between  $(-\pi, \pi]$ , there is a conventional way to select a geometric angle with that measure.
  - Select geometric angle's vertex to be the origin.
  - Select the initial arm of the angle on the  $x$ -axis, pointing in the positive direction.
- Select the terminal arm by rotating the point  $(1, 0)$  on the initial arm by  $|\alpha|$  radians: go clockwise if  $\alpha < 0$ , counter-clockwise if  $\alpha > 0$ .
- To rotate the point, move it along the circle with radius 1 for  $\alpha$  units of arc-length.
- The construction also works for angle measures greater than  $\pi$  rad/smaller than  $-\pi$  rad.
- In this way to every real  $\alpha$  we can assign a geometric angle.
- If  $\alpha$  is in the interval  $(-\pi, \pi]$  the so obtained geometric angle does have measure  $\alpha$ , else the **measure of the geometric angle differs from  $\alpha$  by an even multiple of  $\pi$ .**

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

- In other words, a half-turn is measured by  $\pi \text{ rad}$  or  $180^{\circ}$ .

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

- In other words, a half-turn is measured by  $\pi \text{ rad}$  or  $180^{\circ}$ .

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .
- The relationship between degrees and radians is:

$$\begin{aligned}\pi \text{ rad} &= 180^{\circ} \\ 1 \text{ rad} &= \frac{180^{\circ}}{\pi} \approx 57.3^{\circ}\end{aligned}$$

- In other words, a half-turn is measured by  $\pi \text{ rad}$  or  $180^{\circ}$ .

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^\circ$ .
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^\circ$$

$$1 \text{ rad} = \frac{180^\circ}{\pi} \approx 57.3^\circ$$

$$1^\circ = \frac{\pi}{180} \text{ rad} \approx 0.017 \text{ rad}.$$

- In other words, a half-turn is measured by  $\pi \text{ rad}$  or  $180^\circ$ .



# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .
- The relationship between degrees and radians is:

$$\pi \text{ rad} = 180^{\circ}$$

$$1 \text{ rad} = \frac{180^{\circ}}{\pi} \approx 57.3^{\circ}$$

$$1^{\circ} = \frac{\pi}{180} \text{ rad} \approx 0.017 \text{ rad}.$$

- In other words, a half-turn is measured by  $\pi \text{ rad}$  or  $180^{\circ}$ .
- Degrees are useful because the most frequently encountered fractions of a half turn are measured by a whole number of degrees.

# Degrees and radians

- Degrees is a unit for measuring angles, denoted by  $^{\circ}$ .
- The relationship between degrees and radians is:

$$\begin{aligned}\pi \text{ rad} &= 180^{\circ} \\ 1 \text{ rad} &= \frac{180^{\circ}}{\pi} \approx 57.3^{\circ} \\ 1^{\circ} &= \frac{\pi}{180} \text{ rad} \approx 0.017 \text{ rad}.\end{aligned}$$

- In other words, a half-turn is measured by  $\pi \text{ rad}$  or  $180^{\circ}$ .
- Degrees are useful because the most frequently encountered fractions of a half turn are measured by a whole number of degrees.
- If a measurement unit is not specified, it is implied to be radians. For example, in  $\sin 5$ , the number 5 stands for 5 radians.

$$t^{\circ} = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	45°	36°	-20°	360°	-720°	-225°	2015°
Rad.							

$$x = \frac{x}{\pi} 180^{\circ}.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	?						

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

### Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$						

$$x = \frac{x}{\pi} 180^\circ.$$

### Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	?					

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180} \pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	45°	36°	-20°	360°	-720°	-225°	2015°
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$					

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	?				

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								



$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$				

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	?			

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180} \pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$			

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	?		

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$		

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	?	

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	?

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								



$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.								

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	?							

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$							

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	?						

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$						

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	?					

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$					

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	?				



$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$				

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	?			

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi}180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$			

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$	?		

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$	$390^\circ$		

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$	$390^\circ$	?	

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$	$390^\circ$	$-225^\circ$	

$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$

$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.

Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$	$390^\circ$	$-225^\circ$	?



$$t^\circ = \frac{t}{180}\pi \text{ (radians).}$$

## Example

Convert from degrees to radians.

Deg.	$45^\circ$	$36^\circ$	$-20^\circ$	$360^\circ$	$-720^\circ$	$-225^\circ$	$2015^\circ$
Rad.	$\frac{\pi}{4}$	$\frac{\pi}{5}$	$-\frac{\pi}{9}$	$2\pi$	$-4\pi$	$-\frac{5\pi}{4}$	$\frac{403}{36}\pi$


$$x = \frac{x}{\pi} 180^\circ.$$

## Example

Convert from radians to degrees.


Rad.	$\frac{\pi}{3}$	$\frac{\pi}{10}$	$\frac{11\pi}{6}$	$\frac{7\pi}{4}$	$\frac{\pi}{7}$	$\frac{13\pi}{6}$	$-\frac{5\pi}{4}$	2
Deg.	$60^\circ$	$18^\circ$	$330^\circ$	$315^\circ$	$\frac{180^\circ}{7} \approx 25.7^\circ$	$390^\circ$	$-225^\circ$	$\frac{2}{\pi} 180^\circ \approx 114.6^\circ$

## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .

$$\text{arc-length} = ar$$


## Example



Find the length of an arc of a circle of **radius 3** cut off by an angle of measure  $230^\circ$ .

$$\text{arc-length} = \alpha r = ? \cdot 3$$

## Example




Find the length of an arc of a circle of radius 3 cut off by an **angle of measure  $230^\circ$** .

$$\alpha = 230^\circ$$

$$\text{arc-length} = \alpha r = ? \cdot 3$$

## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


$$\alpha = 230^\circ$$

$$= ?$$

Convert to radians

$$\text{arc-length} = \alpha r = ? \cdot 3$$

## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


$$\alpha = 230^\circ$$

$$= 230^\circ \frac{\pi \text{ rad}}{180^\circ}$$

Convert to radians

$$\text{arc-length} = \alpha r = ? \cdot 3$$

## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


$$\begin{aligned}\alpha &= 230^\circ \\ &= 230^\circ \frac{\pi \text{ rad}}{180^\circ} = \frac{23}{18} \pi \text{ rad}\end{aligned}$$

Convert to radians

$$\text{arc-length} = \alpha r = ? \cdot 3$$



## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


$$\alpha = 230^\circ$$

$$= 230^\circ \frac{\pi \text{ rad}}{180^\circ} = \frac{23}{18} \pi \text{ rad}$$

Convert to radians

$$\text{arc-length} = \alpha r = \frac{23\pi}{18} \cdot 3$$

## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


$$\alpha = 230^\circ$$

$$= 230^\circ \frac{\pi \text{ rad}}{180^\circ} = \frac{23}{18} \pi \text{ rad}$$

$$\text{arc-length} = \alpha r = \frac{23\pi}{18} \cdot 3 = \frac{23\pi}{6}$$

Convert to radians


## Example



Find the length of an arc of a circle of radius 3 cut off by an angle of measure  $230^\circ$ .


$$\begin{aligned}\alpha &= 230^\circ \\ &= 230^\circ \frac{\pi \text{ rad}}{180^\circ} = \frac{23}{18} \pi \text{ rad} \\ \text{arc-length} &= \alpha r = \frac{23\pi}{18} \cdot 3 = \frac{23\pi}{6} \approx 12.043\end{aligned}$$

Convert to radians




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	?										




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0										




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	?									



The most frequently encountered angles are given in the table below.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$									



The most frequently encountered angles are given in the table below.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	?								






The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$								




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	?							




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$							




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	?						




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$						




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	?					



The most frequently encountered angles are given in the table below.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$					



The most frequently encountered angles are given in the table below.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	?				






The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$				




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	?			




The most frequently encountered angles are given in the table below.

Deg.	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$	$120^\circ$	$135^\circ$	$150^\circ$	$180^\circ$	$270^\circ$	$360^\circ$
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$			




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	?		




The most frequently encountered angles are given in the table below.

Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$		




The most frequently encountered angles are given in the table below.

Deg.	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$	$120^\circ$	$135^\circ$	$150^\circ$	$180^\circ$	<b><math>270^\circ</math></b>	$360^\circ$
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$	<b>?</b>	



The most frequently encountered angles are given in the table below.


Deg.	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$	$120^\circ$	$135^\circ$	$150^\circ$	$180^\circ$	<b><math>270^\circ</math></b>	$360^\circ$
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$	<b><math>\frac{3\pi}{2}</math></b>	



The most frequently encountered angles are given in the table below.


Deg.	$0^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$90^\circ$	$120^\circ$	$135^\circ$	$150^\circ$	$180^\circ$	$270^\circ$	$360^\circ$
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$	$\frac{3\pi}{2}$	?






The most frequently encountered angles are given in the table below.


Deg.	0°	30°	45°	60°	90°	120°	135°	150°	180°	270°	360°
Rad.	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	$\pi$	$\frac{3\pi}{2}$	$2\pi$




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.



- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.




- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.






- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.



- Integer quantities of radians are not rational multiples of (the measure of) a half-turn and are not easy to compute with.
- For example to determine in which quadrant is an angle of  $k$  radians located one needs to know the numerical value of  $\frac{k}{\pi}$ , which requires knowledge of  $\pi$  with great numerical accuracy.

## Definition (Coterminal Angles)

Two angles (angle measures) are called coterminal if the corresponding geometric angles have the same initial and terminal sides.



## Observation

*The set of angles coterminal with  $\alpha$  consists of the angles  $\alpha + 2k\pi$ , where  $k$  runs over the set of integers. In other words, the angles coterminal with  $\alpha$  are the angles:*

$$\dots, \alpha - 6\pi, \alpha - 4\pi, \alpha - 2\pi, \alpha, \alpha + 2\pi, \alpha + 4\pi, \alpha + 6\pi, \dots$$

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

To find which among the angles  $\frac{\pi}{4} + 2k\pi$  lie in the interval  $[-2\pi, \pi]$ , we write them as an infinite list

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

To find which among the angles  $\frac{\pi}{4} + 2k\pi$  lie in the interval  $[-2\pi, \pi]$ , we write them as an infinite list

$$\dots, \frac{\pi}{4} - 4\pi, \frac{\pi}{4} - 2\pi, \frac{\pi}{4}, \frac{\pi}{4} + 2\pi, \frac{\pi}{4} + 4\pi, \dots$$

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

To find which among the angles  $\frac{\pi}{4} + 2k\pi$  lie in the interval  $[-2\pi, \pi]$ , we write them as an infinite list (**we indicate the unboundedness of the list by ellipsis dots**)

$$\dots, \frac{\pi}{4} - 4\pi, \frac{\pi}{4} - 2\pi, \frac{\pi}{4}, \frac{\pi}{4} + 2\pi, \frac{\pi}{4} + 4\pi, \dots$$



## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

To find which among the angles  $\frac{\pi}{4} + 2k\pi$  lie in the interval  $[-2\pi, \pi]$ , we write them as an infinite list (we indicate the unboundedness of the list by ellipsis dots) **and cross out the angles that lie outside of the desired interval.**

$$\dots, \frac{\pi}{4} - 4\pi, \frac{\pi}{4} - 2\pi, \frac{\pi}{4}, \frac{\pi}{4} + 2\pi, \frac{\pi}{4} + 4\pi, \dots$$

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

To find which among the angles  $\frac{\pi}{4} + 2k\pi$  lie in the interval  $[-2\pi, \pi]$ , we write them as an infinite list (we indicate the unboundedness of the list by ellipsis dots) **and cross out the angles that lie outside of the desired interval.**

$$\dots, \cancel{\frac{\pi}{4} - 4\pi}, \frac{\pi}{4} - 2\pi, \frac{\pi}{4}, \cancel{\frac{\pi}{4} + 2\pi}, \cancel{\frac{\pi}{4} + 4\pi}, \dots$$

## Example

- Find all angles that are coterminal to  $\frac{\pi}{4}$ .
- Find all angles in the interval  $[-2\pi, \pi]$  that are coterminal to  $\frac{\pi}{4}$ .

By theory, the angles coterminal with  $\frac{\pi}{4}$  are all angles of the form

$$\frac{\pi}{4} + 2k\pi.$$

To find which among the angles  $\frac{\pi}{4} + 2k\pi$  lie in the interval  $[-2\pi, \pi]$ , we write them as an infinite list (we indicate the unboundedness of the list by ellipsis dots) and cross out the angles that lie outside of the desired interval.


$$\dots, \cancel{\frac{\pi}{4} - 4\pi}, \frac{\pi}{4} - 2\pi, \frac{\pi}{4}, \cancel{\frac{\pi}{4} + 2\pi}, \cancel{\frac{\pi}{4} + 4\pi}, \dots$$

Our final answer is  $-\frac{7\pi}{4}, \frac{\pi}{4}$

# Complementary angles

## Definition


Two positive angles are called complementary when they sum to a right angle, i.e., an angle of measure  $\frac{\pi}{2} = 90^\circ$ .




# Supplementary angles

## Definition

Two positive angles are called supplementary when they sum to  $\pi = 180^\circ$ .



A sector of a circle is the region cut off from a circle by an angle whose vertex is at the center of the circle.



### Proposition (Area of a circle sector)

*The area of a circle sector equals*

$$\frac{1}{2}\alpha r^2,$$

*where  $\alpha$  is the angle of the sector.*