

# Precalculus

## Lecture 20

Todor Milev

`https://github.com/tmilev/freecalc`

2020

# Outline

## 1 A Catalog of Essential Functions

- Linear Functions
- Polynomials
- Power Functions
- Rational Functions
- Algebraic Functions
- Transcendental Functions
- Miscellaneous

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:

<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Linear Functions

## Definition (Linear Function)

A linear function is a function the graph of which is a line. We can write any linear function in slope-intercept form:

$$f(x) = mx + b.$$

$m$  is called the slope, and  $b$  is called the  $y$ -intercept.


# Linear Functions

## Definition (Linear Function)

A linear function is a function the graph of which is a line. We can write any linear function in slope-intercept form:

$$f(x) = mx + b.$$

$m$  is called the slope, and  $b$  is called the  $y$ -intercept.



- Any non-vertical line arises as the graph of a linear function.


# Linear Functions

## Definition (Linear Function)


A linear function is a function the graph of which is a line. We can write any linear function in slope-intercept form:

$$f(x) = mx + b.$$


$m$  is called the slope, and  $b$  is called the  $y$ -intercept.




- Any non-vertical line arises as the graph of a linear function.



- Vertical lines fail the vertical line test and therefore are not graphs of a function of  $x$ .




$f(x)$	Direction	y-intercept
$x + 1$ $-0.5x$ $-1$		







$f(x)$	Direction	y-intercept
$x + 1$ $-0.5x$ $-1$	↗	


- $m > 0$  means the graph of  $f$  points up (↗).






$f(x)$	Direction	y-intercept
$x + 1$		
$-0.5x$		
$-1$		

- $m > 0$  means the graph of  $f$  points up ().
- $m < 0$  means the graph of  $f$  points down (.




$f(x)$	Direction	y-intercept
$x + 1$	$\nearrow$	
$-0.5x$	$\searrow$	
$-1$	$\rightarrow$	

- $m > 0$  means the graph of  $f$  points up ( $\nearrow$ ).
- $m < 0$  means the graph of  $f$  points down ( $\searrow$ ).
- $m = 0$  means the graph of  $f$  is horizontal ( $\rightarrow$ ).




$f(x)$	Direction	y-intercept
$x + 1$	$\nearrow$	$1$
$-0.5x$	$\searrow$	
$-1$	$\rightarrow$	

- $m > 0$  means the graph of  $f$  points up ( $\nearrow$ ).
- $m < 0$  means the graph of  $f$  points down ( $\searrow$ ).
- $m = 0$  means the graph of  $f$  is horizontal ( $\rightarrow$ ).
- $b$  tells us the height of the point where the graph hits the y-axis.



$f(x)$	Direction	y-intercept
$x + 1$	$\nearrow$	1
$-0.5x + 0$	$\searrow$	0
$-1$	$\rightarrow$	

- $m > 0$  means the graph of  $f$  points up ( $\nearrow$ ).
- $m < 0$  means the graph of  $f$  points down ( $\searrow$ ).
- $m = 0$  means the graph of  $f$  is horizontal ( $\rightarrow$ ).
- $b$  tells us the height of the point where the graph hits the y-axis.



$f(x)$	Direction	y-intercept
$x + 1$	$\nearrow$	1
$-0.5x$	$\searrow$	0
$-1$	$\rightarrow$	$-1$

- $m > 0$  means the graph of  $f$  points up ( $\nearrow$ ).
- $m < 0$  means the graph of  $f$  points down ( $\searrow$ ).
- $m = 0$  means the graph of  $f$  is horizontal ( $\rightarrow$ ).
- $b$  tells us the height of the point where the graph hits the y-axis.

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

If we interpret  $x$  as an indeterminate formal expression, rather than a number, we say that  $f(x)$  is a polynomial (rather than a polynomial function).

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$					
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	?				
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					



# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes				
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	?			
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4			
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	?		
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1		
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	?	
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	
$6$					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	?
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					



# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6					
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	?				
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	Yes				
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6x^2 - \frac{1}{2}x + \sqrt{x}$	Yes	?			
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	Yes	0			
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	Yes	0	?		
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	Yes	0	6		
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6x^2 - \frac{1}{2}x + \sqrt{x}$	Yes	0	6	?	
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					



# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6x^2 - \frac{1}{2}x + \sqrt{x}$	Yes	0	6	0	
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6x^2 - \frac{1}{2}x + \sqrt{x}$	Yes	0	6	0	?
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$					
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	?				
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative **integer** and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$					
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
$6$	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	?				
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes				
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	?			
$3x^2 - \frac{1}{2x} + \sqrt{2}$					



# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2			
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	?		
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$		
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	?	
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	?
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	3
$3x^2 - \frac{1}{2x} + \sqrt{2}$					

# Polynomials

## Definition (Polynomial Function)

A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a non-negative integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .

$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	3
$3x^2 - \frac{1}{2x} + \sqrt{2}$	?				



# Polynomials

## Definition (Polynomial Function)


A polynomial function is a function  $f$  of the form

$$f(x) = a_0 + a_1x + a_2x^2 + \cdots + a_{n-1}x^{n-1} + a_nx^n,$$

where  $n$  is a **non-negative** integer and  $a_0, \dots, a_n$  are real numbers, called the coefficients. If  $a_n \neq 0$  the integer  $n$  is called the degree of  $f$ .


$f(x)$	Polynomial?	Degree	$a_0$	$a_1$	$a_2$
$x^4 - x + 1$	Yes	4	1	-1	0
6	Yes	0	6	0	0
$3x^2 - \frac{1}{2}x + \sqrt{x}$	No				
$3x^2 - \frac{1}{2}x + \sqrt{2}$	Yes	2	$\sqrt{2}$	$-\frac{1}{2}$	3
$3x^2 - \frac{1}{2x} + \sqrt{2}$	No				

- Linear functions are polynomial (functions).




Linear

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.




Quadratic

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.
- And there are many more.




## Cubic

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.
- And there are many more.



## Quartic

- Linear functions are polynomial (functions).
- So are quadratic functions. Their graphs are parabolas.
- And there are many more.



Quintic

# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = x^a .$$

# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = x^a .$$

$x$  = base.



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = x^a .$$

$x$  = base.  $a$  = **exponent** or **power**.

# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a .$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer  $(1, 2, 3, \dots)$

then  $x^a$  = polynomial function.

$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$

# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer  $(1, 2, 3, \dots)$

then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )

then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )

then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a \quad .$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$





# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b =$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b = ?$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b =$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = ?$$

$$x^{a+b} =$$

$$x^{-a} =$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as


$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.

$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$\begin{aligned} (x^a)^b &= x^{ab} \\ (xy)^b &= x^b y^b \\ x^{a+b} &= \\ x^{-a} &= \end{aligned}$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as


$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.

$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$\begin{aligned} (x^a)^b &= x^{ab} \\ (xy)^b &= x^b y^b \\ x^{a+b} &= ? \\ x^{-a} &= \end{aligned}$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as


$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.

$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$\begin{aligned} (x^a)^b &= x^{ab} \\ (xy)^b &= x^b y^b \\ x^{a+b} &= x^a x^b \\ x^{-a} &= \end{aligned}$$



# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as

$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.


$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$(x^a)^b = x^{ab}$$

$$(xy)^b = x^b y^b$$

$$x^{a+b} = x^a x^b$$

$$x^{-a} = ?$$





# Power Functions

## Definition (Power Function)

Let  $x > 0$ ,  $a$  - arbitrary real number. The power function is defined as


$$f(x) = e^{a \ln x} = x^a.$$

$x$  = base.  $a$  = exponent or power. First equality = one of ways to define for non-integer  $a$  (we study  $\ln x$ ,  $e^x$  later).

If  $a$  - positive integer ( $1, 2, 3, \dots$ )  
then  $x^a$  = polynomial function.

$x^n = \underbrace{x \dots x}_{n \text{ times}}$  when  $n$ -integer.

$$\begin{aligned} (x^a)^b &= x^{ab} \\ (xy)^b &= x^b y^b \\ x^{a+b} &= x^a x^b \\ x^{-a} &= \frac{1}{x^a} \end{aligned}$$




- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .

- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.


- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.
- Let  $x > 0$ . For  $n = 2m + 1$ -odd, we can extend the definition of  $n^{\text{th}}$  root to negative numbers by  $\sqrt[n]{-x} := -\sqrt[n]{x}$ .

- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.
- Let  $x > 0$ . For  $n = 2m + 1$ -odd, we can extend the definition of  $n^{\text{th}}$  root to negative numbers by  ${}^{2m+1}\sqrt{-x} := - {}^{2m+1}\sqrt{x}$ .
- In this course, even roots of negative numbers are not defined.



- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.
- Let  $x > 0$ . For  $n = 2m + 1$ -odd, we can extend the definition of  $n^{\text{th}}$  root to negative numbers by  $\sqrt[n]{-x} := -\sqrt[n]{x}$ .
- In this course, even roots of negative numbers are not defined.
- The graph of  $\sqrt{x}$  is the top half of the parabola  $x = y^2$ .



- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.
- Let  $x > 0$ . For  $n = 2m + 1$ -odd, we can extend the definition of  $n^{\text{th}}$  root to negative numbers by  ${}^{2m+1}\sqrt{-x} := -{}^{2m+1}\sqrt{x}$ .
- In this course, even roots of negative numbers are not defined.
- The graph of  $\sqrt{x}$  is the top half of the parabola  $x = y^2$ . Similarly for  $y = {}^{2m}\sqrt{x}$ , we graph top of  $x = y^{2m}$ .





- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x} =$  the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.
- Let  $x > 0$ . For  $n = 2m + 1$ -odd, we can extend the definition of  $n^{\text{th}}$  root to negative numbers by  $\sqrt[n]{-x} := -\sqrt[n]{x}$ .
- In this course, even roots of negative numbers are not defined.
- The graph of  $\sqrt{x}$  is the top half of the parabola  $x = y^2$ . Similarly for  $y = \sqrt[2m]{x}$ , we graph top of  $x = y^{2m}$ .
- The graph of the cube root  $f(x) = \sqrt[3]{x}$  is the graph of the polynomial  $x = y^3$ .






- $n$  - positive integer,  $f(x) = x^{\frac{1}{n}} = \sqrt[n]{x}$  = the  $n^{\text{th}}$  root function.  
 $\sqrt[n]{x} \geq 0$  for  $x \geq 0$ .
- For  $n = 2$ , we get the square root  $\sqrt{x}$ ; for  $n = 3$  we get the cube root  $\sqrt[3]{x}$ , and so on.
- Let  $x > 0$ . For  $n = 2m + 1$ -odd, we can extend the definition of  $n^{\text{th}}$  root to negative numbers by  $\sqrt[n]{-x} := -\sqrt[n]{x}$ .
- In this course, even roots of negative numbers are not defined.
- The graph of  $\sqrt{x}$  is the top half of the parabola  $x = y^2$ . Similarly for  $y = \sqrt[2m]{x}$ , we graph top of  $x = y^{2m}$ .
- The graph of the cube root  $f(x) = \sqrt[3]{x}$  is the graph of the polynomial  $x = y^3$ . Similarly for  $y = \sqrt[2m+1]{x}$ , we graph  $x = y^{2m+1}$ .



$f(x) = x^{-1} = \frac{1}{x}$  is called the reciprocal function. Its graph has equation  $y = \frac{1}{x}$ , or  $xy = 1$ , and is an hyperbola with the coordinate axes as its



asymptotes.

# Rational Functions

## Definition (Rational Function)

A rational function is a quotient of two polynomials; that is, a function of the form

$$f(x) = \frac{g(x)}{h(x)},$$

where  $g$  and  $h$  are polynomials.


# Rational Functions

## Definition (Rational Function)

A rational function is a quotient of two polynomials; that is, a function of the form

$$f(x) = \frac{g(x)}{h(x)},$$

where  $g$  and  $h$  are polynomials.



## Example ( $x/(x^2 - 1)$ )

The function

$$f(x) = \frac{x}{x^2 - 1}$$

is a rational function.

# Algebraic Functions

## (Algebraic Function)

*A function in  $x$  that can be constructed using  $x$ , constants, and finitely many of the operations  $+$ ,  $-$ ,  $*$ ,  $/$ , and  $\sqrt[n]{\phantom{x}}$  is an algebraic function.*

# Algebraic Functions

## (Algebraic Function)

*A function in  $x$  that can be constructed using  $x$ , constants, and finitely many of the operations  $+$ ,  $-$ ,  $*$ ,  $/$ , and  $\sqrt[n]{\phantom{x}}$  is an algebraic function.*

*Outside of present course: function  $f(x)$  = algebraic if it satisfies a polynomial equation with polynomial coefficients, i.e.,  $a_0(x) + a_1(x)f(x) + \cdots + a_n(x)(f(x))^n = 0$  for some polynomials  $a_i(x)$ .*


# Algebraic Functions

## (Algebraic Function)


A function in  $x$  that can be constructed using  $x$ , constants, and finitely many of the operations  $+$ ,  $-$ ,  $*$ ,  $/$ , and  $\sqrt[n]{\phantom{x}}$  is an algebraic function.

Outside of present course: function  $f(x)$  = algebraic if it satisfies a polynomial equation with polynomial coefficients, i.e.,  $a_0(x) + a_1(x)f(x) + \cdots + a_n(x)(f(x))^n = 0$  for some polynomials  $a_i(x)$ .


Examples.



$$y = (x-1)^{3/2}$$



$$y = \frac{1}{5}(4x - x^2)^{3/2}$$



$$y = (x-1)^{3/2}$$

# Transcendental Functions

Transcendental functions include many classes of functions.



# Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as  $\cos x$ ,  $\sin x$ ,  $\tan x$ , etc.

# Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as  $\cos x$ ,  $\sin x$ ,  $\tan x$ , etc.
- Exponential functions such as  $2^x$ ,  $\left(\frac{1}{2}\right)^x$ ,  $5^x$ ,  $e^x$ , etc.

# Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as  $\cos x$ ,  $\sin x$ ,  $\tan x$ , etc.
- Exponential functions such as  $2^x$ ,  $\left(\frac{1}{2}\right)^x$ ,  $5^x$ ,  $e^x$ , etc.
- The logarithm function  $\ln x$ .

# Transcendental Functions

Transcendental functions include many classes of functions.

- Trigonometric functions such as  $\cos x$ ,  $\sin x$ ,  $\tan x$ , etc.
- Exponential functions such as  $2^x$ ,  $\left(\frac{1}{2}\right)^x$ ,  $5^x$ ,  $e^x$ , etc.
- The logarithm function  $\ln x$ .
- And many more.

# Transcendental Functions


Transcendental functions include many classes of functions.

- Trigonometric functions such as  $\cos x$ ,  $\sin x$ ,  $\tan x$ , etc.
- Exponential functions such as  $2^x$ ,  $\left(\frac{1}{2}\right)^x$ ,  $5^x$ ,  $e^x$ , etc.
- The logarithm function  $\ln x$ .
- And many more.
- Outside of the present course: by definition, a function is transcendental if it is not algebraic, i.e., if it satisfies no polynomial equation with polynomial coefficients.

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .


In computer science this function is called the *floor* function.



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = ?$$

$$\lfloor 4.8 \rfloor =$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor =$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$

$$\left\lfloor -\frac{1}{2} \right\rfloor =$$


$$\lfloor -\pi \rfloor =$$



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = ?$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor =$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = ?$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor =$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = ?$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor =$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor = ?$$

$$\lfloor -\pi \rfloor =$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$

$$\left\lfloor -\frac{1}{2} \right\rfloor = -1$$


$$\lfloor -\pi \rfloor =$$



## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$


$$\left\lfloor -\frac{1}{2} \right\rfloor = -1$$

$$\lfloor -\pi \rfloor = ?$$

## Definition (Greatest Integer Function)

The *greatest integer function*  $\lfloor x \rfloor$  is defined as the largest integer that is less than or equal to  $x$ .

In computer science this function is called the *floor* function.



$$\lfloor 4 \rfloor = 4$$

$$\lfloor 4.8 \rfloor = 4$$

$$\lfloor \pi \rfloor = 3$$

$$\lfloor \sqrt{2} \rfloor = 1$$

$$\left\lfloor -\frac{1}{2} \right\rfloor = -1$$

$$\lfloor -\pi \rfloor = -4$$