

# Calculus II

## Lecture 17

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Basic divergence tests

# Outline

- 1 Basic divergence tests
- 2 The Integral Test and Estimates of Sums
  - The Integral Test
  - Estimating Sums

# Outline

- 1 Basic divergence tests
- 2 The Integral Test and Estimates of Sums
  - The Integral Test
  - Estimating Sums
- 3 The Comparison Test

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .



## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.
- Let  $\lim_{n \rightarrow \infty} s_n = s$ .

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.
- Let  $\lim_{n \rightarrow \infty} s_n = s$ .
- Then  $\lim_{n \rightarrow \infty} s_{n-1} =$

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.
- Let  $\lim_{n \rightarrow \infty} s_n = s$ .
- Then  $\lim_{n \rightarrow \infty} s_{n-1} = s$ .

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.
- Let  $\lim_{n \rightarrow \infty} s_n = s$ .
- Then  $\lim_{n \rightarrow \infty} s_{n-1} = s$ .
- Therefore

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} (s_n - s_{n-1})$$

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.
- Let  $\lim_{n \rightarrow \infty} s_n = s$ .
- Then  $\lim_{n \rightarrow \infty} s_{n-1} = s$ .
- Therefore

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} (s_n - s_{n-1}) = s - s$$

## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

## Proof.

- Let  $s_n = a_1 + a_2 + \cdots + a_n$ .
- Then  $a_n = s_n - s_{n-1}$ .
- Since  $\sum_{n=1}^{\infty} a_n$  is convergent, the sequence  $\{s_n\}$  is convergent.
- Let  $\lim_{n \rightarrow \infty} s_n = s$ .
- Then  $\lim_{n \rightarrow \infty} s_{n-1} = s$ .
- Therefore

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} (s_n - s_{n-1}) = s - s = 0$$



## Theorem

*If the series  $\sum_{n=1}^{\infty} a_n$  is convergent, then  $\lim_{n \rightarrow \infty} a_n = 0$ .*

This is just a restatement of the previous theorem:

## Theorem (The Divergence Test)

*If  $\lim_{n \rightarrow \infty} a_n$  doesn't exist or if  $\lim_{n \rightarrow \infty} a_n \neq 0$ , then the series  $\sum_{n=1}^{\infty} a_n$  is divergent.*



## Example

Show that the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2+4}$  diverges.

## Example

Show that the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2+4}$  diverges.

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^2}{5n^2 + 4}$$

## Example

Show that the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2+4}$  diverges.

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^2}{5n^2 + 4} \cdot \frac{\frac{1}{n^2}}{\frac{1}{n^2}}$$

## Example

Show that the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2+4}$  diverges.

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^2}{5n^2+4} \cdot \frac{\frac{1}{n^2}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{1}{5 + \frac{4}{n^2}}$$

## Example

Show that the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2+4}$  diverges.

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^2}{5n^2+4} \cdot \frac{\frac{1}{n^2}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{1}{5 + \frac{4}{n^2}} = \frac{1}{5} \neq 0$$

## Example

Show that the series  $\sum_{n=1}^{\infty} \frac{n^2}{5n^2+4}$  diverges.

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} \frac{n^2}{5n^2+4} \cdot \frac{\frac{1}{n^2}}{\frac{1}{n^2}} = \lim_{n \rightarrow \infty} \frac{1}{5 + \frac{4}{n^2}} = \frac{1}{5} \neq 0$$

Therefore, by the Divergence Test, the series diverges.

# The Integral Test and Estimates of Sums

- In general, it is not easy to find the sum of a series.
- We could do this for  $\sum_{n=1}^{\infty} \frac{1}{n(n+1)}$  because we found a simple formula for the  $n$ th partial sum  $s_n$ .
- In the next few sections, we'll learn techniques for showing whether a series is convergent or divergent without explicitly computing its sum.

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$



$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$


- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447




$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.




$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .




$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .




$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \dots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .

$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .






$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .


$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .



$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .




$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .
- The improper integral  $\int_1^{\infty} \frac{1}{x^2} dx$  is

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .




$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .
- The improper integral  $\int_1^{\infty} \frac{1}{x^2} dx$  is ?

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .



$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447


- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .
- The improper integral  $\int_1^{\infty} \frac{1}{x^2} dx$  is convergent.

$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .


$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .
- The improper integral  $\int_1^{\infty} \frac{1}{x^2} dx$  is convergent.



$$\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{1}{1^2} + \frac{1}{2^2} + \frac{1}{3^2} + \frac{1}{4^2} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be converging.
- How do we prove it?
- Use  $f(x) = \frac{1}{x^2}$ .



$n$	$s_n = \sum_{i=1}^n \frac{1}{i^2}$
5	1.4636
10	1.5498
50	1.6251
100	1.6350
500	1.6429
1000	1.6439
5000	1.6447

- $\frac{1}{1^2}$  is the area of a rectangle.
- So is  $\frac{1}{2^2} = \frac{1}{4}$ .
- The improper integral  $\int_1^{\infty} \frac{1}{x^2} dx$  is convergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{n^2}$  is convergent.

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.

$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681



$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.

$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$


- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?

$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .

$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681




$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .


$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.



$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .




$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .



$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681


- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \dots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .


$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .



$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .



$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .




$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .


$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .
- $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$  is ?



$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \dots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .




$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .
- $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$  is ?

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .



$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681


- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .
- $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$  is ?

$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \dots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .


$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .
- $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$  is divergent.



$$\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}} = \frac{1}{\sqrt{1}} + \frac{1}{\sqrt{2}} + \frac{1}{\sqrt{3}} + \frac{1}{\sqrt{4}} + \cdots$$

- Use a computer to calculate partial sums.
- Appears to be diverging.
- How do we prove it?
- Use  $f(x) = \frac{1}{\sqrt{x}}$ .



$n$	$s_n = \sum_{i=1}^n \frac{1}{\sqrt{i}}$
5	3.2317
10	5.0210
50	12.7524
100	18.5896
500	43.2834
1000	61.8010
5000	139.9681

- $\frac{1}{\sqrt{1}}$  is the area of a rectangle.
- So is  $\frac{1}{\sqrt{2}}$ .
- $\int_1^{\infty} \frac{1}{\sqrt{x}} dx$  is divergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{\sqrt{n}}$  is divergent.

## Theorem (The Integral Test)

Let  $f$  be a continuous, positive, decreasing function on  $[1, \infty)$  and let  $a_n = f(n)$ . Then the series  $\sum_{n=1}^{\infty} a_n$  is convergent if and only if the improper integral  $\int_1^{\infty} f(x)dx$  is convergent. In other words,

- 1 If  $\int_1^{\infty} f(x)dx$  is convergent, then  $\sum_{n=1}^{\infty} a_n$  is convergent.
- 2 If  $\int_1^{\infty} f(x)dx$  is divergent, then  $\sum_{n=1}^{\infty} a_n$  is divergent.

## Theorem (The Integral Test)

Let  $f$  be a continuous, positive, decreasing function on  $[1, \infty)$  and let  $a_n = f(n)$ . Then the series  $\sum_{n=1}^{\infty} a_n$  is convergent if and only if the improper integral  $\int_1^{\infty} f(x)dx$  is convergent. In other words,

① If  $\int_1^{\infty} f(x)dx$  is convergent, then  $\sum_{n=1}^{\infty} a_n$  is convergent.

② If  $\int_1^{\infty} f(x)dx$  is divergent, then  $\sum_{n=1}^{\infty} a_n$  is divergent.

Note that it is not necessary to start the series or the integral at  $n = 1$ . For instance, to test the series

$$\sum_{n=4}^{\infty} \frac{1}{(n-3)^2}$$

we would use

$$\int_4^{\infty} \frac{1}{(x-3)^2} dx$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.



## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^2 + 1} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} \left[ ? \right]_1^t \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.


$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\ &= \lim_{t \rightarrow \infty} (\arctan t - ?) \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\ &= \lim_{t \rightarrow \infty} (\arctan t - ?) \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned}
 \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\
 &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\
 &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right)
 \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned}
 \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\
 &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\
 &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\
 &= ? - \frac{\pi}{4}
 \end{aligned}$$






## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2+1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\ &= ? - \frac{\pi}{4} \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned}
 \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\
 &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\
 &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\
 &= ? - \frac{\pi}{4}
 \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\ &= \frac{\pi}{2} - \frac{\pi}{4} \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\ &= \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4} \end{aligned}$$




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned}
 \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\
 &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\
 &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\
 &= \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}
 \end{aligned}$$

Therefore  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  is ?




## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  for convergence.

$f(x) = \frac{1}{x^2 + 1}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so use the Integral Test.

$$\begin{aligned} \int_1^{\infty} \frac{1}{x^2 + 1} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{1}{x^2 + 1} dx \\ &= \lim_{t \rightarrow \infty} [\arctan x]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \arctan t - \frac{\pi}{4} \right) \\ &= \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4} \end{aligned}$$

Therefore  $\sum_{n=1}^{\infty} \frac{1}{n^2 + 1}$  is **convergent**.



## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = ?$



## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = ?$

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is ?

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{convergent} & \text{when ?} \\ \text{divergent} & \text{when ?} \end{cases}$$

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{convergent} & \text{when } p > 1 \\ \text{divergent} & \text{when } ? \end{cases}$$



## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{convergent} & \text{when } p > 1 \\ \text{divergent} & \text{when ?} \end{cases}$$

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{convergent} & \text{when } p > 1 \\ \text{divergent} & \text{when } p \leq 1 \end{cases}$$

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore for  $p \leq 0$  the series is divergent.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{convergent} & \text{when } p > 1 \\ \text{divergent} & \text{when } p \leq 1 \end{cases}$$

- $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$  is convergent when  $p > 1$  and divergent when  $p \leq 1$ .

## Example

For which values of  $p$  is the series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  convergent?

- If  $p < 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = \infty$ .
- If  $p = 0$ , then  $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 1$ .
- Therefore **for  $p \leq 0$  the series is divergent**.
- It remains to investigate the case  $p > 0$ . If  $p > 0$ , then  $f(x) = \frac{1}{x^p}$  is continuous, positive, and decreasing on  $[1, \infty)$ , so we can use the Integral Test.

$$\int_1^{\infty} \frac{1}{x^p} dx = \begin{cases} \text{convergent} & \text{when } p > 1 \\ \text{divergent} & \text{when } p \leq 1 \end{cases}$$

- $\Rightarrow \sum_{n=1}^{\infty} \frac{1}{n^p}$  is convergent when  $p > 1$  and **divergent when  $p \leq 1$** .

This theorem summarizes the results of the previous example.

### Theorem ( $p$ -series Convergence)

*The  $p$ -series  $\sum_{n=1}^{\infty} \frac{1}{n^p}$  is convergent if  $p > 1$  and divergent if  $p \leq 1$ .*

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.



## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > ?$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\int_1^{\infty} \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\int_1^{\infty} \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ ? \right]_1^t$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\int_1^{\infty} \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t$$



## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - ? \right) \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - ? \right) \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - 0 \right) \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - 0 \right) = ? \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - 0 \right) = \infty \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - 0 \right) = \infty \end{aligned}$$

Therefore  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  is ?

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  for convergence.

- $f(x) = \frac{\ln x}{x}$  is continuous and positive ( $x > 0$ ).
- To establish where  $f(x)$  is decreasing, take the derivative.

$$f'(x) = \frac{\left(\frac{1}{x}\right)(x) - (\ln x)(1)}{x^2} = \frac{1 - \ln x}{x^2}$$

- This is negative for all  $x > e$ .
- Therefore  $f$  is decreasing for all  $x > e$ .

$$\begin{aligned} \int_1^{\infty} \frac{\ln x}{x} dx &= \lim_{t \rightarrow \infty} \int_1^t \frac{\ln x}{x} dx = \lim_{t \rightarrow \infty} \left[ \frac{(\ln x)^2}{2} \right]_1^t \\ &= \lim_{t \rightarrow \infty} \left( \frac{1}{2} (\ln t)^2 - 0 \right) = \infty \end{aligned}$$

Therefore  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  is **divergent**.

# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.



# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.
- Any partial sum  $s_n$  is an approximation. But how good?

# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.
- Any partial sum  $s_n$  is an approximation. But how good?
- Estimate the size of the remainder
$$R_n = s - s_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots$$

# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.
- Any partial sum  $s_n$  is an approximation. But how good?
- Estimate the size of the remainder
$$R_n = s - s_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots$$
- Suppose  $f(n) = a_n$ . Draw rectangles with heights  $a_{n+1}, a_{n+2}, \dots$

# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.
- Any partial sum  $s_n$  is an approximation. But how good?
- Estimate the size of the remainder
$$R_n = s - s_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots$$
- Suppose  $f(n) = a_n$ . Draw rectangles with heights  $a_{n+1}, a_{n+2}, \dots$
- Use the right endpoints to find the height: then the rectangles are under the curve  $y = f(x)$ .
- Use the left endpoints to find the height: then the rectangles are above the curve  $y = f(x)$ .

# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.
- Any partial sum  $s_n$  is an approximation. But how good?
- Estimate the size of the remainder
$$R_n = s - s_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots$$
- Suppose  $f(n) = a_n$ . Draw rectangles with heights  $a_{n+1}, a_{n+2}, \dots$
- Use the right endpoints to find the height: then the rectangles are under the curve  $y = f(x)$ .
- $R_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots \leq \int_n^\infty f(x)dx$ .
- Use the left endpoints to find the height: then the rectangles are above the curve  $y = f(x)$ .

# Estimating the Sum of a Series

- Suppose we have already used the Integral Test to show that  $\sum a_n$  converges.
- Now we want to find an approximation to the sum of the series.
- Any partial sum  $s_n$  is an approximation. But how good?
- Estimate the size of the remainder
$$R_n = s - s_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots$$
- Suppose  $f(n) = a_n$ . Draw rectangles with heights  $a_{n+1}, a_{n+2}, \dots$
- Use the right endpoints to find the height: then the rectangles are under the curve  $y = f(x)$ .
- $R_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots \leq \int_n^\infty f(x)dx$ .
- Use the left endpoints to find the height: then the rectangles are above the curve  $y = f(x)$ .
- $R_n = a_{n+1} + a_{n+2} + a_{n+3} + \cdots \geq \int_{n+1}^\infty f(x)dx$ .

## Remainder Estimate for the Integral Test

Suppose  $f(k) = a_k$ , where  $f$  is continuous, positive, and decreasing for  $x \geq n$ , and  $\sum a_k$  is convergent with sum  $s$ . If  $R_n = s - s_n$ , then

$$\int_{n+1}^{\infty} f(x)dx \leq R_n \leq \int_n^{\infty} f(x)dx$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?



## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^{\infty} \frac{1}{x^3} dx =$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^{\infty} \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^\infty \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) =$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^{\infty} \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2n^2}$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^\infty \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \approx s_{10} = \frac{1}{1^3} + \frac{1}{2^3} + \cdots + \frac{1}{10^3} \approx 1.975$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^\infty \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \approx s_{10} = \frac{1}{1^3} + \frac{1}{2^3} + \cdots + \frac{1}{10^3} \approx 1.975$$

$$R_{10} \leq \int_{10}^{\infty} \frac{1}{x^3} dx =$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_{\textcolor{red}{n}}^{\infty} \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2\textcolor{red}{n}^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \approx s_{10} = \frac{1}{1^3} + \frac{1}{2^3} + \cdots + \frac{1}{10^3} \approx 1.975$$

$$R_{10} \leq \int_{\textcolor{red}{10}}^{\infty} \frac{1}{x^3} dx = \frac{1}{2(\textcolor{red}{10})^2}$$

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^{\infty} \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \approx s_{10} = \frac{1}{1^3} + \frac{1}{2^3} + \cdots + \frac{1}{10^3} \approx 1.975$$

$$R_{10} \leq \int_{10}^{\infty} \frac{1}{x^3} dx = \frac{1}{2(10)^2} = \frac{1}{200}$$



## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^{\infty} \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \approx s_{10} = \frac{1}{1^3} + \frac{1}{2^3} + \cdots + \frac{1}{10^3} \approx 1.975$$

$$R_{10} \leq \int_{10}^{\infty} \frac{1}{x^3} dx = \frac{1}{2(10)^2} = \frac{1}{200}$$

Therefore the error is at most 0.005.

## Example (Example 5, p. 737)

Approximate the sum of  $\sum \frac{1}{n^3}$  using the first 10 terms. Estimate the error involved in this approximation. How many terms are required to get an accuracy of 0.0005 or better?

$$\int_n^{\infty} \frac{1}{x^3} dx = \lim_{t \rightarrow \infty} \left[ -\frac{1}{2x^2} \right]_n^t = \lim_{t \rightarrow \infty} \left( -\frac{1}{2t^2} + \frac{1}{2n^2} \right) = \frac{1}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{1}{n^3} \approx s_{10} = \frac{1}{1^3} + \frac{1}{2^3} + \cdots + \frac{1}{10^3} \approx 1.975$$

$$R_{10} \leq \int_{10}^{\infty} \frac{1}{x^3} dx = \frac{1}{2(10)^2} = \frac{1}{200}$$

Therefore the error is at most 0.005.

To get an accuracy of 0.0005 or better, we want  $R_n \leq 0.0005$ . Since  $R_n \leq \frac{1}{2n^2}$ , we want

$$\frac{1}{2n^2} \leq 0.0005, \quad \text{or} \quad n \geq \sqrt{1000} \approx 31.6$$

$$\int_{n+1}^{\infty} f(x)dx \leq R_n \leq \int_n^{\infty} f(x)dx$$

$$\begin{array}{ccccc} \int_{n+1}^{\infty} f(x)dx & \leq & R_n & \leq & \int_n^{\infty} f(x)dx \\ \textcolor{red}{S}_n + \int_{n+1}^{\infty} f(x)dx & \leq & \textcolor{red}{S}_n + R_n & \leq & \textcolor{red}{S}_n + \int_n^{\infty} f(x)dx \end{array}$$

- Add  $s_n$  to both sides of both inequalities.

$$\begin{array}{rclclcl}
 \int_{n+1}^{\infty} f(x)dx & \leq & R_n & \leq & \int_n^{\infty} f(x)dx \\
 s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s_n + R_n & \leq & s_n + \int_n^{\infty} f(x)dx \\
 s_n + \int_{n+1}^{\infty} f(x)dx & \leq & & \leq & s_n + \int_n^{\infty} f(x)dx
 \end{array}$$

- Add  $s_n$  to both sides of both inequalities.

$$\begin{array}{rclclcl}
 \int_{n+1}^{\infty} f(x)dx & \leq & R_n & \leq & \int_n^{\infty} f(x)dx \\
 s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s_n + R_n & \leq & s_n + \int_n^{\infty} f(x)dx \\
 s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s & \leq & s_n + \int_n^{\infty} f(x)dx
 \end{array}$$

- Add  $s_n$  to both sides of both inequalities.

$$\begin{array}{rclclcl} \int_{n+1}^{\infty} f(x)dx & \leq & R_n & \leq & \int_n^{\infty} f(x)dx \\ s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s_n + R_n & \leq & s_n + \int_n^{\infty} f(x)dx \\ s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s & \leq & s_n + \int_n^{\infty} f(x)dx \end{array}$$

- Add  $s_n$  to both sides of both inequalities.
- This gives upper and lower bounds for  $s$ .

$$\begin{array}{rclclcl} \int_{n+1}^{\infty} f(x)dx & \leq & R_n & \leq & \int_n^{\infty} f(x)dx \\ s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s_n + R_n & \leq & s_n + \int_n^{\infty} f(x)dx \\ s_n + \int_{n+1}^{\infty} f(x)dx & \leq & s & \leq & s_n + \int_n^{\infty} f(x)dx \end{array}$$

- Add  $s_n$  to both sides of both inequalities.
- This gives upper and lower bounds for  $s$ .
- This is a better approximation than just using  $s_n$ .



# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a =$  and  $r =$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r =$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r =$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is



# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} \qquad \frac{1}{2^i}$$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^n+1}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} < \frac{1}{2^i}$$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} < \frac{1}{2^i}$$
$$\sum_{i=1}^n \frac{1}{2^i + 1} < \sum_{i=1}^n \frac{1}{2^i}$$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} < \frac{1}{2^i}$$
$$\sum_{i=1}^n \frac{1}{2^i + 1} < \sum_{i=1}^n \frac{1}{2^i} < \sum_{i=1}^{\infty} \frac{1}{2^i}$$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^{i+1}} < \frac{1}{2^i}$$
$$\sum_{i=1}^n \frac{1}{2^{i+1}} < \sum_{i=1}^n \frac{1}{2^i} < \sum_{i=1}^{\infty} \frac{1}{2^i} =$$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\begin{aligned} \frac{1}{2^i + 1} &< \frac{1}{2^i} \\ \sum_{i=1}^n \frac{1}{2^i + 1} &< \sum_{i=1}^n \frac{1}{2^i} < \sum_{i=1}^{\infty} \frac{1}{2^i} = 1 \end{aligned}$$

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} < \frac{1}{2^i}$$
$$\sum_{i=1}^n \frac{1}{2^i + 1} < \sum_{i=1}^n \frac{1}{2^i} < \sum_{i=1}^{\infty} \frac{1}{2^i} = 1$$

- The partial sums of  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$  are increasing and are bounded above by 1.



# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} < \frac{1}{2^i}$$
$$\sum_{i=1}^n \frac{1}{2^i + 1} < \sum_{i=1}^n \frac{1}{2^i} < \sum_{i=1}^{\infty} \frac{1}{2^i} = 1$$

- The partial sums of  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$  are increasing and are bounded above by 1.
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$  is

# The Comparison Tests

- In the Comparison Tests, the idea is to compare a given series with another series that is known to be convergent or divergent.
- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$ .
- This reminds us of the series  $\sum_{n=1}^{\infty} \frac{1}{2^n}$ .
- $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is a geometric series with  $a = \frac{1}{2}$  and  $r = \frac{1}{2}$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^n}$  is convergent.

$$\frac{1}{2^i + 1} < \frac{1}{2^i}$$
$$\sum_{i=1}^n \frac{1}{2^i + 1} < \sum_{i=1}^n \frac{1}{2^i} < \sum_{i=1}^{\infty} \frac{1}{2^i} = 1$$

- The partial sums of  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$  are increasing and are bounded above by 1.
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{2^{n+1}}$  is convergent.

## Theorem (The Comparison Test)

*Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.*

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.*
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.*

## Theorem (The Comparison Test)

*Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.*

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.*
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.*

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series
- A geometric series

## Theorem (The Comparison Test)

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  **converges if**                      and diverges if                      )
- A geometric series

## Theorem (The Comparison Test)

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  **converges if  $p > 1$**  and diverges if )
- A geometric series

## Theorem (The Comparison Test)

*Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.*

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.*
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.*

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  converges if  $p > 1$  and **diverges if** )
- A geometric series

## Theorem (The Comparison Test)

*Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.*

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.*
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.*

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  converges if  $p > 1$  and **diverges if  $p \leq 1$** )
- A geometric series



## Theorem (The Comparison Test)

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  converges if  $p > 1$  and diverges if  $p \leq 1$ )
- A geometric series ( $\sum ar^{n-1}$  **converges if** and diverges if )

## Theorem (The Comparison Test)

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  converges if  $p > 1$  and diverges if  $p \leq 1$ )
- A geometric series ( $\sum ar^{n-1}$  converges if  $|r| < 1$  and diverges if  $|r| \geq 1$ )

## Theorem (The Comparison Test)

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  converges if  $p > 1$  and diverges if  $p \leq 1$ )
- A geometric series ( $\sum ar^{n-1}$  converges if  $|r| < 1$  and **diverges if**  
)

## Theorem (The Comparison Test)

Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms.

- 1 If  $\sum b_n$  is convergent and  $a_n \leq b_n$  for all  $n$ , then  $\sum a_n$  is also convergent.
- 2 If  $\sum b_n$  is divergent and  $a_n \geq b_n$  for all  $n$ , then  $\sum a_n$  is also divergent.

When we use the Comparison Test, we need to have some series  $\sum b_n$  that we know in order to make a comparison. Usually  $\sum b_n$  is one of

- A  $p$ -series ( $\sum \frac{1}{n^p}$  converges if  $p > 1$  and diverges if  $p \leq 1$ )
- A geometric series ( $\sum ar^{n-1}$  converges if  $|r| < 1$  and **diverges if  $|r| \geq 1$** )

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} \quad \frac{5}{2n^2}$$

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$



## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a constant times a  $p$ -series with  $p =$

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a constant times a  $p$ -series with  $p = 2 > 1$ .

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a constant times a  $p$ -series with  $p = 2 > 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{5}{2n^2}$  is

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a constant times a  $p$ -series with  $p = 2 > 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{5}{2n^2}$  is convergent.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a constant times a  $p$ -series with  $p = 2 > 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{5}{2n^2}$  is convergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  is by the Comparison Test.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  converges or diverges.

- As  $n \rightarrow \infty$ , the dominant term in the denominator is  $2n^2$ , so compare with  $\frac{5}{2n^2}$ .

$$\frac{5}{2n^2 + 7n + 3} < \frac{5}{2n^2}$$

$$\sum_{n=1}^{\infty} \frac{5}{2n^2} = \frac{5}{2} \sum_{n=1}^{\infty} \frac{1}{n^2}$$

- This is a constant times a  $p$ -series with  $p = 2 > 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{5}{2n^2}$  is convergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{5}{2n^2+7n+3}$  is convergent by the Comparison Test.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.



## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

- $\sum_{n=1}^{\infty} \frac{1}{n}$  is a  $p$ -series with  $p =$

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

- $\sum_{n=1}^{\infty} \frac{1}{n}$  is a  $p$ -series with  $p = 1$ .

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

- $\sum_{n=1}^{\infty} \frac{1}{n}$  is a  $p$ -series with  $p = 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{n}$  is

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

- $\sum_{n=1}^{\infty} \frac{1}{n}$  is a  $p$ -series with  $p = 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{n}$  is divergent.

## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

- $\sum_{n=1}^{\infty} \frac{1}{n}$  is a  $p$ -series with  $p = 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{n}$  is divergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  is **divergent** by the Comparison Test.



## Example

Determine if  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  converges or diverges.

- We could use the Integral Test to find this.
- The Comparison Test is even easier.

$$\frac{\ln n}{n} > \frac{1}{n} \quad \text{if } n \geq 3$$

- $\sum_{n=1}^{\infty} \frac{1}{n}$  is a  $p$ -series with  $p = 1$ .
- Therefore  $\sum_{n=1}^{\infty} \frac{1}{n}$  is divergent.
- Therefore  $\sum_{n=1}^{\infty} \frac{\ln n}{n}$  is divergent by the Comparison Test.

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 **smaller** than the terms of a **convergent** series, or
- 2 **bigger** than the terms of a **divergent** series.

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 **smaller** than the terms of a **convergent** series, or
- 2 **bigger** than the terms of a **divergent** series.

If the terms  $a_n$  are

- 1 **bigger** than the terms of a **convergent** series, or
- 2 **smaller** than the terms of a **divergent** series,

then the Comparison Test gives no information.

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 smaller than the terms of a convergent series, or
- 2 bigger than the terms of a divergent series.

If the terms  $a_n$  are

- 1 bigger than the terms of a convergent series, or
- 2 smaller than the terms of a divergent series,

then the Comparison Test gives no information.

- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$ .

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 smaller than the terms of a convergent series, or
- 2 bigger than the terms of a divergent series.

If the terms  $a_n$  are

- 1 bigger than the terms of a convergent series, or
- 2 smaller than the terms of a divergent series,

then the Comparison Test gives no information.

- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$ .

$$\frac{1}{2^n - 1} \quad \frac{1}{2^n}$$

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 smaller than the terms of a convergent series, or
- 2 bigger than the terms of a divergent series.

If the terms  $a_n$  are

- 1 bigger than the terms of a convergent series, or
- 2 smaller than the terms of a divergent series,

then the Comparison Test gives no information.

- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$ .

$$\frac{1}{2^n - 1} > \frac{1}{2^n}$$

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 smaller than the terms of a convergent series, or
- 2 bigger than the terms of a divergent series.

If the terms  $a_n$  are

- 1 bigger than the terms of a convergent series, or
- 2 smaller than the terms of a divergent series,

then **the Comparison Test gives no information.**

- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$ .

$$\frac{1}{2^n - 1} > \frac{1}{2^n}$$

- The Comparison Test tells us nothing here.

In order to use the comparison test to see if  $\sum a_n$  is convergent or divergent, we need the terms  $a_n$  to be

- 1 smaller than the terms of a convergent series, or
- 2 bigger than the terms of a divergent series.

If the terms  $a_n$  are

- 1 bigger than the terms of a convergent series, or
- 2 smaller than the terms of a divergent series,

then the Comparison Test gives no information.

- Consider the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$ .

$$\frac{1}{2^n - 1} > \frac{1}{2^n}$$

- The Comparison Test tells us nothing here.
- Nevertheless, we think  $\sum \frac{1}{2^n - 1}$  should converge, because it's so close to  $\sum \frac{1}{2^n}$ .



## Theorem (The Limit Comparison Test)

*Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms. If*

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$$

*where  $c$  is a finite number and  $c > 0$ , then either both series converge or both series diverge.*

## Theorem (The Limit Comparison Test)

*Suppose that  $\sum a_n$  and  $\sum b_n$  are series with positive terms. If*

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$$

*where  **$c$  is a finite number and  $c > 0$** , then either both series converge or both series diverge.*

The main thing to check is that  $c$  is finite and non-zero.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^{n-1}}$  for convergence or divergence.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \cdot \frac{1}{2^n} \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \cdot \frac{1}{2^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{1}{2^n}} \end{aligned}$$



## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \cdot \frac{1}{2^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{1}{2^n}} = 1 > 0 \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \cdot \frac{1}{2^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{1}{2^n}} = 1 > 0 \end{aligned}$$

- $\sum \frac{1}{2^n}$  is a geometric series.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \cdot \frac{1}{2^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{1}{2^n}} = 1 > 0 \end{aligned}$$

- $\sum \frac{1}{2^n}$  is a convergent geometric series.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{1}{2^n - 1}$  for convergence or divergence.  
Use the Limit Comparison Test with

$$a_n = \frac{1}{2^n - 1}, \quad b_n = \frac{1}{2^n}$$

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{\frac{1}{2^n - 1}}{\frac{1}{2^n}} \\ &= \lim_{n \rightarrow \infty} \frac{2^n}{2^n - 1} \cdot \frac{1}{2^n} \\ &= \lim_{n \rightarrow \infty} \frac{1}{1 - \frac{1}{2^n}} = 1 > 0 \end{aligned}$$

- $\sum \frac{1}{2^n}$  is a convergent geometric series.
- By the Limit Comparison Test  $\sum \frac{1}{2^n - 1}$  is convergent too.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7} + n^5}$  for convergence or divergence.

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is and the dominant part of the denominator is

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \text{---}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{\sqrt{7 + n^5}}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{\sqrt{7 + n^5}}$$



## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$a_n = \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, \quad b_n = \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}}$$
$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned} a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\ \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned} a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\ \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}} \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned} a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\ \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}} \\ &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned} a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\ \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}} \\ &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0 \end{aligned}$$

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^2}$  is a constant multiple of a  $p$ -series with  $p =$



## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \frac{\frac{1}{n^{5/2}}}{\frac{1}{n^{5/2}}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^{\frac{1}{2}}}$  is a constant multiple of a  $p$ -series with  $p = \frac{1}{2}$ .

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned} a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\ \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\ &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0 \end{aligned}$$

- $\sum \frac{2}{n^2}$  is a constant multiple of a  $p$ -series with  $p = \frac{1}{2}$ .
- Therefore  $\sum \frac{2}{n^2}$  is

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned} a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\ \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\ &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0 \end{aligned}$$

- $\sum \frac{2}{n^2}$  is a constant multiple of a  $p$ -series with  $p = \frac{1}{2}$ .
- Therefore  $\sum \frac{2}{n^2}$  is divergent

## Example

Test the series  $\sum_{n=1}^{\infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$  for convergence or divergence.

- The dominant part of the numerator is  $2n^2$  and the dominant part of the denominator is  $\sqrt{n^5} = n^{5/2}$ .

$$\begin{aligned}
 a_n &= \frac{2n^2 + 3n}{\sqrt{7 + n^5}}, & b_n &= \frac{2n^2}{n^{5/2}} = \frac{2}{n^{1/2}} \\
 \lim_{n \rightarrow \infty} \frac{a_n}{b_n} &= \lim_{n \rightarrow \infty} \frac{2n^2 + 3n}{\sqrt{7 + n^5}} \cdot \frac{n^{1/2}}{2} = \lim_{n \rightarrow \infty} \frac{2n^{5/2} + 3n^{3/2}}{2\sqrt{7 + n^5}} \cdot \frac{1}{n^{5/2}} \\
 &= \lim_{n \rightarrow \infty} \frac{2 + \frac{3}{n}}{2\sqrt{\frac{7}{n^5} + 1}} = 1 > 0
 \end{aligned}$$

- $\sum \frac{2}{n^2}$  is a constant multiple of a  $p$ -series with  $p = \frac{1}{2}$ .
- Therefore  $\sum \frac{2}{n^2}$  is divergent, and so is  $\sum \frac{2n^2 + 3n}{\sqrt{7 + n^5}}$ .