

Calculus III

Lecture 12

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1 Minima, Maxima

Outline

- 1 Minima, Maxima
- 2 Lagrange Multipliers

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
`https://creativecommons.org/licenses/by/3.0/us/`
and the links therein.

Minima, Maxima

Function $f: D \rightarrow \mathbb{R}$ defined on a region D in \mathbb{R}^2 , we want to know:

- The largest and the smallest values of f attained on D , if any;
- The points where these extreme values are attained.

Minima, Maxima

Function $f: D \rightarrow \mathbb{R}$ defined on a region D in \mathbb{R}^2 , we want to know:

- The largest and the smallest values of f attained on D , if any;
- The points where these extreme values are attained.

A point P_0 in D is a point of:

- absolute maximum, if $f(P) \leq f(P_0)$ for all P in D ;
- absolute minimum, if $f(P) \geq f(P_0)$ for all P in D .

Minima, Maxima

Function $f: D \rightarrow \mathbb{R}$ defined on a region D in \mathbb{R}^2 , we want to know:

- The largest and the smallest values of f attained on D , if any;
- The points where these extreme values are attained.

A point P_0 in D is a point of:

- absolute maximum, if $f(P) \leq f(P_0)$ for all P in D ;
- absolute minimum, if $f(P) \geq f(P_0)$ for all P in D .

These notions are relative to the domain D . A point P_0 that is not an extreme point might become one if we focus only around that point.

Minima, Maxima

Function $f: D \rightarrow \mathbb{R}$ defined on a region D in \mathbb{R}^2 , we want to know:

- The largest and the smallest values of f attained on D , if any;
- The points where these extreme values are attained.

A point P_0 in D is a point of:

- absolute maximum, if $f(P) \leq f(P_0)$ for all P in D ;
- absolute minimum, if $f(P) \geq f(P_0)$ for all P in D .

These notions are relative to the domain D . A point P_0 that is not an extreme point might become one if we focus only around that point.

A point P_0 in D is a point of:

- local maximum, if there exists an open disk $B = B_r(P_0)$ centered at P_0 such that $f(P) \leq f(P_0)$ for all P in $B \cap D$;
- local minimum, if there exists an open disk $B = B_r(P_0)$ centered at P_0 such that $f(P) \geq f(P_0)$ for all P in $B \cap D$.

Minima, Maxima

Function $f: D \rightarrow \mathbb{R}$ defined on a region D in \mathbb{R}^2 , we want to know:

- The largest and the smallest values of f attained on D , if any;
- The points where these extreme values are attained.

A point P_0 in D is a point of:

- absolute maximum, if $f(P) \leq f(P_0)$ for all P in D ;
- absolute minimum, if $f(P) \geq f(P_0)$ for all P in D .

These notions are relative to the domain D . A point P_0 that is not an extreme point might become one if we focus only around that point.

A point P_0 in D is a point of:

- local maximum, if there exists an open disk $B = B_r(P_0)$ centered at P_0 such that $f(P) \leq f(P_0)$ for all P in $B \cap D$;
- local minimum, if there exists an open disk $B = B_r(P_0)$ centered at P_0 such that $f(P) \geq f(P_0)$ for all P in $B \cap D$.

How do we find points of extreme?

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.
If:

- P_0 is a point of extreme (minimum or maximum);
- P_0 is an *interior point* of D , which means that there exists an open disk centered at P_0 and completely included in D ;
- directional derivatives at P_0 exist in all directions

then

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.
If:

- P_0 is a point of extreme (minimum or maximum);
- P_0 is an *interior point* of D , which means that there exists an open disk centered at P_0 and completely included in D ;
- directional derivatives at P_0 exist in all directions

then $(\nabla f)(P_0) = \mathbf{0}$. In particular, $f_x(P_0) = f_y(P_0) = 0$.

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.
If:

- P_0 is a point of extreme (minimum or maximum);
- P_0 is an *interior point* of D , which means that there exists an open disk centered at P_0 and completely included in D ;
- directional derivatives at P_0 exist in all directions

then $(\nabla f)(P_0) = \mathbf{0}$. In particular, $f_x(P_0) = f_y(P_0) = 0$.

Geometric Interpretation:

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.
If:

- P_0 is a point of extreme (minimum or maximum);
- P_0 is an *interior point* of D , which means that there exists an open disk centered at P_0 and completely included in D ;
- directional derivatives at P_0 exist in all directions

then $(\nabla f)(P_0) = \mathbf{0}$. In particular, $f_x(P_0) = f_y(P_0) = 0$.

Geometric Interpretation: At an interior point of extreme, the tangent plane to the graph surface is horizontal.

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.
If:

- P_0 is a point of extreme (minimum or maximum);
- P_0 is an *interior point* of D , which means that there exists an open disk centered at P_0 and completely included in D ;
- directional derivatives at P_0 exist in all directions

then $(\nabla f)(P_0) = \mathbf{0}$. In particular, $f_x(P_0) = f_y(P_0) = 0$.

Geometric Interpretation: At an interior point of extreme, the tangent plane to the graph surface is horizontal.

The converse is not true:

Critical Points

If $\mathbf{u} = (\nabla f)(P_0)$ exists and is non-zero, then

- f increases along \mathbf{u} ;
- f decreases along $-\mathbf{u}$;

If we can move along $\pm\mathbf{u}$ and stay in D , then P_0 is not an extreme.
If:

- P_0 is a point of extreme (minimum or maximum);
- P_0 is an *interior point* of D , which means that there exists an open disk centered at P_0 and completely included in D ;
- directional derivatives at P_0 exist in all directions

then $(\nabla f)(P_0) = \mathbf{0}$. In particular, $f_x(P_0) = f_y(P_0) = 0$.

Geometric Interpretation: At an interior point of extreme, the tangent plane to the graph surface is horizontal.

The converse is not true: if $f_x(P_0) = f_y(P_0) = 0$, then P_0 is not necessarily a point of extreme.

Where else can one find extreme points?

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D .
Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D . Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Examples:

- D =open unit disk \implies

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D .
Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Examples:

- D =open unit disk \implies set of boundary points = unit circle;

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D . Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Examples:

- D =open unit disk \implies set of boundary points = unit circle;
- D =closed unit disk \implies

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D . Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Examples:

- D =open unit disk \implies set of boundary points = unit circle;
- D =closed unit disk \implies set of boundary points = unit circle;

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D . Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Examples:

- D =open unit disk \implies set of boundary points = unit circle;
- D =closed unit disk \implies set of boundary points = unit circle;

Notice that a boundary point may or may not be included in D .

Where else can one find extreme points?

- At points P_0 where some directional derivatives do not exist (suffices that one of $f_x(P_0)$ or $f_y(P_0)$ does not exist.);
- At points P_0 in D that are not interior points of D .

Important concept: A point P in \mathbb{R}^2 is a *boundary point* for a region D if every open disk centered at P has points both in D and outside of D . Similar definition for \mathbb{R}^3 , but replace open disk with open ball.

Examples:

- D =open unit disk \implies set of boundary points = unit circle;
- D =closed unit disk \implies set of boundary points = unit circle;

Notice that a boundary point may or may not be included in D .

Strategy for finding extreme points:

- Check the *critical points* of f :
 - Points P_0 for which $f_x(P_0)$ or $f_y(P_0)$ does not exist;
 - Points P_0 for which $f_x(P_0) = f_y(P_0) = 0$.
- Check boundary points included in the domain.

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior;

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.
- It remains to find the points (x, y) for which $f_x(x, y) = f_y(x, y) = 0$.

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{cases} \iff \begin{cases} x^3 = y \\ y^3 = x \end{cases}$$

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.
- It remains to find the points (x, y) for which $f_x(x, y) = f_y(x, y) = 0$.

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{cases} \iff \begin{cases} x^3 = y \\ y^3 = x \end{cases}$$

- This system is a non-linear, haven't studied methods for those.

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.
- It remains to find the points (x, y) for which $f_x(x, y) = f_y(x, y) = 0$.

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{cases} \iff \begin{cases} x^3 = y \\ y^3 = x \end{cases}$$

- This system is a non-linear, haven't studied methods for those. This system can be solved using ad-hoc methods.

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.
- It remains to find the points (x, y) for which $f_x(x, y) = f_y(x, y) = 0$.

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{cases} \iff \begin{cases} x^3 = y \\ y^3 = x \end{cases}$$

- This system is a non-linear, haven't studied methods for those. This system can be solved using ad-hoc methods.
- There are three values of x that work:

$$x = 0 \implies y = 0 \implies \text{Point } (0, 0)$$

$$x = 1 \implies y = 1 \implies \text{Point } (1, 1)$$

$$x = -1 \implies y = -1 \implies \text{Point } (-1, -1)$$

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.
- It remains to find the points (x, y) for which $f_x(x, y) = f_y(x, y) = 0$.

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{cases} \iff \begin{cases} x^3 = y \\ y^3 = x \end{cases}$$

- This system is a non-linear, haven't studied methods for those. This system can be solved using ad-hoc methods.
- There are three values of x that work:

$$x = 0 \implies y = 0 \implies \text{Point } (0, 0)$$

$$x = 1 \implies y = 1 \implies \text{Point } (1, 1)$$

$$x = -1 \implies y = -1 \implies \text{Point } (-1, -1)$$

Typical mistake:

Example

Find the critical points of $f(x, y) = x^4 + y^4 - 4xy$ on $D = \mathbb{R}^2$.

- All points are interior; the function is differentiable everywhere.
- It remains to find the points (x, y) for which $f_x(x, y) = f_y(x, y) = 0$.

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} 4x^3 - 4y = 0 \\ 4y^3 - 4x = 0 \end{cases} \iff \begin{cases} x^3 = y \\ y^3 = x \end{cases}$$

- This system is a non-linear, haven't studied methods for those. This system can be solved using ad-hoc methods.
- There are three values of x that work:

$$x = 0 \implies y = 0 \implies \text{Point } (0, 0)$$

$$x = 1 \implies y = 1 \implies \text{Point } (1, 1)$$

$$x = -1 \implies y = -1 \implies \text{Point } (-1, -1)$$

Typical mistake: $x^9 = x \iff x^8 = 1$.

Second Derivative Test

When is an interior critical point a pt. of min/max?

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows.

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix}$$

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. **Example:?**

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. **Example:** crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. **Example: ?**

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. **Example:** crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. Example: crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.
- If $D(x_0, y_0) < 0$, then (x_0, y_0) is neither a minimum nor a maximum. Such points are called *saddle points*.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. Example: crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.
- If $D(x_0, y_0) < 0$, then (x_0, y_0) is neither a minimum nor a maximum. Such points are called *saddle points*. Example:?

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. Example: crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.
- If $D(x_0, y_0) < 0$, then (x_0, y_0) is neither a minimum nor a maximum. Such points are called *saddle points*. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 - y^2$.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. Example: crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.
- If $D(x_0, y_0) < 0$, then (x_0, y_0) is neither a minimum nor a maximum. Such points are called *saddle points*. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 - y^2$.
- If $D(x_0, y_0) = 0$, then the test is inconclusive.

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. Example: crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.
- If $D(x_0, y_0) < 0$, then (x_0, y_0) is neither a minimum nor a maximum. Such points are called *saddle points*. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 - y^2$.
- If $D(x_0, y_0) = 0$, then the test is inconclusive. Examples: ?

Second Derivative Test

When is an interior critical point a pt. of min/max? Define the *Hessian matrix* H of f as follows. Denote by D the determinant of H .

$$H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} \quad D = \det H = f_{xx}f_{yy} - f_{xy}^2$$

Test: Let $P(x_0, y_0)$ be an interior critical point of f and suppose that f has continuous second order derivatives around P .

- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) > 0$, then (x_0, y_0) is a local minimum. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 + y^2$.
- If $D(x_0, y_0) > 0$ and $f_{xx}(x_0, y_0) < 0$, then (x_0, y_0) is a local maximum. Example: crit. pt. $(0, 0)$ for $f(x, y) = -x^2 - y^2$.
- If $D(x_0, y_0) < 0$, then (x_0, y_0) is neither a minimum nor a maximum. Such points are called *saddle points*. Example: crit. pt. $(0, 0)$ for $f(x, y) = x^2 - y^2$.
- If $D(x_0, y_0) = 0$, then the test is inconclusive. **Examples:**

$$x^4 + y^4, -x^4 - y^4, x^4 - y^4.$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} =$$

$$f_{xy} =$$

$$f_{yy} =$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} =$$

$$f_{xy} =$$

$$f_{yy} =$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} =$$

$$f_{yy} =$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} =$$

$$f_{yy} =$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} =$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} =$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 =$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.


$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$					
$(1, 1)$					
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$?	?	?	?	
$(1, 1)$					
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	
$(1, 1)$					
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$?
$(1, 1)$					
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$					
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$?	?	?	?	
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$?
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$					

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$?	?	?	?	

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$	

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$?

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$	Local min

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$	Local min

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$	Local min

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$


$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$	Local min

Example (Finding maxima, minima)



Find the local and global maxima and minima of $f(x, y) = x^4 + y^4 - 4xy$.

$$f_{xx} = 12x^2$$

$$f_{xy} = -4$$

$$f_{yy} = 12y^2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = 144x^2y^2 - 16.$$

The critical points were previously computed.

(x_0, y_0)	f_{xx}	f_{yy}	f_{xy}	D	Extremum ?
$(0, 0)$	0	0	-4	$-16 < 0$	Saddle point
$(1, 1)$	12	12	-4	$128 > 0$	Local min
$(-1, -1)$	12	12	-4	$128 > 0$	Local min

In this case it turns out that the two local minimum points are actually global minimum points, because

$$f(x, y) = x^4 + y^4 - 4xy = (x^2 - 1)^2 + (y^2 - 1)^2 + 2(x - y)^2 - 2 \geq -2.$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} .

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$$d = \sqrt{(x - 2)^2 + (y - 1)^2 + z^2},$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x - 2)^2 + (y - 1)^2 + z^2}$, equivalently to minimize $f = d^2$:

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2 .$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) =$$

$$0 = f_y(x, y) =$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2 .$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = ?$$

$$0 = f_y(x, y) =$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) =$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = ?$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, **solve the system**:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, **solve the system**:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$$\frac{14}{5}y - 2 = 0. \text{ Finally } x = \frac{11}{7}, y = \frac{5}{7}.$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = ?$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = \begin{pmatrix} 20 & 12 \\ 12 & 10 \end{pmatrix}$.

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = \begin{pmatrix} 20 & 12 \\ 12 & 10 \end{pmatrix}$.

$D = \det H = ?$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = \begin{pmatrix} 20 & 12 \\ 12 & 10 \end{pmatrix}$.

$$D = \det H = 200 - 144 = 56$$

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = \begin{pmatrix} 20 & 12 \\ 12 & 10 \end{pmatrix}$.

$$D = \det H = 200 - 144 = 56$$

$$\text{at } x = \frac{11}{7}, y = \frac{5}{7},$$

Therefore we have a local ?

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = \begin{pmatrix} 20 > 0 & 12 \\ 12 & 10 \end{pmatrix}$.

$D = \det H = 200 - 144 = 56 > 0$. Therefore we have a local **minimum** at $x = \frac{11}{7}$, $y = \frac{5}{7}$,

Example

Let $P(2, 1, 0)$ and let \mathcal{P} be the plane $3x + 2y + z = 6$. Find the shortest distance between P and a point on \mathcal{P} .

Let $Q(x, y, z)$ be a point on \mathcal{P} . We seek to minimize

$d = \sqrt{(x-2)^2 + (y-1)^2 + z^2}$, equivalently to minimize $f = d^2$:

$$f(x, y) = (x-2)^2 + (y-1)^2 + z^2 = (x-2)^2 + (y-1)^2 + (6-3x-2y)^2.$$

To find the critical points, solve the system:

$$0 = f_x(x, y) = 2(x-2) - 6(6-3x-2y) = 20x + 12y - 40$$

$$0 = f_y(x, y) = 2(y-1) - 4(6-3x-2y) = 12x + 10y - 26$$

From first equation $x = \frac{10-3y}{5}$. Substitute in the second eqn.:

$\frac{14}{5}y - 2 = 0$. Finally $x = \frac{11}{7}$, $y = \frac{5}{7}$. To find whether $(\frac{11}{7}, \frac{5}{7})$ is local

extremum, compute Hessian: $H = \begin{pmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{pmatrix} = \begin{pmatrix} 20 & 12 \\ 12 & 10 \end{pmatrix}$.

$D = \det H = 200 - 144 = 56 > 0$. Therefore we have a local minimum at $x = \frac{11}{7}$, $y = \frac{5}{7}$, and the min. is: $f(\frac{11}{7}, \frac{5}{7}) = \frac{\sqrt{14}}{7}$.

Extreme Value Theorem

Global extreme points are guaranteed to exist if:

- $f: D \rightarrow \mathbb{R}$ is continuous, and
- the domain D has the following properties:
 - D is *bounded*: The points in D don't go farther than a certain fixed, finite distance from a fixed point.
 - D is *closed*: D contains all its boundary points.

The statement above is the **Extreme Value Theorem**.

Extreme Value Theorem

Global extreme points are guaranteed to exist if:

- $f: D \rightarrow \mathbb{R}$ is continuous, and
- the domain D has the following properties:
 - D is *bounded*: The points in D don't go farther than a certain fixed, finite distance from a fixed point.
 - D is *closed*: D contains all its boundary points.

The statement above is the **Extreme Value Theorem**.

- Why does D have to be bounded:

Extreme Value Theorem

Global extreme points are guaranteed to exist if:

- $f: D \rightarrow \mathbb{R}$ is continuous, and
- the domain D has the following properties:
 - D is *bounded*: The points in D don't go farther than a certain fixed, finite distance from a fixed point.
 - D is *closed*: D contains all its boundary points.

The statement above is the **Extreme Value Theorem**.

- Why does D have to be bounded: to exclude $f: \mathbb{R}^2 \rightarrow \mathbb{R}$,
 $f(x, y) = x$;

Extreme Value Theorem

Global extreme points are guaranteed to exist if:

- $f: D \rightarrow \mathbb{R}$ is continuous, and
- the domain D has the following properties:
 - D is *bounded*: The points in D don't go farther than a certain fixed, finite distance from a fixed point.
 - D is *closed*: D contains all its boundary points.

The statement above is the **Extreme Value Theorem**.

- Why does D have to be bounded: to exclude $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $f(x, y) = x$;
- Why does D have to be closed:

Extreme Value Theorem



Global extreme points are guaranteed to exist if:

- $f: D \rightarrow \mathbb{R}$ is continuous, and
- the domain D has the following properties:
 - D is *bounded*: The points in D don't go farther than a certain fixed, finite distance from a fixed point.
 - D is *closed*: D contains all its boundary points.

The statement above is the **Extreme Value Theorem**.



- Why does D have to be bounded: to exclude $f: \mathbb{R}^2 \rightarrow \mathbb{R}$, $f(x, y) = x$;
- Why does D have to be closed: to exclude $f: \mathbb{R}^2 \setminus \{(0, 0)\} \rightarrow \mathbb{R}$, $f(x, y) = (x^2 + y^2)^{-1}$. In this situation the boundary of D is $\{(0, 0)\}$ and is not included in D , so D is not closed.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.



Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z .



Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Example





Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10$.

Example





Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}.$

Example





Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$

Example





Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$

$(x, y) \in ?$

Example





Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}.$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.



Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.
 $(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x$$

$$0 = V_y$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.



Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$
 $(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = ?$$

$$0 = V_y$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.



Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.



Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = ?$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.



Restrictions: $xy + 2(zx + zy) = 10. \Rightarrow z = \frac{10-xy}{2(x+y)}. \Rightarrow V = xy \frac{10-xy}{2(x+y)}.$

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$



$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then

$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$



$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then

$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$



We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then

$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Therefore $x^2 = y^2$ and so $x = y$ (both quantities are positive).

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then



$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Therefore $x^2 = y^2$ and so $x = y$ (both quantities are positive).

Therefore $\frac{3}{2}x^2 = 5$,

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then



$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Therefore $x^2 = y^2$ and so $x = y$ (both quantities are positive).

Therefore $\frac{3}{2}x^2 = 5$, and so $x = \sqrt{\frac{10}{3}} = y$.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then



$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Therefore $x^2 = y^2$ and so $x = y$ (both quantities are positive).

Therefore $\frac{3}{2}x^2 = 5$, and so $x = \sqrt{\frac{10}{3}} = y$. By EVT max exists

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then

$$0 = -xy - \frac{1}{2}x^2 + 5$$



$$0 = -yx - \frac{1}{2}y^2 + 5$$

Therefore $x^2 = y^2$ and so $x = y$ (both quantities are positive).

Therefore $\frac{3}{2}x^2 = 5$, and so $x = \sqrt{\frac{10}{3}} = y$. By EVT max exists \Rightarrow is

achieved for $x = y = \sqrt{\frac{10}{3}}$.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let dimensions of box be x, y, z . We seek to maximize $V = xyz$.

Restrictions: $xy + 2(zx + zy) = 10 \Rightarrow z = \frac{10-xy}{2(x+y)} \Rightarrow V = xy \frac{10-xy}{2(x+y)}$.

$(x, y) \in \mathcal{R} = \{(x, y) | xy \leq 10, x \geq 0, y \geq 0\}$. We're solving:

$$0 = V_x = \frac{-xy^3 - \frac{1}{2}x^2y^2 + 5y^2}{(x+y)^2}$$

$$0 = V_y = \frac{-yx^3 - \frac{1}{2}x^2y^2 + 5x^2}{(x+y)^2}$$

We can assume $y \neq 0, x \neq 0$ (else the volume is zero). Then

$$0 = -xy - \frac{1}{2}x^2 + 5$$

$$0 = -yx - \frac{1}{2}y^2 + 5$$

Therefore $x^2 = y^2$ and so $x = y$ (both quantities are positive).

Therefore $\frac{3}{2}x^2 = 5$, and so $x = \sqrt{\frac{10}{3}} = y$. By EVT max exists \Rightarrow is

achieved for $x = y = \sqrt{\frac{10}{3}}$. Max volume $V_{max} = \frac{5}{9}\sqrt{30}$.

Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Proof.

Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a curve in S . Compute:



Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Proof.

Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a curve in S . Compute:

$$F(x(t), y(t), z(t)) = 0 \quad \text{apply } \frac{d}{dt}$$



Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Proof.

Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a curve in S . Compute:

$$\begin{aligned} F(x(t), y(t), z(t)) &= 0 && \text{apply } \frac{d}{dt} \\ \left(F_x \frac{dx}{dt} + F_y \frac{dy}{dt} + F_z \frac{dz}{dt} \right)_{|x=x(t), y=y(t), z=z(t)} &= 0 \end{aligned}$$



Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Proof.

Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a curve in S . Compute:

$$\begin{aligned}
 F(x(t), y(t), z(t)) &= 0 && \text{apply } \frac{d}{dt} \\
 \left(F_x \frac{dx}{dt} + F_y \frac{dy}{dt} + F_z \frac{dz}{dt} \right)_{|x=x(t), y=y(t), z=z(t)} &= 0 \\
 \mathbf{r}'(t) \cdot (F_x, F_y, F_z)_{|x=x(t), y=y(t), z=z(t)} &= 0
 \end{aligned}$$



Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Proof.

Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a curve in S . Compute:

$$F(x(t), y(t), z(t)) = 0 \quad \text{apply } \frac{d}{dt}$$

$$\left(F_x \frac{dx}{dt} + F_y \frac{dy}{dt} + F_z \frac{dz}{dt} \right)_{|x=x(t), y=y(t), z=z(t)} = 0$$

$$\mathbf{r}'(t) \cdot (F_x, F_y, F_z)_{|x=x(t), y=y(t), z=z(t)} = 0$$

$$\mathbf{r}'(t) \cdot (\nabla F)_{|x=x(t), y=y(t), z=z(t)} = 0$$



Let S be a surface that has implicit equation $F(x, y, z) = 0$ for some differentiable function F . Let P be a point on the surface.

Theorem

Suppose $\nabla F(P) \neq \mathbf{0}$. Then $\nabla F(P)$ is perpendicular to the tangent vector at P of every differentiable curve lying in S passing through P .

Proof.

Suppose $\mathbf{r}(t) = (x(t), y(t), z(t))$ is a curve in S . Compute:

$$\begin{aligned}
 F(x(t), y(t), z(t)) &= 0 && \text{apply } \frac{d}{dt} \\
 \left(F_x \frac{dx}{dt} + F_y \frac{dy}{dt} + F_z \frac{dz}{dt} \right)_{|x=x(t), y=y(t), z=z(t)} &= 0 \\
 \mathbf{r}'(t) \cdot (F_x, F_y, F_z)_{|x=x(t), y=y(t), z=z(t)} &= 0 \\
 \mathbf{r}'(t) \cdot (\nabla F)_{|x=x(t), y=y(t), z=z(t)} &= 0
 \end{aligned}$$



Definition (Tangent plane to level surface)

Suppose $\nabla F(P) \neq \mathbf{0}$. We define the tangent plane to the surface S at P to be the plane passing through P with normal vector $\nabla F(P)$.

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

$$\left. \frac{d}{dt} \right|_{t=0} (G(\mathbf{r}(t))) = 0$$

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

$$\begin{aligned} \frac{d}{dt}\bigg|_{t=0} (G(\mathbf{r}(t))) &= 0 \\ \left(\frac{\partial G}{\partial x} \frac{dx}{dt} + \frac{\partial G}{\partial y} \frac{dy}{dt} + \frac{\partial G}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} &= 0 \end{aligned}$$

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

$$\begin{aligned} \frac{d}{dt}\bigg|_{t=0} (G(\mathbf{r}(t))) &= 0 \\ \left(\frac{\partial G}{\partial x} \frac{dx}{dt} + \frac{\partial G}{\partial y} \frac{dy}{dt} + \frac{\partial G}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} &= 0 \\ \nabla G \cdot \mathbf{r}'(0) &= 0 \end{aligned}$$

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

$$\begin{aligned} \frac{d}{dt}\bigg|_{t=0} (G(\mathbf{r}(t))) &= 0 \\ \left(\frac{\partial G}{\partial x} \frac{dx}{dt} + \frac{\partial G}{\partial y} \frac{dy}{dt} + \frac{\partial G}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} &= 0 \\ \nabla G \cdot \mathbf{r}'(0) &= 0 \end{aligned}$$

- Therefore ∇G is \perp to tangent at P of every curve in S through P .

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

$$\begin{aligned} \frac{d}{dt}\bigg|_{t=0} (G(\mathbf{r}(t))) &= 0 \\ \left(\frac{\partial G}{\partial x} \frac{dx}{dt} + \frac{\partial G}{\partial y} \frac{dy}{dt} + \frac{\partial G}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} &= 0 \\ \nabla G \cdot \mathbf{r}'(0) &= 0 \end{aligned}$$

- Therefore ∇G is \perp to tangent at P of every curve in S through P .
- Therefore $\nabla F(P)$ and $\nabla G(P)$ are parallel,

Method of Lagrange Multipliers

Problem

Find the maximum of a function $G(x, y, z)$ subject to the variable restriction $F(x, y, z) = 0$.

- Let $S = \{(x, y, z) | F(x, y, z) = 0\}$.
- Suppose the max is achieved at $P(x_0, y_0, z_0)$. Let $\mathbf{r}(t) = (x(t), y(t), z(t))$ be a smooth curve on S such that $\mathbf{r}(0) = P$.
- Then $G(\mathbf{r}(t)) = G(x(t), y(t), z(t))$ has maximum at $t = 0$.

$$\begin{aligned} \frac{d}{dt}\bigg|_{t=0} (G(\mathbf{r}(t))) &= 0 \\ \left(\frac{\partial G}{\partial x} \frac{dx}{dt} + \frac{\partial G}{\partial y} \frac{dy}{dt} + \frac{\partial G}{\partial z} \frac{dz}{dt} \right)\bigg|_{t=0} &= 0 \\ \nabla G \cdot \mathbf{r}'(0) &= 0 \end{aligned}$$

- Therefore ∇G is \perp to tangent at P of every curve in S through P .
- Therefore $\nabla F(P)$ and $\nabla G(P)$ are parallel, i.e., there exists λ s.t.: $(\nabla G)(P) = \lambda(\nabla F)(P)$.

Example

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Example

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Region:

Example

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Region: closed square of vertices $(2, 0)$, $(0, 2)$, $(-2, 0)$, and $(0, -2)$.

Example

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Region: closed square of vertices $(2, 0)$, $(0, 2)$, $(-2, 0)$, and $(0, -2)$.

The function is continuous and the domain is bounded and closed.

Extreme Value Theorem $\implies f$ has global minimum and maximum points.

Example

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Region: closed square of vertices $(2, 0)$, $(0, 2)$, $(-2, 0)$, and $(0, -2)$.

The function is continuous and the domain is bounded and closed. Extreme Value Theorem $\implies f$ has global minimum and maximum points.

Strategy:

- Find critical points in the interior of the disk;
- Find extreme points on the boundary of the disk;
- Compare the values.

Example

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Region: closed square of vertices $(2, 0)$, $(0, 2)$, $(-2, 0)$, and $(0, -2)$.

The function is continuous and the domain is bounded and closed. Extreme Value Theorem $\implies f$ has global minimum and maximum points.

Strategy:

- Find critical points in the interior of the disk;
- Find extreme points on the boundary of the disk;
- Compare the values.

Since f is differentiable everywhere, the interior extreme points are among the solutions of the system

$$\begin{cases} f_x(x, y) = 0 \\ f_y(x, y) = 0 \end{cases} \iff \begin{cases} y = 0 \\ x = 0 \end{cases}$$

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Extreme points on the boundary: check each of the four sides.

For the segment joining $(2, 0)$ with $(0, 2)$ we get:

Find min/max of $f(x, y) = xy$

Subject to $g(x, y) = x + y - 2 = 0$

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Extreme points on the boundary: check each of the four sides.

For the segment joining $(2, 0)$ with $(0, 2)$ we get:

Find min/max of $f(x, y) = xy$

Subject to $g(x, y) = x + y - 2 = 0$

The Lagrange function is

$$F(x, y, \lambda) = f(x, y) - \lambda g(x, y) = xy - \lambda(x + y - 2)$$

Find the maximum and the minimum values of $f(x, y) = xy$ on the region $D = \{(x, y) \mid |x| + |y| \leq 2\}$.

Extreme points on the boundary: check each of the four sides.

For the segment joining $(2, 0)$ with $(0, 2)$ we get:

Find min/max of $f(x, y) = xy$

Subject to $g(x, y) = x + y - 2 = 0$

The Lagrange function is

$$F(x, y, \lambda) = f(x, y) - \lambda g(x, y) = xy - \lambda(x + y - 2)$$

The critical points of F are the solutions of the system

$$\begin{cases} F_x(x, y, \lambda) = 0 \\ F_y(x, y, \lambda) = 0 \\ F_\lambda(x, y, \lambda) = 0 \end{cases} \iff \begin{cases} y - \lambda = 0 \\ x - \lambda = 0 \\ x + y - 2 = 0 \end{cases} \iff \begin{cases} x = 1 \\ y = 1 \\ \lambda = 1 \end{cases}$$

Gradient Analysis

Gradient Analysis

- $(\nabla f)_{(1,1)} = \langle y, x \rangle|_{x=1,y=1} = \langle 1, 1 \rangle.$

Gradient Analysis

- $(\nabla f)_{(1,1)} = \langle y, x \rangle|_{x=1,y=1} = \langle 1, 1 \rangle$.
- If we move along the direction of the gradient at $(1, 1)$:
 - the value of the objective would increase;
 - the level curves of f we cross no longer intersect the constraint
 - those levels of f are unattainable on the constraint set $x + y = 2$.

Gradient Analysis

- $(\nabla f)_{(1,1)} = \langle y, x \rangle|_{x=1,y=1} = \langle 1, 1 \rangle$.
- If we move along the direction of the gradient at $(1, 1)$:
 - the value of the objective would increase;
 - the level curves of f we cross no longer intersect the constraint
 - those levels of f are unattainable on the constraint set $x + y = 2$.
- The point $(1, 1)$ corresponds to a local maxim.

Gradient Analysis

- $(\nabla f)_{(1,1)} = \langle y, x \rangle|_{x=1,y=1} = \langle 1, 1 \rangle$.
- If we move along the direction of the gradient at $(1, 1)$:
 - the value of the objective would increase;
 - the level curves of f we cross no longer intersect the constraint
 - those levels of f are unattainable on the constraint set $x + y = 2$.
- The point $(1, 1)$ corresponds to a local maxim.

Three more critical points on the boundary: $(-1, 1)$, $(-1, -1)$, $(1, -1)$.


Gradient Analysis

- $(\nabla f)_{(1,1)} = \langle y, x \rangle|_{x=1,y=1} = \langle 1, 1 \rangle$.
- If we move along the direction of the gradient at $(1, 1)$:
 - the value of the objective would increase;
 - the level curves of f we cross no longer intersect the constraint
 - those levels of f are unattainable on the constraint set $x + y = 2$.
- The point $(1, 1)$ corresponds to a local maxim.

Three more critical points on the boundary: $(-1, 1)$, $(-1, -1)$, $(1, -1)$.
Compare the values at all points:


- the global maximum is 1, attained at $(1, 1)$ and $(-1, -1)$;
- the global minimum is -1, attained at $(1, -1)$ and $(-1, 1)$;
- the critical point $(0, 0)$ is a saddle point.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ under the restriction

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ **under the restriction**

?

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ **under the restriction**

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$

By the Lagrange multiplier method, we need to solve the system

$$\nabla V = \lambda \nabla g$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$

By the Lagrange multiplier method, we need to solve the system

$$\nabla V = \lambda \nabla g$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$

By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$

By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g = \lambda ?$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$

By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g = \lambda (2z + y, 2z + x, 2y + 2x) \quad .$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$


By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g = \lambda(2z + y, 2z + x, 2y + 2x) \quad .$$

In other words we are solving the following system.

$$\begin{cases} yz &= \lambda(2z + y) \\ xz &= \lambda(2z + x) \\ xy &= \lambda(2x + 2y) \\ 10 &= xy + 2(zx + yz) \end{cases}$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$


By the Lagrange multiplier method, we need to solve the system

$$(yz, \textcolor{red}{xz}, xy) = \nabla V = \lambda \nabla g = \textcolor{red}{\lambda}(2z + y, \textcolor{red}{2z} + \textcolor{red}{x}, 2y + 2x) \quad .$$

In other words we are solving the following system.

$$\left| \begin{array}{lcl} yz & = & \lambda(2z + y) \\ \textcolor{red}{xz} & = & \textcolor{red}{\lambda}(2\textcolor{red}{z} + \textcolor{red}{x}) \\ xy & = & \lambda(2x + 2y) \\ 10 & = & xy + 2(zx + yz) \end{array} \right.$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$


By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g = \lambda (2z + y, 2z + x, 2y + 2x) \quad .$$

In other words we are solving the following system.

$$\begin{cases} yz &= \lambda(2z + y) \\ xz &= \lambda(2z + x) \\ xy &= \lambda(2x + 2y) \\ 10 &= xy + 2(zx + yz) \end{cases}$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$


By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g = \lambda(2z + y, 2z + x, 2y + 2x) \quad .$$

In other words we are solving the following system.

$$\begin{array}{lcl} yz & = & \lambda(2z + y) \\ xz & = & \lambda(2z + x) \\ xy & = & \lambda(2x + 2y) \\ 10 & = & xy + 2(zx + yz) \end{array}$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

We seek to maximize $V = xyz$ under the restriction

$$g(x, y, z) = xy + 2(zx + yz) - 10 = 0 \quad .$$


By the Lagrange multiplier method, we need to solve the system

$$(yz, zx, xy) = \nabla V = \lambda \nabla g = \lambda(2z + y, 2z + x, 2y + 2x) \quad .$$

In other words we are solving the following system.

$$\begin{aligned} yz &= \lambda(2z + y) \\ xz &= \lambda(2z + x) \\ xy &= \lambda(2x + 2y) \\ 10 &= xy + 2(zx + yz) \end{aligned}$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


$$yz = \lambda(2z + y)$$

$$xz = \lambda(2z + x)$$

$$xy = \lambda(2x + 2y)$$

$$10 = xy + 2(zx + yz)$$

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\begin{array}{l|l}
 \begin{array}{l}
 yz = \lambda(2z + y) \\
 xz = \lambda(2z + x) \\
 xy = \lambda(2x + 2y) \\
 10 = xy + 2(zx + yz)
 \end{array}
 &
 \Rightarrow
 \begin{array}{l}
 xyz = \lambda(2z + y)x \\
 xyz = \lambda(2z + x)y \\
 xy = \lambda(2x + 2y) \\
 10 = xy + 2(zx + yz)
 \end{array}
 \end{array}$$

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\begin{array}{l|l}
 \begin{array}{l}
 yz = \lambda(2z + y) \\
 xz = \lambda(2z + x) \\
 xy = \lambda(2x + 2y) \\
 10 = xy + 2(zx + yz)
 \end{array}
 &
 \Rightarrow
 \begin{array}{l}
 xyz = \lambda(2z + y)x \\
 xyz = \lambda(2z + x)y \\
 xy = \lambda(2x + 2y) \\
 10 = xy + 2(zx + yz)
 \end{array}
 \end{array}$$

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

From the first two equalities we get $2\lambda xz + \lambda xy = 2\lambda yz + \lambda xy$

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\begin{array}{l|l}
 yz = \lambda(2z + y) & xyz = \lambda(2z + y)x \\
 xz = \lambda(2z + x) & xyz = \lambda(2z + x)y \\
 xy = \lambda(2x + 2y) & xy = \lambda(2x + 2y) \\
 10 = xy + 2(zx + yz) & 10 = xy + 2(zx + yz)
 \end{array} \Rightarrow$$

From the first two equalities we get $2\lambda xz + \lambda xy = 2\lambda yz + \lambda xy$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

From the first two equalities we get $2\lambda xz + \lambda xy = 2\lambda yz + \lambda xy$ and so

$$\lambda xz = \lambda yz .$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$


From the first two equalities we get $2\lambda xz + \lambda xy = 2\lambda yz + \lambda xy$ and so

$$\lambda xz = \lambda yz .$$

We have that $\lambda \neq 0, z \neq 0$ (else the volume would be zero). Therefore

$$x = y$$

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\begin{array}{l|l}
 yz = \lambda(2z + y) & xyz = \lambda(2z + y)x \\
 xz = \lambda(2z + x) & xyz = \lambda(2z + x)y \\
 xy = \lambda(2x + 2y) & \textcolor{red}{xy} = \textcolor{red}{\lambda(2x + 2y)} \\
 10 = xy + 2(zx + yz) & 10 = xy + 2(zx + yz)
 \end{array} \Rightarrow$$

$$x = y$$

We substitute $y = x$ in the third equality to get $\textcolor{red}{x^2} = 4\lambda x$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y$$

We substitute $y = x$ in the third equality to get $x^2 = 4\lambda x$ and since $x \neq 0$ we get $\lambda = \frac{x}{4}$.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y$$

We substitute $y = x$ in the third equality to get $x^2 = 4\lambda x$ and since $x \neq 0$ we get $\lambda = \frac{x}{4}$. We substitute $\lambda = \frac{x}{4}$ in the original second equality to get $zx = \frac{x}{4}(2z + x)$.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y$$

We substitute $y = x$ in the third equality to get $x^2 = 4\lambda x$ and since $x \neq 0$ we get $\lambda = \frac{x}{4}$. We substitute $\lambda = \frac{x}{4}$ in the original second equality to get $zx = \frac{x}{4}(2z + x)$. Since $x \neq 0$ it follows that

$$z = \frac{1}{4}(2z + x)$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y$$

We substitute $y = x$ in the third equality to get $x^2 = 4\lambda x$ and since $x \neq 0$ we get $\lambda = \frac{x}{4}$. We substitute $\lambda = \frac{x}{4}$ in the original second equality to get $zx = \frac{x}{4}(2z + x)$. Since $x \neq 0$ it follows that $z = \frac{1}{4}(2z + x)$ and so $z = \frac{x}{2}$.

Example




Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y \qquad z = \frac{x}{2} .$$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\begin{array}{l|l}
 yz = \lambda(2z + y) & \\
 xz = \lambda(2z + x) & \\
 xy = \lambda(2x + 2y) & \\
 10 = xy + 2(zx + yz) &
 \end{array}
 \Rightarrow
 \begin{array}{l|l}
 xyz = \lambda(2z + y)x & \\
 xyz = \lambda(2z + x)y & \\
 xy = \lambda(2x + 2y) & \\
 10 = xy + 2(zx + yz) &
 \end{array}
 .$$

$$x = y \qquad z = \frac{x}{2} .$$

Finally we substitute $y = x, z = \frac{x}{2}$ in the last equality

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y \qquad z = \frac{x}{2} .$$

Finally we substitute $y = x, z = \frac{x}{2}$ in the last equality to get $10 = 3x^2$.

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .


$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y \quad z = \frac{x}{2} .$$

Finally we substitute $y = x, z = \frac{x}{2}$ in the last equality to get $10 = 3x^2$.

Thus $x = \frac{\sqrt{30}}{3}$

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.


Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y \quad z = \frac{x}{2} .$$

Finally we substitute $y = x, z = \frac{x}{2}$ in the last equality to get $10 = 3x^2$.
Thus $x = \frac{\sqrt{30}}{3}$ and therefore $y = \frac{\sqrt{30}}{3}, z = \frac{\sqrt{30}}{6}$,

Example



Find the maximal volume of a box with no lid whose surface area is $10m^2$.

Let the three dimensions of the box be x, y, z .

$$\left| \begin{array}{l} yz = \lambda(2z + y) \\ xz = \lambda(2z + x) \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. \Rightarrow \left| \begin{array}{l} xyz = \lambda(2z + y)x \\ xyz = \lambda(2z + x)y \\ xy = \lambda(2x + 2y) \\ 10 = xy + 2(zx + yz) \end{array} \right. .$$

$$x = y \qquad z = \frac{x}{2} .$$

Finally we substitute $y = x, z = \frac{x}{2}$ in the last equality to get $10 = 3x^2$.

Thus $x = \frac{\sqrt{30}}{3}$ and therefore $y = \frac{\sqrt{30}}{3}, z = \frac{\sqrt{30}}{6}$, our final answer.

Multiple Constraints

Find $\min / \max f(x, y, z)$

Subject to $g(x, y, z) = 0$

$h(x, y, z) = 0$

Multiple Constraints

Find $\min / \max f(x, y, z)$

Subject to $g(x, y, z) = 0$

$h(x, y, z) = 0$

Each constraint defines a surface \implies their intersection defines a curve.

Multiple Constraints

Find $\min / \max f(x, y, z)$

Subject to $g(x, y, z) = 0$

$h(x, y, z) = 0$

Each constraint defines a surface \implies their intersection defines a curve.

Condition: $(\nabla g)_P (\nabla h)_P$ are non-collinear for each intersection point P

The level surface of f through a point of extreme P_0 is tangent to the constraint curve, so $(\nabla f)(P_0)$ is perpendicular to the curve at P_0 .

Constraint curve included in both surfaces \implies

$(\nabla g)(P_0)$ and $(\nabla h)(P_0)$ are perpendicular to the curve \implies

Multiple Constraints

Find $\min / \max f(x, y, z)$

Subject to $g(x, y, z) = 0$

$h(x, y, z) = 0$

Each constraint defines a surface \implies their intersection defines a curve.

Condition: $(\nabla g)_P(\nabla h)_P$ are non-collinear for each intersection point P

The level surface of f through a point of extreme P_0 is tangent to the constraint curve, so $(\nabla f)(P_0)$ is perpendicular to the curve at P_0 .

Constraint curve included in both surfaces \implies

$(\nabla g)(P_0)$ and $(\nabla h)(P_0)$ are perpendicular to the curve \implies

there exist constants λ and μ such that

$$(\nabla f)(P_0) = \lambda(\nabla g)(P_0) + \mu(\nabla h)(P_0) .$$

Multiple Constraints

Find $\min / \max f(x, y, z)$

Subject to $g(x, y, z) = 0$

$h(x, y, z) = 0$

Each constraint defines a surface \implies their intersection defines a curve.

Condition: $(\nabla g)_P (\nabla h)_P$ are non-collinear for each intersection point P

The level surface of f through a point of extreme P_0 is tangent to the constraint curve, so $(\nabla f)(P_0)$ is perpendicular to the curve at P_0 .

Constraint curve included in both surfaces \implies

$(\nabla g)(P_0)$ and $(\nabla h)(P_0)$ are perpendicular to the curve \implies

there exist constants λ and μ such that

$$(\nabla f)(P_0) = \lambda(\nabla g)(P_0) + \mu(\nabla h)(P_0) .$$

The Lagrange function is in this case

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function:

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints:

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

$$F(x, y, z, \lambda, \mu) = x + 2y - \lambda(y^2 + z^2 - 5) - \mu(x + y + z - 1) .$$

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

$$F(x, y, z, \lambda, \mu) = x + 2y - \lambda(y^2 + z^2 - 5) - \mu(x + y + z - 1).$$

- Critical points of F :

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

$$F(x, y, z, \lambda, \mu) = x + 2y - \lambda(y^2 + z^2 - 5) - \mu(x + y + z - 1).$$

- Critical points of F : $(1, \sqrt{5/2}, -\sqrt{5/2})$ and $(1, -\sqrt{5/2}, \sqrt{5/2})$

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

$$F(x, y, z, \lambda, \mu) = x + 2y - \lambda(y^2 + z^2 - 5) - \mu(x + y + z - 1).$$

- Critical points of F : $(1, \sqrt{5/2}, -\sqrt{5/2})$ and $(1, -\sqrt{5/2}, \sqrt{5/2})$
- Values of objective function at these points:

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

$$F(x, y, z, \lambda, \mu) = x + 2y - \lambda(y^2 + z^2 - 5) - \mu(x + y + z - 1).$$

- Critical points of F : $(1, \sqrt{5/2}, -\sqrt{5/2})$ and $(1, -\sqrt{5/2}, \sqrt{5/2})$
- Values of objective function at these points:

$$f(1, \sqrt{5/2}, -\sqrt{5/2}) = 1 + 2\sqrt{5/2}, \quad f(1, -\sqrt{5/2}, \sqrt{5/2}) = 1 - 2\sqrt{5/2}$$

Example

Find the extreme points of $x + 2y$ on the intersection of the cylinder $y^2 + z^2 = 5$ and the plane $x + y + z = 1$.

- Objective function: $f(x, y, z) = x + 2y$.
- Constraints: $g(x, y, z) = y^2 + z^2 - 5$ and $h(x, y, z) = x + y + z - 1$.
- Lagrange function:

$$F(x, y, z, \lambda, \mu) = x + 2y - \lambda(y^2 + z^2 - 5) - \mu(x + y + z - 1).$$

- Critical points of F : $(1, \sqrt{5/2}, -\sqrt{5/2})$ and $(1, -\sqrt{5/2}, \sqrt{5/2})$
- Values of objective function at these points:

$$f(1, \sqrt{5/2}, -\sqrt{5/2}) = 1 + 2\sqrt{5/2}, \quad f(1, -\sqrt{5/2}, \sqrt{5/2}) = 1 - 2\sqrt{5/2}$$

- Constraint set is bounded and closed, function f is continuous $\implies f$ attains its extreme on the constraint \implies
 $(1, -\sqrt{5/2}, \sqrt{5/2})$ corresponds to an absolute minimum and
 $(1, \sqrt{5/2}, -\sqrt{5/2})$ corresponds to an absolute maximum.
- The minimum value is $f(1, -\sqrt{5/2}, \sqrt{5/2}) = 1 - 2\sqrt{5/2}$ and the
 maximal value is $f(1, \sqrt{5/2}, -\sqrt{5/2}) = 1 + 2\sqrt{5/2}$