

# Calculus III

## Lecture 15

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

1

## Parallelotopes

# Outline

1 Parallelotopes

2 Variable Changes in Multivariable Integrals

# License to use and redistribute


These lecture slides and their L<sup>A</sup>T<sub>E</sub>X source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,


as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.


- Let  $\bullet$  be a marked point. If omitted, we assume  $\bullet$  is the origin.




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .



- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .




- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region **spanned by** the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .



- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )


We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.



- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	?

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2		

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2	2	?

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2	2	parallelogram

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2	2	parallelogram
2	3	?

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2	2	parallelogram
2	3	parallelogram in space

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .

### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.

$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2	2	parallelogram
2	3	parallelogram in space
3	3	?

- Let  $\mathbf{o}$  be a marked point. If omitted, we assume  $\mathbf{o}$  is the origin.
- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_k = (v_{k1}, \dots, v_{kn})$  be  $k$  vectors in  $n$ -dimensional space,  $k \leq n$ .
- Let  $\mathcal{R}$  be region spanned by the vectors at  $\mathbf{o}$ , coefficients in  $[0, 1]$ .
- $\mathcal{R} = \{\mathbf{o} + t_1\mathbf{v}_1 + t_2\mathbf{v}_2 + \dots + t_k\mathbf{v}_k | t_1 \in [0, 1], \dots, t_k \in [0, 1]\}$ .


### Definition (parallelotope at $\mathbf{o}$ )

We call a region  $\mathcal{R}$  of the above form a  $k$ -dimensional parallelotope at the point  $\mathbf{o}$  in  $n$ -dimensional space.




- When  $k, n, \mathbf{o}$  are clear from context we can omit them.


$k$	$n$	parallelotope name
1	any	segment (in $n$ -dim space)
2	2	parallelogram
2	3	parallelogram in space
3	3	parallelepiped




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .



- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .



- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .

## Definition ( $k$ -volume of a parallelotope)


Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ .



- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .

### Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .



- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .


### Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .

- Let the height vector  $\mathbf{h}_k$  be the vector of the form  

$$\mathbf{h}_k = \mathbf{v}_k + a_1\mathbf{v}_1 + \dots + a_{k-1}\mathbf{v}_{k-1}$$
 for which  

$$\mathbf{h}_k \cdot \mathbf{v}_1 = 0, \dots, \mathbf{h}_k \cdot \mathbf{v}_{k-1} = 0.$$




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .

### Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .

- Let the height vector  $\mathbf{h}_k$  be the vector of the form  $\mathbf{h}_k = \mathbf{v}_k + a_1\mathbf{v}_1 + \dots + a_{k-1}\mathbf{v}_{k-1}$  for which  $\mathbf{h}_k \cdot \mathbf{v}_1 = 0, \dots, \mathbf{h}_k \cdot \mathbf{v}_{k-1} = 0$ .
- Then  $h_k$  is computed as the length of  $\mathbf{h}_k$ .




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .

### Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .

- Let the height vector  $\mathbf{h}_k$  be the vector of the form  $\mathbf{h}_k = \mathbf{v}_k + a_1\mathbf{v}_1 + \dots + a_{k-1}\mathbf{v}_{k-1}$  for which  $\mathbf{h}_k \cdot \mathbf{v}_1 = 0, \dots, \mathbf{h}_k \cdot \mathbf{v}_{k-1} = 0$ .
- Then  $h_k$  is computed as the length of  $\mathbf{h}_k$ .
- For the largest parallelotope  $\mathcal{R}_n$ , we already have definition of volume:




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .

### Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .

- Let the height vector  $\mathbf{h}_k$  be the vector of the form  $\mathbf{h}_k = \mathbf{v}_k + a_1\mathbf{v}_1 + \dots + a_{k-1}\mathbf{v}_{k-1}$  for which  $\mathbf{h}_k \cdot \mathbf{v}_1 = 0, \dots, \mathbf{h}_k \cdot \mathbf{v}_{k-1} = 0$ .
- Then  $h_k$  is computed as the length of  $\mathbf{h}_k$ .
- For the largest parallelotope  $\mathcal{R}_n$ , we already have definition of volume: the integral of 1 over  $\mathcal{R}_n$ .




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- $\mathcal{R}_k$  can be regarded as “prism” with base  $\mathcal{R}_{k-1}$ .
- Let  $h_k$  be the height from  $\mathbf{v}_k$  to the base  $\mathcal{R}_{k-1}$ .

### Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .

- Let the height vector  $\mathbf{h}_k$  be the vector of the form  $\mathbf{h}_k = \mathbf{v}_k + a_1\mathbf{v}_1 + \dots + a_{k-1}\mathbf{v}_{k-1}$  for which  $\mathbf{h}_k \cdot \mathbf{v}_1 = 0, \dots, \mathbf{h}_k \cdot \mathbf{v}_{k-1} = 0$ .
- Then  $h_k$  is computed as the length of  $\mathbf{h}_k$ .
- For the largest parallelotope  $\mathcal{R}_n$ , we already have definition of volume: the integral of 1 over  $\mathcal{R}_n$ .
- We will see that  $\text{Vol}_n(\mathcal{R}_n)$  equals that integral.

# Length, Surface Area, Volume as $k$ -volumes




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Let  $h_k$  be the height of  $\mathcal{R}_k$  with base  $\mathcal{R}_{k-1}$ .

## Definition ( $k$ -volume of a parallelotope)

Define  $\text{Vol}_1(\mathcal{R}_1) = |\mathbf{v}_1|$ . For  $k > 1$ , define  $\text{Vol}_k(\mathcal{R}_k) = h_k \text{Vol}_{k-1}(\mathcal{R}_{k-1})$ .

	spanned by	$\text{Vol}_k(\mathcal{R}_k)$	volume name
$\mathcal{R}_1$	$\mathbf{v}_1$	$h_1 =  \mathbf{v}_1 $	length
$\mathcal{R}_2$	$\mathbf{v}_1, \mathbf{v}_2$	$h_1 h_2$	(surface) area
$\mathcal{R}_3$	$\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3$	$h_1 h_2 h_3$	volume
$\vdots$	$\vdots$	$\vdots$	$\vdots$
$\mathcal{R}_k$	$\mathbf{v}_1, \dots, \mathbf{v}_k$	$h_1 \dots h_k$	$k$ -volume

# Integral and Algebraic Volume Definitions Agree




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Let  $h_k$  be the height of  $\mathcal{R}_k$  with base  $\mathcal{R}_{k-1}$ .

## Theorem

$$\text{Vol}_n(\mathcal{R}_n) = h_n \text{Vol}_{n-1}(\mathcal{R}_{n-1}) = \int \cdots \int_{\mathcal{R}_n} 1 \cdot dx_1 \dots dx_n.$$

# Integral and Algebraic Volume Definitions Agree




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Let  $h_k$  be the height of  $\mathcal{R}_k$  with base  $\mathcal{R}_{k-1}$ .

## Theorem

$$\text{Vol}_n(\mathcal{R}_n) = h_n \text{Vol}_{n-1}(\mathcal{R}_{n-1}) = \int \cdots \int_{\mathcal{R}_n} 1 \cdot dx_1 \dots dx_n.$$

- Right hand side: approx. vol. with boxes, sides along coord. axes.

# Integral and Algebraic Volume Definitions Agree




- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Let  $h_k$  be the height of  $\mathcal{R}_k$  with base  $\mathcal{R}_{k-1}$ .

## Theorem

$$\text{Vol}_n(\mathcal{R}_n) = h_n \text{Vol}_{n-1}(\mathcal{R}_{n-1}) = \int \cdots \int_{\mathcal{R}_n} 1 \cdot dx_1 \dots dx_n.$$

- Right hand side: approx. vol. with boxes, sides along coord. axes.
- Left hand side: approximate volume with slabs parallel to base.**

# Integral and Algebraic Volume Definitions Agree



- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors in  $n$ -dimensional space.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Let  $h_k$  be the height of  $\mathcal{R}_k$  with base  $\mathcal{R}_{k-1}$ .

## Theorem

$$\text{Vol}_n(\mathcal{R}_n) = h_n \text{Vol}_{n-1}(\mathcal{R}_{n-1}) = \int \cdots \int_{\mathcal{R}_n} 1 \cdot dx_1 \dots dx_n.$$

- Right hand side: approx. vol. with boxes, sides along coord. axes.
- Left hand side: approximate volume with slabs parallel to base.
- Theorem is fully intuitive but its proof is surprisingly laborious.


- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .

- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = ?$

- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = v_{i1}v_{j1} + \dots + v_{in}v_{jn}$ .

- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = v_{i1}v_{j1} + \dots + v_{in}v_{jn}$ .


Theorem ( $k$ -volume = Gram determinant)



$$\text{Vol}_k(\mathcal{R}_k) = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \dots & \mathbf{v}_1 \cdot \mathbf{v}_k \\ \vdots & \dots & \vdots \\ \mathbf{v}_k \cdot \mathbf{v}_1 & \dots & \mathbf{v}_k \cdot \mathbf{v}_k \end{vmatrix}}.$$

- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = v_{i1}v_{j1} + \dots + v_{in}v_{jn}$ .


Theorem ( $k$ -volume = Gram determinant)



$$\text{Vol}_k(\mathcal{R}_k) = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \dots & \mathbf{v}_1 \cdot \mathbf{v}_k \\ \vdots & \dots & \vdots \\ \mathbf{v}_k \cdot \mathbf{v}_1 & \dots & \mathbf{v}_k \cdot \mathbf{v}_k \end{vmatrix}}.$$

- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = v_{i1}v_{j1} + \dots + v_{in}v_{jn}$ .

Theorem ( $k$ -volume = Gram determinant)




$$\text{Vol}_k(\mathcal{R}_k) = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \dots & \mathbf{v}_1 \cdot \mathbf{v}_k \\ \vdots & \dots & \vdots \\ \mathbf{v}_k \cdot \mathbf{v}_1 & \dots & \mathbf{v}_k \cdot \mathbf{v}_k \end{vmatrix}}.$$

Proof: studied in Linear algebra ( $\text{Vol}_k$  - defined by algebra only).

- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = v_{i1}v_{j1} + \dots + v_{in}v_{jn}$ .

Theorem ( $k$ -volume = Gram determinant)




$$\text{Vol}_k(\mathcal{R}_k) = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \dots & \mathbf{v}_1 \cdot \mathbf{v}_k \\ \vdots & \dots & \vdots \\ \mathbf{v}_k \cdot \mathbf{v}_1 & \dots & \mathbf{v}_k \cdot \mathbf{v}_k \end{vmatrix}}.$$

Proof: studied in Linear algebra ( $\text{Vol}_k$  - defined by algebra only).  $\text{Vol}_n(\mathcal{R}_n)$  is a perfect square for all  $n$ .


- Let  $\mathbf{v}_1 = (v_{11}, \dots, v_{1n}), \dots, \mathbf{v}_n = (v_{n1}, \dots, v_{nn})$  be  $n$ -vectors.
- Let  $\mathcal{R}_k$  be the parallelotope spanned by  $\mathbf{v}_1, \dots, \mathbf{v}_k$ .
- Recall that  $\mathbf{v}_i \cdot \mathbf{v}_j = v_{i1}v_{j1} + \dots + v_{in}v_{jn}$ .

Theorem ( $k$ -volume = Gram determinant)



$$\text{Vol}_k(\mathcal{R}_k) = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \dots & \mathbf{v}_1 \cdot \mathbf{v}_k \\ \vdots & \dots & \vdots \\ \mathbf{v}_k \cdot \mathbf{v}_1 & \dots & \mathbf{v}_k \cdot \mathbf{v}_k \end{vmatrix}}.$$

Proof: studied in Linear algebra ( $\text{Vol}_k$  - defined by algebra only).  $\text{Vol}_n(\mathcal{R}_n)$  is a perfect square for all  $n$ .



Theorem

$$\text{Vol}_n(\mathcal{R}_n) = \pm \begin{vmatrix} v_{11} & \dots & v_{n1} \\ \vdots & \dots & \vdots \\ v_{1n} & \dots & v_{nn} \end{vmatrix}.$$

# Properties of determinants

- Multiplying a column of a matrix by a number changes multiplies the determinant by the same number. In precise notation:

## Lemma

$$\begin{vmatrix} a_{11} & \dots & xa_{1k} & \dots & a_{1n} \\ a_{21} & \dots & xa_{2k} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & \dots & xa_{nk} & \dots & a_{nn} \end{vmatrix} = x \begin{vmatrix} a_{11} & \dots & a_{1k} & \dots & a_{1n} \\ a_{21} & \dots & a_{2k} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & \dots & a_{nk} & \dots & a_{nn} \end{vmatrix}$$

## Example

Find the 1-dimensional volume (length) of the segment through the origin spanned by  $\mathbf{v} = (1, 2, 3)$ .

## Example

Find the 1-dimensional volume (length) of the segment  $\mathcal{R}_1$  through the origin spanned by  $\mathbf{v} = (v_1, v_2, v_3)$ .

$$\text{Vol}_1 = \sqrt{\underbrace{\mathbf{v} \cdot \mathbf{v}}_{1 \times 1 \text{ Gram determinant}}} = \sqrt{v_1^2 + v_2^2 + v_3^2}.$$

## Example

Let  $\mathcal{R}_2$  be the parallelogram in 2-dimensional space spanned by  $\mathbf{v}_1 = (2, 3)$ ,  $\mathbf{v}_2 = (5, 7)$ . Find the area of  $\mathcal{R}_2$ .

## Example

Let  $\mathcal{R}_2$  be the parallelogram in 2-dimensional space spanned by  $\mathbf{v}_1 = (v_{11}, v_{12})$ ,  $\mathbf{v}_2 = (v_{21}, v_{22})$ . Find the area of  $\mathcal{R}_2$ .

$$\text{Vol}_2 = \pm \begin{vmatrix} v_{11} & v_{21} \\ v_{12} & v_{22} \end{vmatrix}$$
$$\text{Vol}_2 = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \mathbf{v}_1 \cdot \mathbf{v}_2 \\ \mathbf{v}_2 \cdot \mathbf{v}_1 & \mathbf{v}_2 \cdot \mathbf{v}_2 \end{vmatrix}}$$

## Example

Find the surface area of the parallelogram spanned by  $\mathbf{v}_1 = (1, 2, 3)$  and  $\mathbf{v}_2 = (5, 7, 11)$ .

$$\text{Vol}_2 = \sqrt{\begin{vmatrix} \mathbf{v}_1 \cdot \mathbf{v}_1 & \mathbf{v}_1 \cdot \mathbf{v}_2 \\ \mathbf{v}_2 \cdot \mathbf{v}_1 & \mathbf{v}_2 \cdot \mathbf{v}_2 \end{vmatrix}} = \sqrt{\begin{vmatrix} 14 & 52 \\ 52 & 195 \end{vmatrix}} = \sqrt{26}.$$

## Example

Find the surface area of the parallelogram spanned by  
 $\mathbf{v}_1 = (v_{11}, v_{12}, v_{13})$  and  $\mathbf{v}_2 = (v_{21}, v_{22}, v_{23})$ .

## Example

Find the volume of the parallelepiped with vertex at the origin and spanned by  $\mathbf{v}_1 = (1, 2, 3)$ ,  $\mathbf{v}_2 = (3, 5, 7)$ ,  $\mathbf{v}_3 = (5, 7, 11)$ .

$$\text{Vol}_3 = \left| \det \begin{pmatrix} 1 & 2 & 3 \\ 3 & 5 & 7 \\ 5 & 7 & 11 \end{pmatrix} \right| = |-2| = 2.$$

## Example

Find the volume of the parallelepiped spanned by  $\mathbf{v}_1 = (v_{11}, v_{12}, v_{13})$ ,  $\mathbf{v}_2 = (v_{21}, v_{22}, v_{23})$ ,  $\mathbf{v}_3 = (v_{31}, v_{32}, v_{33})$ .

- Recall the polar coordinate variable change

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta.\end{aligned}$$

- Recall the polar coordinate variable change

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta.\end{aligned}$$

- This variable change can be thought of as two functions:  
 $x = h(r, \theta) = r \cos \theta$  and  $y = g(r, \theta) = r \sin \theta.$

- Recall the polar coordinate variable change

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta.\end{aligned}$$

- This variable change can be thought of as two functions:  
 $x = h(r, \theta) = r \cos \theta$  and  $y = g(r, \theta) = r \sin \theta$ .
- The functions  $h, g$  map the two-dimensional plane with coordinates  $r, \theta$  into the two-dimensional plane with coordinates  $x, y$ .

- Recall the polar coordinate variable change

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta.\end{aligned}$$

- This variable change can be thought of as two functions:  
 $x = h(r, \theta) = r \cos \theta$  and  $y = g(r, \theta) = r \sin \theta$ .
- The functions  $h, g$  map the two-dimensional plane with coordinates  $r, \theta$  into the two-dimensional plane with coordinates  $x, y$ .
- Let  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an infinitely differentiable map.

- Recall the polar coordinate variable change

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta.\end{aligned}$$

- This variable change can be thought of as two functions:  
 $x = h(r, \theta) = r \cos \theta$  and  $y = g(r, \theta) = r \sin \theta$ .
- The functions  $h, g$  map the two-dimensional plane with coordinates  $r, \theta$  into the two-dimensional plane with coordinates  $x, y$ .
- Let  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an infinitely differentiable map.
- In other words,  $\mathbf{f}$  takes  $n$  scalar inputs and produces  $n$  scalar outputs.

- Recall the polar coordinate variable change

$$\begin{aligned}x &= r \cos \theta \\y &= r \sin \theta.\end{aligned}$$

- This variable change can be thought of as two functions:  
 $x = h(r, \theta) = r \cos \theta$  and  $y = g(r, \theta) = r \sin \theta$ .
- The functions  $h, g$  map the two-dimensional plane with coordinates  $r, \theta$  into the two-dimensional plane with coordinates  $x, y$ .
- Let  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be an infinitely differentiable map.
- In other words,  $\mathbf{f}$  takes  $n$  scalar inputs and produces  $n$  scalar outputs.

### Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ .

## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{vmatrix} & f_1(y_1, \dots, y_n) \\ & \vdots \\ & f_n(y_1, \dots, y_n) \end{vmatrix} .$$


## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & f_1(y_1, \dots, y_n) \\ \hline & \vdots \\ & f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c} \color{red}{x_1} = f_1(y_1, \dots, y_n) \\ \vdots \\ \color{red}{x_n} = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.
- Do similarly with respect to the remaining variables.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.
- Do similarly with respect to the remaining variables.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.
- Do similarly with respect to the remaining variables.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.

- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.
- Do similarly with respect to the remaining variables.




## Definition (Infinitely Smooth Variable Change)

An infinitely differentiable map  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is called an (infinitely) smooth variable change.


- Variable change  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is given by f-ns  $f_1, \dots, f_n$ . We write:

$$\mathbf{f} : \begin{array}{c|c} & x_1 = f_1(y_1, \dots, y_n) \\ & \vdots \\ & x_n = f_n(y_1, \dots, y_n) \end{array} .$$

- The variables  $y_1, \dots, y_n$  denote coordinates in the domain of  $\mathbf{f}$ .
- We may include vars.  $x_1, \dots, x_n$  denoting coords. in codomain of  $\mathbf{f}$ .
- Fix  $y_2, \dots, y_n$  and view  $\mathbf{f}$  as curve with respect to  $y_1$ ; plot.
- Do similarly with respect to the remaining variables.



$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$




## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix}$$

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$




## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$




## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$

- Consider curve given by  $\mathbf{f}$  with **parameter  $y_1$**  (other  $y_j$ 's fixed).

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$




## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$

- Consider **curve given by  $\mathbf{f}$**  with parameter  $y_1$  (other  $y_j$ 's fixed).

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$




## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$

- Consider curve given by  $\mathbf{f}$  with parameter  $y_1$  (other  $y_j$ 's fixed).
- Then the tangent vector of that curve is  $\left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right)$ .

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$




## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$

- Consider curve given by  $\mathbf{f}$  with parameter  $y_1$  (other  $y_j$ 's fixed).
- Then the tangent vector of that curve is  $\left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right)$ .
- Similar considerations hold for  $y_2$

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$



## Definition (Jacobian matrix)

The Jacobian matrix of a variable change  $\mathbf{f}$  is defined as the matrix


$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial f_1}{\partial y_1} & \dots & \frac{\partial f_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_n}{\partial y_1} & \dots & \frac{\partial f_n}{\partial y_n} \end{pmatrix} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$

- Consider curve given by  $\mathbf{f}$  with parameter  $y_1$  (other  $y_j$ 's fixed).
- Then the tangent vector of that curve is  $\left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right)$ .
- Similar considerations hold for  $y_2, \dots, y_n$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \cdots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{array}{l} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{array}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .
- The point  $\mathbf{y}$  and the corners  $\mathbf{y} + \Delta y_1 \mathbf{e}_1, \dots, \mathbf{y} + \Delta y_n \mathbf{e}_n$  suffice to identify  $B$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .
- The point  $\mathbf{y}$  and the corners  $\mathbf{y} + \Delta y_1 \mathbf{e}_1, \dots, \mathbf{y} + \Delta y_n \mathbf{e}_n$  suffice to identify  $B$ .
- $\text{Vol}(B) = \Delta y_1 \dots \Delta y_n$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .
- The point  $\mathbf{y}$  and the corners  $\mathbf{y} + \Delta y_1 \mathbf{e}_1, \dots, \mathbf{y} + \Delta y_n \mathbf{e}_n$  suffice to identify  $B$ .
- $\text{Vol}(B) = \Delta y_1 \dots \Delta y_n$ .
- Let the image of  $B$  be  $\mathbf{f}(B) = \mathcal{C}$ .  $\mathcal{C}$  is a “curvilinear box”.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .
- The point  $\mathbf{y}$  and the corners  $\mathbf{y} + \Delta y_1 \mathbf{e}_1, \dots, \mathbf{y} + \Delta y_n \mathbf{e}_n$  suffice to identify  $B$ .
- $\text{Vol}(B) = \Delta y_1 \dots \Delta y_n$ .
- Let the image of  $B$  be  $\mathbf{f}(B) = \mathcal{C}$ .  $\mathcal{C}$  is a “curvilinear box”.
- Let  $\mathcal{E}$  be the parallelopiped at  $\mathbf{f}(\mathbf{y})$  spanned by images of the corners of  $B$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .
- The point  $\mathbf{y}$  and the corners  $\mathbf{y} + \Delta y_1 \mathbf{e}_1, \dots, \mathbf{y} + \Delta y_n \mathbf{e}_n$  suffice to identify  $B$ .
- $\text{Vol}(B) = \Delta y_1 \dots \Delta y_n$ .
- Let the image of  $B$  be  $\mathbf{f}(B) = \mathbf{C}$ .  $\mathbf{C}$  is a “curvilinear box”.
- Let  $E$  be the parallelopiped at  $\mathbf{f}(\mathbf{y})$  spanned by images of the corners of  $B$ . Then  $\text{Vol}(\mathbf{C}) \approx \text{Vol}_n(E)$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $\mathbf{e}_1, \dots, \mathbf{e}_n$  be basis vectors. Fix point  $\mathbf{y} = (y_1, \dots, y_n)$ .
- Let  $\Delta y_1, \dots, \Delta y_n$  be small numbers. Construct small box  $B$  with corner  $\mathbf{y}$  spanned by the vectors  $\Delta y_1 \mathbf{e}_1, \dots, \Delta y_n \mathbf{e}_n$ .
- The point  $\mathbf{y}$  and the corners  $\mathbf{y} + \Delta y_1 \mathbf{e}_1, \dots, \mathbf{y} + \Delta y_n \mathbf{e}_n$  suffice to identify  $B$ .
- $\text{Vol}(B) = \Delta y_1 \dots \Delta y_n$ .
- Let the image of  $B$  be  $\mathbf{f}(B) = C$ .  $C$  is a “curvilinear box”.
- Let  $E$  be the parallelopiped at  $\mathbf{f}(\mathbf{y})$  spanned by images of the corners of  $B$ . Then  $\text{Vol}(C) \approx \text{Vol}_n(E)$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(C) \approx \text{Vol}_n(E)$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(C) \approx \text{Vol}_n(E)$ .
- The first edge of  $E$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(C) \approx \text{Vol}_n(E)$ .
- The first edge of  $E$  corresponds to the vector  $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y})$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(C) \approx \text{Vol}_n(E)$ .
- The first edge of  $E$  corresponds to the vector  $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y})$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector  $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y})$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector  $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y})$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector  $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y})$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{array}{l} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{array} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(C) \approx \text{Vol}_n(E)$ .
- The first edge of  $E$  corresponds to the vector  $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y})))$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector  
 $\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector

$$\begin{aligned} \mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) &\approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1} \\ &= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) \end{aligned}$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector


$$\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$$

$$= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right).$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector
 
$$\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$$

$$= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right).$$
- Similar considerations hold for the other edges of  $\mathcal{E}$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector
 
$$\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$$

$$= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right).$$
- Similar considerations holds for the other edges of  $\mathcal{E}$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$





- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector
 
$$\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$$

$$= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right).$$
- Similar considerations holds for the other edges of  $\mathcal{E}$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .

- The first edge of  $\mathcal{E}$  corresponds to the vector

$$\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$$

$$= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right).$$


- Similar considerations holds for the other edges of  $\mathcal{E}$ .
- Let  $\mathcal{J}$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors

$$\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right).$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E})$ .
- The first edge of  $\mathcal{E}$  corresponds to the vector
 
$$\mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) \approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1}$$

$$= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right).$$
- Similar considerations hold for the other edges of  $\mathcal{E}$ .
- Let  $\mathcal{J}$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors
 
$$\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right).$$
- Then  $\text{Vol}(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E}) \approx \text{Vol}_n(\mathcal{J})$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



- $\text{Vol}(C) \approx \text{Vol}_n(E)$ .

- The first edge of  $E$  corresponds to the vector


$$\begin{aligned} \mathbf{f}(\mathbf{y} + \Delta y_1 \mathbf{e}_1) - \mathbf{f}(\mathbf{y}) &\approx \Delta y_1 (D_{\mathbf{e}_1} (\mathbf{f}(\mathbf{y}))) = \Delta y_1 \frac{\partial \mathbf{f}}{\partial y_1} \\ &= \Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right) = \left( \Delta y_1 \frac{\partial x_1}{\partial y_1}, \dots, \Delta y_1 \frac{\partial x_n}{\partial y_1} \right). \end{aligned}$$

- Similar considerations holds for the other edges of  $E$ .
- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- Then  $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- $\text{Vol}_n(\mathcal{C}) \approx \text{Vol}_n(\mathcal{E}) \approx \text{Vol}_n(\mathcal{J})$

$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $J$  be the parallelopiped at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$

$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$

$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix} = \pm \Delta y_1 \dots \Delta y_n \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$

$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix} = \pm \Delta y_1 \dots \Delta y_n \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$


$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix} = \pm \Delta y_1 \dots \Delta y_n \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

$$= \pm \det(J_{\mathbf{f}}) \Delta y_1 \dots \Delta y_n$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ . Suppose  $\det J_{\mathbf{f}} \geq 0$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J)$


$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix} = \pm \Delta y_1 \dots \Delta y_n \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

$$= \pm \det(J_{\mathbf{f}}) \Delta y_1 \dots \Delta y_n$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



- Let  $J$  be the parallelotope at  $\mathbf{f}(\mathbf{y})$  spanned by the vectors  $\Delta y_1 \left( \frac{\partial x_1}{\partial y_1}, \dots, \frac{\partial x_n}{\partial y_1} \right), \dots, \Delta y_n \left( \frac{\partial x_1}{\partial y_n}, \dots, \frac{\partial x_n}{\partial y_n} \right)$ . Suppose  $\det J_{\mathbf{f}} \geq 0$ .
- $\text{Vol}(C) \approx \text{Vol}_n(E) \approx \text{Vol}_n(J) = \det J_{\mathbf{f}} \Delta y_1 \dots \Delta y_n$


$$\text{Vol}_n(J) = \pm \begin{vmatrix} \Delta y_1 \frac{\partial x_1}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \Delta y_1 \frac{\partial x_n}{\partial y_1} & \dots & \Delta y_n \frac{\partial x_n}{\partial y_n} \end{vmatrix} = \pm \Delta y_1 \dots \Delta y_n \begin{vmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{vmatrix}$$

$$= \pm \det(J_{\mathbf{f}}) \Delta y_1 \dots \Delta y_n$$

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




$\text{Vol}(C) \approx \det(J_f) \Delta y_1 \dots \Delta y_n$



# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(C(\mathbf{y})) \approx \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$




- Regard  $\mathbf{y}$  as a variable

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{vmatrix} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{vmatrix} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(C(\mathbf{y})) \approx \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$




- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




•  $\sum_{\mathbf{y}} \text{Vol}(\mathbf{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$\text{Vol}(\mathcal{S}) = \sum_{\mathbf{y}} \text{Vol}(C(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$




- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases}$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(\mathcal{S}) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$




- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$\text{Vol}(\mathcal{S}) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$


- $\int_S 1 \cdot dx_1 \dots dx_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(\mathcal{S}) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$


- $\int_{\mathcal{S}} 1 \cdot dx_1 \dots dx_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.
- Let  $\Delta y_1 \rightarrow 0, \dots, \Delta y_n \rightarrow 0$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(\mathcal{S}) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$


- $\int_{\mathcal{S}} 1 \cdot dx_1 \dots dx_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.
- Let  $\Delta y_1 \rightarrow 0, \dots, \Delta y_n \rightarrow 0$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(\mathcal{S}) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$


- $\int_{\mathcal{S}} 1 \cdot dx_1 \dots dx_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.
- Let  $\Delta y_1 \rightarrow 0, \dots, \Delta y_n \rightarrow 0$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \cdots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(S) = \sum_{\mathbf{y}} \text{Vol}(\mathbf{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$


- $\int_S 1 \cdot dx_1 \dots dx_n$
- $\int \dots \int_{\mathcal{R}} \det(J_{\mathbf{f}}(y)) dy_1 \dots dy_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.
- Let  $\Delta y_1 \rightarrow 0, \dots, \Delta y_n \rightarrow 0$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$$\text{Vol}(S) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$$


- $\int_S 1 \cdot dx_1 \dots dx_n = \int_{\mathcal{R}} \dots \int_{\mathcal{R}} \det(J_{\mathbf{f}}(y)) dy_1 \dots dy_n$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.
- Let  $\Delta y_1 \rightarrow 0, \dots, \Delta y_n \rightarrow 0$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \cdots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



$\bullet$   $\text{Vol}(S) = \sum_{\mathbf{y}} \text{Vol}(\mathcal{C}(\mathbf{y})) \approx \sum_{\mathbf{y}} \det(J_{\mathbf{f}}(\mathbf{y})) \Delta y_1 \dots \Delta y_n$


$$\int_S 1 \cdot dx_1 \dots dx_n = \int_{\mathcal{R}} \dots \int_{\mathcal{R}} \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n$$

- Regard  $\mathbf{y}$  as a variable and let it traverse a rectangular mesh.
- Sum over the rectangular mesh.
- Let  $\Delta y_1 \rightarrow 0, \dots, \Delta y_n \rightarrow 0$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \cdots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \cdots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$



## Theorem (Variable change in multivariable integrals)

Let  $\mathbf{f}$  be a smooth one to one variable change. Let  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ .

Then


$$\int \cdots \int_{\mathcal{S}} d\mathbf{x}_1 \dots d\mathbf{x}_n = \int \cdots \int_{\mathcal{R}} \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

provided that  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$  for all  $\mathbf{y} \in \mathcal{R}$ .

# Variable change in multivariable integrals

$$\mathbf{f} : \begin{cases} x_1 = f_1(y_1, \dots, y_n) \\ \vdots \\ x_n = f_n(y_1, \dots, y_n) \end{cases} .$$

$$J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x_1}{\partial y_1} & \dots & \frac{\partial x_1}{\partial y_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial y_1} & \dots & \frac{\partial x_n}{\partial y_n} \end{pmatrix}$$




## Theorem (Variable change in multivariable integrals)

Let  $\mathbf{f}$  be a smooth one to one variable change. Let  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ . Let  $h$  be an integrable function. Then

$$\int_S h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$


provided that  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$  for all  $\mathbf{y} \in \mathcal{R}$ .

## Example




Find the volume of a ball of radius  $r$ .

## Example




Find the volume of a spherical curvilinear box, given by the spherical coordinate inequalities

$$\begin{aligned}\rho_{min} \leq \rho &\leq \rho_{max}, \\ \phi_{min} \leq \phi &\leq \phi_{max}, \\ \theta_{min} \leq \theta &\leq \theta_{max}.\end{aligned}$$




## Example




Find the volume of the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ ,  
 $a, b, c > 0$ .

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn.


## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ .


## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .


## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .


## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane.

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ .

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles.

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$|OC| = ?$$

the indicated angles. We have .

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$|OC| = R$$

the indicated angles. We have

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$\begin{aligned} |OC| &= R \\ |\textcolor{red}{PC}| &= ? \end{aligned}$$

the indicated angles. We have .

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .


Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$|OC| = R$$

$$|PC| = r$$

the indicated angles. We have .

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$\begin{aligned} |OC| &= R \\ |PC| &= r \\ |\textcolor{red}{PH}| &= ? \end{aligned} .$$

the indicated angles. We have

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$\begin{aligned} |OC| &= R \\ |PC| &= r \\ |\textcolor{red}{PH}| &= \textcolor{red}{r \sin \theta} \end{aligned} .$$

the indicated angles. We have

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

$$\begin{aligned} |OC| &= R \\ |PC| &= r \\ |PH| &= r \sin \theta \\ |OH| &= ? \end{aligned} .$$

the indicated angles. We have

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = ? \\ y = ? \\ z = ? \end{cases}$$


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = \\ z = \end{cases}$$


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = ? \\ z = \end{cases}$$


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = \end{cases}$$


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = ? \end{cases}$$


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$


Suppose the toroid sits in space as drawn. Let  $P(x, y, z) \in S$ . Let  $\mathcal{P}$  be the plane through the  $z$ -axis and  $P$ .

Let  $H$  be the heel of the perpendicular from  $P$  to the  $x, y$ -plane. Let  $C$  be the center of the circle cross-section of  $\mathcal{P}$  with  $T$ . Let  $\phi$  and  $\theta$  be

the indicated angles. We have

$ OC $	$=$	$R$
$ PC $	$=$	$r$
$ PH $	$=$	$r \sin \theta$
$ OH $	$=$	$R + r \cos \theta$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, ?), \theta \in [0, ?] \\ z = \rho \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, ?), \theta \in [0, ?] \\ z = \rho \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, ?), \theta \in [0, ?] \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, ?), \theta \in [0, ?] \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, ?] \\ z = \rho \sin \theta \end{cases}.$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, ?] \\ z = \rho \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, ?] \\ z = \rho \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, ?] \\ z = \rho \sin \theta \end{cases}.$$

## Example (Volume of toroid)




$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, 2\pi). \\ z = \rho \sin \theta \end{cases}$$

Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

## Example (Volume of toroid)




$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases} \quad \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, 2\pi).$$

Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, 2\pi). \\ z = \rho \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, 2\pi). \\ z = \rho \sin \theta \end{cases}$$

## Example (Volume of toroid)




$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, ?], \phi \in [0, 2\pi), \theta \in [0, 2\pi). \\ z = \rho \sin \theta \end{cases}$$

Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

## Example (Volume of toroid)




$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases} \quad \rho \in [0, \ r], \phi \in [0, \ 2\pi), \theta \in [0, \ 2\pi).$$

Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

## Example (Volume of toroid)




Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$

$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, r], \phi \in [0, 2\pi], \theta \in [0, 2\pi]. \\ z = \rho \sin \theta \end{cases}$$

Let  $\mathbf{f}$  be the map participating in the parametrization of  $T$ .

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$


$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, r], \phi \in [0, 2\pi], \theta \in [0, 2\pi]. \\ z = \rho \sin \theta \end{cases}$$

Let  $\mathbf{f}$  be the map participating in the parametrization of  $T$ .

$$\theta = ? \quad \phi = ? \quad \rho = ?$$

$$\text{Vol}(T) = \int_{\theta=?}^{} \int_{\phi=?}^{} \int_{\rho=?}^{} \det(J_{\mathbf{f}}) d\rho d\phi d\theta$$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$


$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, r], \phi \in [0, 2\pi], \theta \in [0, 2\pi]. \\ z = \rho \sin \theta \end{cases}$$

Let  $\mathbf{f}$  be the map participating in the parametrization of  $T$ .

$$\theta = 2\pi, \phi = 2\pi, \rho = r$$

$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(J_{\mathbf{f}}) d\rho d\phi d\theta$$

## Example (Volume of toroid)



Find the volume of a toroid  $T$  (the inside of a torus  $S$ ) with major radius  $R$  and minor radius  $r$ .

$$S : \begin{cases} x = (R + r \cos \theta) \cos \phi \\ y = (R + r \cos \theta) \sin \phi \\ z = r \sin \theta \end{cases}$$


$$T : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi & \rho \in [0, r], \phi \in [0, 2\pi], \theta \in [0, 2\pi]. \\ z = \rho \sin \theta \end{cases}$$

Let  $\mathbf{f}$  be the map participating in the parametrization of  $T$ .

$$\theta = 2\pi, \phi = 2\pi, \rho = r$$

$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$

## Example (Volume of toroid)




Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$\mathbf{f} : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^{r=r} \det(J_f) d\rho d\phi d\theta$$

## Example (Volume of toroid)



$$\theta = 2\pi \quad \phi = 2\pi \quad \rho = r$$


$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(\mathbf{J}_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$\mathbf{f} : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$\mathbf{J}_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)




$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{aligned} x &= (R + \rho \cos \theta) \cos \phi \\ y &= (R + \rho \cos \theta) \sin \phi \\ z &= \rho \sin \theta \end{aligned}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)




$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)




$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{aligned} x &= (R + \rho \cos \theta) \cos \phi \\ y &= (R + \rho \cos \theta) \sin \phi \\ z &= \rho \sin \theta \end{aligned}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)




$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^{\rho=r} \det(J_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)




$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^{r} \det(J_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)




$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^{\rho=r} \det(J_f) d\rho d\phi d\theta$$


$$J_f = \begin{pmatrix} ? & ? & ? \\ ? & ? & ? \\ ? & ? & ? \end{pmatrix}$$

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{\theta=2\pi} \int_{\phi=0}^{\phi=2\pi} \int_{\rho=0}^{\rho=r} \det(J_f) d\rho d\phi d\theta$$

$$J_f = \begin{pmatrix} \cos \theta \cos \phi & -A \sin \phi & -\rho \sin \theta \cos \phi \\ \cos \theta \sin \phi & A \cos \phi & -\rho \sin \theta \sin \phi \\ \sin \theta & 0 & \rho \cos \theta \end{pmatrix}$$


where we have set  $A = R + \rho \cos \theta$ .

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \int_0^r ? d\rho d\phi d\theta$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

where we have set  $A = R + \rho \cos \theta$ .

Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$


$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

$$2\pi \quad 2\pi \quad r$$

$$\int_0^{2\pi} \int_0^{2\pi} \int_0^r ? d\rho d\phi d\theta$$

$$J_f = \begin{pmatrix} \cos \theta \cos \phi & -A \sin \phi & -\rho \sin \theta \cos \phi \\ \cos \theta \sin \phi & A \cos \phi & -\rho \sin \theta \sin \phi \\ \sin \theta & 0 & \rho \cos \theta \end{pmatrix}$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho A d\rho d\phi d\theta$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

where we have set  $A = R + \rho \cos \theta$ .


Find volume of toroid  $T$ , major radius  $R$  minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

$$J_f = \begin{pmatrix} \cos \theta \cos \phi & -A \sin \phi & -\rho \sin \theta \cos \phi \\ \cos \theta \sin \phi & A \cos \phi & -\rho \sin \theta \sin \phi \\ \sin \theta & 0 & \rho \cos \theta \end{pmatrix}$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$


Find volume of toroid  $T$ , major radius  $R$ , minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

$$\int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta$$

# Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta$$

$$= \int_0^{2\pi} \int_0^{2\pi} \left[ ? \right]_{\rho=0}^{\rho=r} d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$ , minor radius  $r$ .


$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

$$\int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta$$

$$= \int_0^{2\pi} \int_0^{2\pi} \left[ ? \right]_{\rho=0}^{\rho=r} d\phi d\theta$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta$$


$$= \int_0^{2\pi} \int_0^{2\pi} \left[ \frac{R\rho^2}{2} + \frac{\rho^3}{3} \cos \theta \right]_{\rho=0}^r d\phi d\theta$$

Find volume of toroid  $T$ , major radius  $R$ , minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

## Example (Volume of toroid)



$$\text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta$$

$$= \int_0^{2\pi} \int_0^{2\pi} \left[ \frac{R\rho^2}{2} + \frac{\rho^3}{3} \cos \theta \right]_{\rho=0}^r d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\phi d\theta$$

$$= 2\pi \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\theta$$


Find volume of toroid  $T$ , major radius  $R$ , minor radius  $r$ .

$$f : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$

$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$


$$= \int_0^{2\pi} \int_0^{2\pi} \left[ \frac{R\rho^2}{2} + \frac{\rho^3}{3} \cos \theta \right]_{\rho=0}^r d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\phi d\theta$$

# Example (Volume of toroid)



$$\begin{aligned}
 & \text{Find volume of toroid } T, \text{ major radius } R, \text{ minor radius } r. \\
 & \mathbf{f} : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases} \\
 & J_{\mathbf{f}} = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix} \\
 & \text{Vol}(T) = \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_{\mathbf{f}}) d\rho d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta \\
 & = \int_0^{2\pi} \int_0^{2\pi} \left[ \frac{R\rho^2}{2} + \frac{\rho^3}{3} \cos \theta \right]_{\rho=0}^r d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\phi d\theta \\
 & = 2\pi \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\theta = 2\pi \int_0^{2\pi} \frac{Rr^2}{2} d\theta
 \end{aligned}$$

## Example (Volume of toroid)



$$\begin{aligned}
 & \theta = 2\pi \quad \phi = 2\pi \quad \rho = r \\
 \text{Vol}(T) &= \int_{\theta=0}^{2\pi} \int_{\phi=0}^{2\pi} \int_{\rho=0}^r \det(J_f) d\rho d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta \\
 &= \int_0^{2\pi} \int_0^{2\pi} \left[ \frac{R\rho^2}{2} + \frac{\rho^3}{3} \cos \theta \right]_{\rho=0}^r d\phi d\theta = \int_0^{2\pi} \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\phi d\theta \\
 &= 2\pi \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\theta = 2\pi \int_0^{2\pi} \frac{Rr^2}{2} d\theta = 2Rr^2\pi^2
 \end{aligned}$$

Find volume of toroid  $T$ , major radius  $R$ , minor radius  $r$ .

$$\mathbf{f} : \begin{cases} x = (R + \rho \cos \theta) \cos \phi \\ y = (R + \rho \cos \theta) \sin \phi \\ z = \rho \sin \theta \end{cases}$$


$$J_f = \begin{pmatrix} \frac{\partial x}{\partial \rho} & \frac{\partial x}{\partial \phi} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial \rho} & \frac{\partial y}{\partial \phi} & \frac{\partial y}{\partial \theta} \\ \frac{\partial z}{\partial \rho} & \frac{\partial z}{\partial \phi} & \frac{\partial z}{\partial \theta} \end{pmatrix}$$

$$\int_0^{2\pi} \int_0^{2\pi} \int_0^r \rho(R + \rho \cos \theta) d\rho d\phi d\theta$$

$$\int_0^{2\pi} \int_0^{2\pi} \left( \frac{Rr^2}{2} + \frac{r^3}{3} \cos \theta \right) d\phi d\theta$$

$$2\pi \int_0^{2\pi} \frac{Rr^2}{2} d\theta = 2Rr^2\pi^2$$

## Example



Find the volume of the horn given by

$$\begin{aligned}x &= (2 + \rho \cos \theta) \cos \phi \\y &= (2 + \rho \cos \theta) \sin \phi \\z &= \rho \sin \theta + \frac{\phi}{3}\end{aligned},$$

$$\theta \in [0, 2\pi], \phi \in [0, 3\pi], \rho \in \left[0, \frac{\phi}{9}\right].$$

## Theorem (Variable change in multivariable integrals)

*f - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .*

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy.$

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy.$

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid
  - without positivity requirements (arranged by compensating with minus sign when changing boundaries of integration)

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, **one-to-one**,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid
  - without positivity requirements (arranged by compensating with minus sign when changing boundaries of integration)
  - and **without requiring that  $f$  be one to one** (compensated by neutralizing contributions arising from sign changes of  $f'(y)$ ).

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid
  - without positivity requirements (arranged by compensating with minus sign when changing boundaries of integration)
  - and without requiring that  $f$  be one to one (compensated by neutralizing contributions arising from sign changes of  $f'(y)$ ).
- Similarly integration can be generalized so multivar. subst. holds

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid
  - without positivity requirements (arranged by compensating with minus sign when changing boundaries of integration)
  - and without requiring that  $f$  be one to one (compensated by neutralizing contributions arising from sign changes of  $f'(y)$ ).
- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid
  - without positivity requirements (arranged by compensating with minus sign when changing boundaries of integration)
  - and without requiring that  $f$  be one to one (compensated by neutralizing contributions arising from sign changes of  $f'(y)$ ).
- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- One-variable subst. rule:  $\int_{f(a)}^{f(b)} h(x) dx = \int_a^b h(f(y)) f'(y) dy$ .
- The one-variable substitution rule is valid
  - without positivity requirements (arranged by compensating with minus sign when changing boundaries of integration)
  - and without requiring that  $f$  be one to one (compensated by neutralizing contributions arising from sign changes of  $f'(y)$ ).
- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).
- When using the above generalization of  $\int$ , one writes

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \wedge \dots \wedge dx_n .$$

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).
- When using the above generalization of  $\int$ , one writes
 
$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \wedge \dots \wedge dx_n .$$
- The wedge sign  $\wedge$  stands for exterior product.

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).

- When using the above generalization of  $\int$ , one writes

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \wedge \dots \wedge dx_n .$$

- The wedge sign  $\wedge$  stands for exterior product.
- We will skip the theoretical details of the above generalization.

## Theorem (Variable change in multivariable integrals)

$f$  - smooth, one-to-one,  $\mathbf{f}(\mathcal{R}) = \mathcal{S}$ ,  $\det(J_{\mathbf{f}}(\mathbf{y})) \geq 0$ .

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \dots dx_n = \int_{\mathcal{R}} \cdots \int h(f_1, \dots, f_n) \det(J_{\mathbf{f}}(\mathbf{y})) dy_1 \dots dy_n,$$

- Similarly integration can be generalized so multivar. subst. holds
  - without positivity of  $\det(J_{\mathbf{f}})$  (arranged by compensating with minus sign when changing orientation of spaces),
  - without requiring that  $\mathbf{f}$  be one to one (compensated by neutralizing contributions arising from sign changes of  $\det J_{\mathbf{f}}$ ).

- When using the above generalization of  $\int$ , one writes

$$\int_S \cdots \int h(x_1, \dots, x_n) dx_1 \wedge \dots \wedge dx_n .$$

- The wedge sign  $\wedge$  stands for exterior product.
- We will skip the theoretical details of the above generalization.
- However we will learn to compute with  $\wedge$  in practice.