

Calculus III

Lecture 4

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

1

Equations of Lines

- Line from point and direction
- Line from two points

Outline

1 Equations of Lines

- Line from point and direction
- Line from two points

2 Equations of planes

- Plane from point and normal
- Plane from two directions
- Plane from three points

Outline

1 Equations of Lines

- Line from point and direction
- Line from two points

2 Equations of planes

- Plane from point and normal
- Plane from two directions
- Plane from three points

3 Distances, Angles, Parallelism, Incidence

- Distance Between Point and Line
- Parallel Lines
- Angle Between Lines
- Distance Between Skew Lines
- Distance Between Plane and Parallel Line
- Angle Between Plane and Line
- Parallel Planes
- Angle Between Planes

License to use and redistribute


These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
`https://creativecommons.org/licenses/by/3.0/us/`
and the links therein.

Main Questions



What condition(s) should

- the position vector
- the coordinates

of a point satisfy for it to be on a specific


- line L
- plane \mathcal{P} ?

Condition(s) in terms of:


- position vector \Rightarrow vector (system of) equations;
- coordinates \Rightarrow scalar equations.

Line from Point and Direction

- Suppose we have line L that passes through point P_0 and has non-zero direction \mathbf{u} .




Line from Point and Direction




- Suppose we have line L that passes through point P_0 and has non-zero direction \mathbf{u} .
- Denote by $\mathbf{r}_0 = \mathbf{OP}_0$ the position vector of P_0 .

Line from Point and Direction




- Suppose we have line L that passes through point P_0 and has non-zero direction \mathbf{u} .
- Denote by $\mathbf{r}_0 = \mathbf{OP}_0$ the position vector of P_0 .
- P with position vector \mathbf{r} is on $L \Leftrightarrow$

Line from Point and Direction




- Suppose we have line L that passes through point P_0 and has non-zero direction \mathbf{u} .
- Denote by $\mathbf{r}_0 = \mathbf{OP}_0$ the position vector of P_0 .
- P with position vector \mathbf{r} is on $L \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P}$ has the same direction as $\mathbf{u} \Leftrightarrow$

Line from Point and Direction




- Suppose we have line L that passes through point P_0 and has non-zero direction \mathbf{u} .
- Denote by $\mathbf{r}_0 = \mathbf{OP}_0$ the position vector of P_0 .
- P with position vector \mathbf{r} is on $L \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P}$ has the same direction as $\mathbf{u} \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P}$ is a scalar multiple of $\mathbf{u} \Leftrightarrow$

Line from Point and Direction



- Suppose we have line L that passes through point P_0 and has non-zero direction \mathbf{u} .
- Denote by $\mathbf{r}_0 = \mathbf{OP}_0$ the position vector of P_0 .
- P with position vector \mathbf{r} is on $L \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P}$ has the same direction as $\mathbf{u} \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P}$ is a scalar multiple of $\mathbf{u} \Leftrightarrow$
- $\mathbf{r} - \mathbf{r}_0 = t\mathbf{u}$ for some real number t .

Line from Point and Direction




- Point $P_0(x_0, y_0, z_0)$, $\mathbf{r}_0 = (x_0, y_0, z_0)$;
- Direction $\mathbf{u} = (u_1, u_2, u_3)$.

L - line with direction
 \mathbf{u} passing through P_0

Definition

Line from Point and Direction




- Point $P_0(x_0, y_0, z_0)$, $\mathbf{r}_0 = (x_0, y_0, z_0)$;
- Direction $\mathbf{u} = (u_1, u_2, u_3)$. $P(x, y, z)$ with position vector \mathbf{r} is on $L \Leftrightarrow$

L - line with direction \mathbf{u} passing through P_0

Definition

Line from Point and Direction




- Point $P_0(x_0, y_0, z_0)$, $\mathbf{r}_0 = (x_0, y_0, z_0)$;
- Direction $\mathbf{u} = (u_1, u_2, u_3)$. $P(x, y, z)$ with position vector \mathbf{r} is on $L \Leftrightarrow$

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{u} \Leftrightarrow$$

L - line with direction \mathbf{u} passing through P_0

Definition

Line from Point and Direction



L - line with direction \mathbf{u} passing through P_0


- Point $P_0(x_0, y_0, z_0)$, $\mathbf{r}_0 = (x_0, y_0, z_0)$;
- Direction $\mathbf{u} = (u_1, u_2, u_3)$. $P(x, y, z)$ with position vector \mathbf{r} is on $L \Leftrightarrow$

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{u} \Leftrightarrow$$

$$(x, y, z) = (x_0, y_0, z_0) + t(u_1, u_2, u_3) \Leftrightarrow$$

Definition

Line from Point and Direction



L - line with direction \mathbf{u} passing through P_0

- Point $P_0(x_0, y_0, z_0)$, $\mathbf{r}_0 = (x_0, y_0, z_0)$;
- Direction $\mathbf{u} = (u_1, u_2, u_3)$. $P(x, y, z)$ with position vector \mathbf{r} is on $L \Leftrightarrow$

$$\mathbf{r} = \mathbf{r}_0 + t\mathbf{u} \Leftrightarrow$$

$$(x, y, z) = (x_0, y_0, z_0) + t(u_1, u_2, u_3) \Leftrightarrow$$

Definition

The equations

$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases}, t \in \mathbb{R}$$

are called **parametric scalar equations** of the line L .

$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases} \implies \boxed{\frac{x - x_0}{u_1} = \frac{y - y_0}{u_2} = \frac{z - z_0}{u_3}} \quad \text{Symmetric equations}$$

$$\begin{cases} x = x_0 + tu_1 \\ y = y_0 + tu_2 \\ z = z_0 + tu_3 \end{cases}$$

 \implies

$$\boxed{\frac{x - x_0}{u_1} = \frac{y - y_0}{u_2} = \frac{z - z_0}{u_3}}$$

Symmetric equations

- Caution! Symmetric equations are valid for $u_1, u_2, u_3 \neq 0$. For example if $u_2 = 0$ the equations should be:

$$\frac{x - x_0}{u_1} = \frac{z - z_0}{u_3} \quad \text{and} \quad y = y_0$$

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Parametric vectorial equation:

$\mathbf{r} =$

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Parametric vectorial equation:

$$\mathbf{r} = (1, 2, 3) + t(4, 5, 6) \leftrightarrow \mathbf{r} = (1 + 4t, 2 + 5t, 3 + 6t)$$

Parametric scalar equations:

$$\begin{cases} x = \\ y = \\ z = \end{cases}, \quad t \text{ real number.}$$

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Parametric vectorial equation:

$$\mathbf{r} = (1, 2, 3) + t(4, 5, 6) \leftrightarrow \mathbf{r} = (1 + 4t, 2 + 5t, 3 + 6t)$$

Parametric scalar equations:

$$x =$$

$$y =$$

$$z =$$

, t real number.

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Parametric vectorial equation:

$$\mathbf{r} = (1, 2, 3) + t(4, 5, 6) \leftrightarrow \mathbf{r} = (1 + 4t, 2 + 5t, 3 + 6t)$$

Parametric scalar equations:

$$\begin{cases} x = 1 + 4t \\ y = 2 + 5t \\ z = 3 + 6t \end{cases}, \quad t \text{ real number.}$$

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Parametric vectorial equation:

$$\mathbf{r} = (1, 2, 3) + t(4, 5, 6) \leftrightarrow \mathbf{r} = (1 + 4t, 2 + 5t, 3 + 6t)$$

Parametric scalar equations:

$$\begin{cases} x = 1 + 4t \\ y = 2 + 5t \\ z = 3 + 6t \end{cases}, \quad t \text{ real number.}$$

Symmetric equations:

Example

L - line with direction $\mathbf{u} = (4, 5, 6)$ passing through $P_0(1, 2, 3)$. Find

- a parametric vectorial equation of L ;
- a parametric scalar equation of L ;
- symmetric equations of L .

Parametric vectorial equation:

$$\mathbf{r} = (1, 2, 3) + t(4, 5, 6) \leftrightarrow \mathbf{r} = (1 + 4t, 2 + 5t, 3 + 6t)$$


Parametric scalar equations:

$$\begin{cases} x = 1 + 4t \\ y = 2 + 5t \\ z = 3 + 6t \end{cases}, \quad t \text{ real number.}$$

Symmetric equations:


$$\frac{x - 1}{4} = \frac{y - 2}{5} = \frac{z - 3}{6}.$$

Line from Two Points




- Given: distinct points P_0 and P_1 , position vectors \mathbf{r}_0 and \mathbf{r}_1 .
- Goal: write equations of line L through P_0 and P_1 .

Line from Two Points



- Given: distinct points P_0 and P_1 , position vectors \mathbf{r}_0 and \mathbf{r}_1 .
- Goal: write equations of line L through P_0 and P_1 .
- Direction of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0$.

Line from Two Points




- Given: distinct points P_0 and P_1 , position vectors \mathbf{r}_0 and \mathbf{r}_1 .
- Goal: write equations of line L through P_0 and P_1 .
- Direction of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0$.

Definition

Parametric equation of a line L :

$$\mathbf{r} = \mathbf{r}_0 + t(\mathbf{r}_1 - \mathbf{r}_0) \quad \Leftrightarrow \quad \mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1$$

Line from Two Points




- Given: distinct points P_0 and P_1 , position vectors \mathbf{r}_0 and \mathbf{r}_1 .
- Goal: write equations of line L through P_0 and P_1 .
- Direction of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0$.

Definition

Parametric equation of a line L :

$$\mathbf{r} = \mathbf{r}_0 + t(\mathbf{r}_1 - \mathbf{r}_0) \quad \Leftrightarrow \quad \mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1$$

Line from Two Points



- Given: distinct points P_0 and P_1 , position vectors \mathbf{r}_0 and \mathbf{r}_1 .
- Goal: write equations of line L through P_0 and P_1 .
- Direction of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0$.
- $\mathbf{u} = (x_1 - x_0, y_1 - y_0, z_1 - z_0)$.

Definition

Parametric equation of a line L :

$$\mathbf{r} = \mathbf{r}_0 + t(\mathbf{r}_1 - \mathbf{r}_0) \Leftrightarrow \mathbf{r} = (1 - t)\mathbf{r}_0 + t\mathbf{r}_1$$

Parametric scalar equations of a line L :

$$\begin{cases} x = x_0 + t(x_1 - x_0) \\ y = y_0 + t(y_1 - y_0) \\ z = z_0 + t(z_1 - z_0) \end{cases} \Leftrightarrow \begin{cases} x = (1 - t)x_0 + tx_1 \\ y = (1 - t)y_0 + ty_1 \\ z = (1 - t)z_0 + tz_1 \end{cases}, \quad t \text{ real number.}$$

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L :

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.
- Parametric vector equation:

$\mathbf{r} =$

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.

- Parametric vector equation:

$$\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t).$$

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.
- Parametric vector equation:
 $\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t)$.
- Parametric scalar equations:

$$\begin{cases} x = \\ y = \\ z = \end{cases}, \quad t \text{ real number.}$$

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.
- Parametric vector equation:
 $\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t)$.
- Parametric scalar equations:

$$\begin{cases} x = 1 + 4t \\ y = 2 \\ z = 3 - 2t \end{cases}, \quad t \text{ real number.}$$

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.

- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.
- Parametric vector equation:
 $\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t)$.
- Parametric scalar equations:

$$\begin{cases} x = 1 + 4t \\ y = 2 \\ z = 3 - 2t \end{cases}, \quad t \text{ real number.}$$

- Symmetric equations:

Example

Write the equations of line L through $P_0(1, 2, 3)$ and $P_1(5, 2, 1)$.


- Direction \mathbf{u} of L : $\mathbf{u} = \mathbf{r}_1 - \mathbf{r}_0 = (4, 0, -2)$.
- Parametric vector equation:
 $\mathbf{r} = (1, 2, 3) + t(4, 0, -2) \Leftrightarrow \mathbf{r} = (1 + 4t, 2, 3 - 2t)$.
- Parametric scalar equations:

$$\begin{cases} x = 1 + 4t \\ y = 2 \\ z = 3 - 2t \end{cases}, \quad t \text{ real number.}$$

- Symmetric equations:


$$\frac{x - 1}{4} = \frac{z - 3}{-2} \quad \text{and} \quad y = 2.$$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
- Direction \mathbf{n} , non-zero vector.
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
- Direction \mathbf{n} , non-zero vector.
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
- Direction \mathbf{n} , non-zero vector.
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P}$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
- Direction \mathbf{n} , non-zero vector.
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P_0P} = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
 $\mathbf{r}_0 = (x_0, y_0, z_0)$
- Direction \mathbf{n} , non-zero vector. $\mathbf{n} = (a, b, c)$
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P}_0P = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$
- **Implicit vectorial equation:** $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.

Plane from Point and Normal



- Point P_0 , with position vector \mathbf{r}_0 ;
 $\mathbf{r}_0 = (x_0, y_0, z_0)$
- Direction \mathbf{n} , non-zero vector. $\mathbf{n} = (a, b, c)$
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P_0P} = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$
- **Implicit vectorial equation:** $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- A point $P(x, y, z)$ is on $\mathcal{P} \Leftrightarrow$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
 $\mathbf{r}_0 = (x_0, y_0, z_0)$
- Direction \mathbf{n} , non-zero vector. $\mathbf{n} = (a, b, c)$
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P} = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$
- **Implicit vectorial equation:** $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- A point $P(x, y, z)$ is on $\mathcal{P} \Leftrightarrow$

Definition (Implicit scalar equation)

$$(x - x_0, y - y_0, z - z_0) \cdot (a, b, c) = 0$$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
 $\mathbf{r}_0 = (x_0, y_0, z_0)$
- Direction \mathbf{n} , non-zero vector. $\mathbf{n} = (a, b, c)$
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P} = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$
- **Implicit vectorial equation:** $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- A point $P(x, y, z)$ is on $\mathcal{P} \Leftrightarrow$

Definition (Implicit scalar equation)

$$(x - x_0, y - y_0, z - z_0) \cdot (a, b, c) = 0$$

Plane from Point and Normal




- Point P_0 , with position vector \mathbf{r}_0 ;
 $\mathbf{r}_0 = (x_0, y_0, z_0)$
- Direction \mathbf{n} , non-zero vector. $\mathbf{n} = (a, b, c)$
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P_0P} = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$
- **Implicit vectorial equation:** $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- A point $P(x, y, z)$ is on $\mathcal{P} \Leftrightarrow$

Definition (Implicit scalar equation)

$$(x - x_0, y - y_0, z - z_0) \cdot (a, b, c) = 0$$

Plane from Point and Normal



- Point P_0 , with position vector \mathbf{r}_0 ;
 $\mathbf{r}_0 = (x_0, y_0, z_0)$
- Direction \mathbf{n} , non-zero vector. $\mathbf{n} = (a, b, c)$
- Goal: describe plane passing through P_0 and orthogonal to \mathbf{n} .
- Point P with position \mathbf{r} is on $\mathcal{P} \Leftrightarrow$
- $\mathbf{P}_0\mathbf{P} = \mathbf{r} - \mathbf{r}_0$ is orthogonal (normal) to $\mathbf{n} \Leftrightarrow$
- **Implicit vectorial equation:** $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- A point $P(x, y, z)$ is on $\mathcal{P} \Leftrightarrow$

Definition (Implicit scalar equation)

$$(x - x_0, y - y_0, z - z_0) \cdot (a, b, c) = 0 \Leftrightarrow$$

$$\mathbf{a}(x - x_0) + \mathbf{b}(y - y_0) + \mathbf{c}(z - z_0) = 0$$

Example

Find an equation of the plane

- passing through $P_0(1, 2, 3)$
- and perpendicular (normal) to the direction $\mathbf{n} = (6, 5, 4)$.

Example

Find an equation of the plane

- passing through $P_0(1, 2, 3)$
- and perpendicular (normal) to the direction $\mathbf{n} = (6, 5, 4)$.

$$6(x - 1) + 5(y - 2) + 4(z - 3) = 0$$

$$6x + 5y + 4z = 28$$

The general equation of a plane is given by:

$$ax + by + cz = d.$$

The coefficients a, b, c are

Example

Find an equation of the plane

- passing through $P_0(1, 2, 3)$
- and perpendicular (normal) to the direction $\mathbf{n} = (6, 5, 4)$.

$$6(x - 1) + 5(y - 2) + 4(z - 3) = 0$$

$$6x + 5y + 4z = 28$$


The general equation of a plane is given by:

$$ax + by + cz = d.$$

The coefficients a, b, c are the components of normal to the plane,


$$\mathbf{n} = (a, b, c) .$$

Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .


Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .




Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .


Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Normal direction $\mathbf{n} = \mathbf{u} \times \mathbf{v} \neq \mathbf{0}$.

Plane from Point and two Directions




- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Normal direction $\mathbf{n} = \mathbf{u} \times \mathbf{v} \neq \mathbf{0}$.

Implicit equation: $P(\mathbf{r})$ is on $\mathcal{P} \iff (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$

Plane from Point and two Directions




- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Normal direction $\mathbf{n} = \mathbf{u} \times \mathbf{v} \neq \mathbf{0}$.

Implicit equation: $P(\mathbf{r})$ is on $\mathcal{P} \iff (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$ Interpretation:

$$\text{Vol}(R(\mathbf{r} - \mathbf{r}_0, \mathbf{u}, \mathbf{v})) = 0$$


Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

$P(\mathbf{r})$ is on the plane $\mathcal{P} \Leftrightarrow$


Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

$P(\mathbf{r})$ is on the plane $\mathcal{P} \Leftrightarrow$
 $\mathbf{P}_0\mathbf{P}$ is a combination of $\mathbf{u}, \mathbf{v} \Leftrightarrow$

Plane from Point and two Directions




- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

$P(\mathbf{r})$ is on the plane $\mathcal{P} \Leftrightarrow$

$\mathbf{P}_0\mathbf{P}$ is a combination of $\mathbf{u}, \mathbf{v} \Leftrightarrow$

There are scalars s, t such that $\mathbf{r} - \mathbf{r}_0 = s\mathbf{u} + t\mathbf{v} \Leftrightarrow$

Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

$P(\mathbf{r})$ is on the plane $\mathcal{P} \Leftrightarrow$

$\mathbf{P}_0\mathbf{P}$ is a combination of \mathbf{u} , $\mathbf{v} \Leftrightarrow$


There are scalars s , t such that $\mathbf{r} - \mathbf{r}_0 = s\mathbf{u} + t\mathbf{v} \Leftrightarrow$

Parametric equation:

$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

for some parameters s and t .

Plane from Point and two Directions




- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Parametric equation:

$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

Plane from Point and two Directions




- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Parametric equation:

$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

Let $P_0(x_0, y_0, z_0)$, $P(x, y, z)$ $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$.

Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Parametric equation:


$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

Let $P_0(x_0, y_0, z_0)$, $P(x, y, z)$ $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$. \Rightarrow

Parametric scalar equations:

$$\begin{cases} x = x_0 + su_1 + tv_1 \\ y = y_0 + su_2 + tv_2 \\ z = z_0 + su_3 + tv_3 \end{cases}, \text{ for } s, t \text{ real parameters.}$$

Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Parametric equation:


$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

Let $P_0(x_0, y_0, z_0)$, $P(x, y, z)$ $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$. \Rightarrow

Parametric scalar equations:

$$\left| \begin{array}{l} x = x_0 + su_1 + tv_1 \\ y = y_0 + su_2 + tv_2 \\ z = z_0 + su_3 + tv_3 \end{array} \right. , \text{ for } s, t \text{ real parameters.}$$

Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Parametric equation:


$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

Let $P_0(x_0, y_0, z_0)$, $P(x, y, z)$ $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$. \Rightarrow

Parametric scalar equations:

$$\begin{cases} x = x_0 + su_1 + tv_1 \\ y = y_0 + su_2 + tv_2 \\ z = z_0 + su_3 + tv_3 \end{cases}, \text{ for } s, t \text{ real parameters.}$$

Plane from Point and two Directions



- Given: point P_0 with position vector \mathbf{r}_0 .
- Non-parallel directions \mathbf{u} and \mathbf{v} .
- Goal: give equations of plane \mathcal{P} through P_0 and parallel to both \mathbf{u} and \mathbf{v} .

Parametric equation:

$$\mathbf{r} = \mathbf{r}_0 + s\mathbf{u} + t\mathbf{v}$$

Let $P_0(x_0, y_0, z_0)$, $P(x, y, z)$ $\mathbf{u} = (u_1, u_2, u_3)$, $\mathbf{v} = (v_1, v_2, v_3)$. \Rightarrow

Parametric scalar equations:

$$\left| \begin{array}{lcl} x = & x_0 + su_1 + tv_1 \\ y = & y_0 + su_2 + tv_2 \\ z = & z_0 + su_3 + tv_3 \end{array} \right. , \text{ for } s, t \text{ real parameters.}$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$\mathbf{n} =$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0 \iff 4x + y + 2z = 12$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0 \iff 4x + y + 2z = 12$$

Parametric vectorial equation:

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0 \iff 4x + y + 2z = 12$$

Parametric vectorial equation:

$$(x, y, z) = (1, 2, 3) + s(-1, 0, 2) + t(0, -2, 1)$$

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0 \iff 4x + y + 2z = 12$$

Parametric vectorial equation:

$$(x, y, z) = (1, 2, 3) + s(-1, 0, 2) + t(0, -2, 1)$$

Parametric scalar equations:

Example

Find equations of a plane passing through $P_0(1, 2, 3)$ and parallel to the vectors $\mathbf{u} = (-1, 0, 2)$, $\mathbf{v} = (0, -2, 1)$.

$$\mathbf{n} = \mathbf{u} \times \mathbf{v} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -1 & 0 & 2 \\ 0 & -2 & 1 \end{vmatrix} = 4\mathbf{i} + \mathbf{j} + 2\mathbf{k}$$

\Rightarrow implicit scalar equation given by:

$$4(x - 1) + 1(y - 2) + 2(z - 3) = 0 \iff 4x + y + 2z = 12$$


Parametric vectorial equation:

$$(x, y, z) = (1, 2, 3) + s(-1, 0, 2) + t(0, -2, 1)$$

Parametric scalar equations:


$$\begin{cases} x = 1 - s \\ y = 2 - 2t \\ z = 3 + 2s + t \end{cases} \quad s, t \text{ real parameters.}$$

Plane from Three Points




- Given: three non-collinear points $P_0(\mathbf{r}_0)$, $P_1(\mathbf{r}_1)$, $P_2(\mathbf{r}_2)$.
- Goal: find equations for plane \mathcal{P} passing through P_0 , P_1 , and P_2 .

Plane from Three Points



- Given: three non-collinear points $P_0(\mathbf{r}_0)$, $P_1(\mathbf{r}_1)$, $P_2(\mathbf{r}_2)$.
- Goal: find equations for plane \mathcal{P} passing through P_0 , P_1 , and P_2 .
- The plane is parallel to $\mathbf{u} = \mathbf{P}_1 - \mathbf{P}_0 = \mathbf{r}_1 - \mathbf{r}_0$ and passing through $P_0 \Rightarrow$ this problem was solved previously.

Plane from Three Points



- Given: three non-collinear points $P_0(\mathbf{r}_0)$, $P_1(\mathbf{r}_1)$, $P_2(\mathbf{r}_2)$.
- Goal: find equations for plane \mathcal{P} passing through P_0 , P_1 , and P_2 .
- The plane is parallel to $\mathbf{u} = \mathbf{P}_0\mathbf{P}_1 = \mathbf{r}_1 - \mathbf{r}_0$ and passing through $P_0 \Rightarrow$ this problem was solved previously.

Normal $\mathbf{n} = \mathbf{u} \times \mathbf{v} = (\mathbf{r}_1 - \mathbf{r}_0) \times (\mathbf{r}_2 - \mathbf{r}_0)$


Implicit equation:

$$(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

$$(\mathbf{r} - \mathbf{r}_0) \cdot [(\mathbf{r}_1 - \mathbf{r}_0) \times (\mathbf{r}_2 - \mathbf{r}_0)] = 0$$

$$\text{Vol}(R(\mathbf{P}_0\mathbf{P}, \mathbf{P}_0\mathbf{P}_1, \mathbf{P}_0\mathbf{P}_2)) = 0$$

Plane from Three Points



- Given: three non-collinear points $P_0(\mathbf{r}_0)$, $P_1(\mathbf{r}_1)$, $P_2(\mathbf{r}_2)$.
- Goal: find equations for plane \mathcal{P} passing through P_0 , P_1 , and P_2 .
- The plane is parallel to $\mathbf{u} = \mathbf{P}_0\mathbf{P}_1 = \mathbf{r}_1 - \mathbf{r}_0$ and passing through $P_0 \Rightarrow$ this problem was solved previously.

Implicit equation: $(\mathbf{r} - \mathbf{r}_0) \cdot [(\mathbf{r}_1 - \mathbf{r}_0) \times (\mathbf{r}_2 - \mathbf{r}_0)] = 0$


Let the points have coordinates $P_0(x_0, y_0, z_0)$, $P_1(x_1, y_1, z_1)$, $P_2(x_2, y_2, z_2)$. $P(x, y, z)$ is on plane \mathcal{P} :

Implicit scalar equation:

$$\begin{vmatrix} x - x_0 & y - y_0 & z - z_0 \\ x_1 - x_0 & y_1 - y_0 & z_1 - z_0 \\ x_2 - x_0 & y_2 - y_0 & z_2 - z_0 \end{vmatrix} = 0.$$

Example


Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0 , P_1 , P_2 (i.e., plane with prescribed x, y, z -intercepts).



Example

Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0 , P_1 , P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to




Example

Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$




Example

Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$

Normal:



Example


Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$

Normal:

$$\mathbf{n} = \mathbf{P}_0\mathbf{P}_1 \times \mathbf{P}_0\mathbf{P}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a & b & 0 \\ -a & 0 & c \end{vmatrix} = bci + acj + abk.$$



Example


Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$

Normal:

$$\mathbf{n} = \mathbf{P}_0\mathbf{P}_1 \times \mathbf{P}_0\mathbf{P}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a & b & 0 \\ -a & 0 & c \end{vmatrix} = bci + acj + abk.$$



Implicit scalar equation of plane:

Example


Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$

Normal:

$$\mathbf{n} = \mathbf{P}_0\mathbf{P}_1 \times \mathbf{P}_0\mathbf{P}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a & b & 0 \\ -a & 0 & c \end{vmatrix} = bci + acj + abk.$$



Implicit scalar equation of plane:

$$(x - a, y, z) \cdot (bc, ac, ab) = 0$$

Example


Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$

Normal:

$$\mathbf{n} = \mathbf{P}_0\mathbf{P}_1 \times \mathbf{P}_0\mathbf{P}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a & b & 0 \\ -a & 0 & c \end{vmatrix} = bci + acj + abk.$$



Implicit scalar equation of plane:

$$\begin{aligned} (x - a, y, z) \cdot (bc, ac, ab) &= 0 \\ bcx + acy + abz &= abc \end{aligned}$$

Example


Let $P_0(a, 0, 0)$, $P_1(0, b, 0)$, $P_2(0, 0, c)$ be three points, $a, b, c \neq 0$. Find plane \mathcal{P} passing through P_0, P_1, P_2 (i.e., plane with prescribed x, y, z -intercepts).

\mathcal{P} : parallel to

$$\mathbf{P}_0\mathbf{P}_1 = (-a, b, 0), \mathbf{P}_0\mathbf{P}_2 = (-a, 0, c).$$

Normal:

$$\mathbf{n} = \mathbf{P}_0\mathbf{P}_1 \times \mathbf{P}_0\mathbf{P}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ -a & b & 0 \\ -a & 0 & c \end{vmatrix} = bci + acj + abk.$$




Implicit scalar equation of plane:

$$\begin{aligned} (x - a, y, z) \cdot (bc, ac, ab) &= 0 \\ bcx + acy + abz &= abc \\ \frac{x}{a} + \frac{y}{b} + \frac{z}{c} &= 1 \end{aligned}$$

Relationships between points lines and planes

- So far we studied the following geometric objects/
 - Points: $P(\mathbf{r})$.
 - Lines: $L: \mathbf{r} = \mathbf{r}_0 + t\mathbf{u}$
 - Planes: $\mathcal{P}: (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$
- We investigate the following relationships/geometric quantities:
 - Parallelism
 - Perpendicularity
 - Angles
 - Distances
 - Intersections


Point and line



Distance from P to L :

- Given: Point $P(r_1)$,
- line $L : \mathbf{r} = \mathbf{r}_0 + t\mathbf{u}$.
- Goal: find the distance between P and L .

Point and line




Distance from P to L :

- Given: Point $P(\mathbf{r}_1)$,
- line $L : \mathbf{r} = \mathbf{r}_0 + t\mathbf{u}$.
- Goal: find the distance between P and L .

$$d(P, L) = |\mathbf{orth}_{\mathbf{u}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

Point and line




- Given: Point $P(\mathbf{r}_1)$,
- line $L : \mathbf{r} = \mathbf{r}_0 + t\mathbf{u}$.
- Goal: find the distance between P and L .

Distance from P to L :

$$d(P, L) = |\mathbf{orth}_{\mathbf{u}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, L) = \left| \mathbf{r}_1 - \mathbf{r}_0 - \frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{u}}{\mathbf{u} \cdot \mathbf{u}} \mathbf{u} \right|$$

Point and line



- Given: Point $P(\mathbf{r}_1)$,
- line $L : \mathbf{r} = \mathbf{r}_0 + t\mathbf{u}$.
- Goal: find the distance between P and L .

Distance from P to L :


$$d(P, L) = |\text{orth}_{\mathbf{u}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, L) = \left| \mathbf{r}_1 - \mathbf{r}_0 - \frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{u}}{\mathbf{u} \cdot \mathbf{u}} \mathbf{u} \right|$$

$$d(P, L) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \times \mathbf{u}|}{|\mathbf{u}|}$$

Valid only in 3 dimensions!


Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

Distance between point and plane




- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :


$$d(P, \mathcal{P}) = |\mathbf{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$

Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .


Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = \frac{|\mathbf{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|}{1} = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$

Scalar equation:

$$\begin{aligned} \mathcal{P} : \quad a x + b y + c z + d &= (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0 \\ \mathbf{n} &= (a, b, c) \end{aligned}$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$


$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$

Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

$$\mathbf{n} = (a, b, c)$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$


$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$

Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

$$\mathbf{n} = (a, b, c)$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$


Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

$$\mathbf{n} = (a, b, c)$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$


Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

$$\mathbf{n} = (a, b, c)$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$


Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

$$\mathbf{n} = (a, b, c)$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

Distance between point and plane



- Given: point $P(\mathbf{r}_1) = (x_1, y_1, z_1)$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: find the distance between P and \mathcal{P} .

Distance from P to \mathcal{P} :

$$d(P, \mathcal{P}) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$

Scalar equation:

$$\mathcal{P} : ax + by + cz + d = (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$


$$\mathbf{n} = (a, b, c)$$

$$d(P, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|ax_1 + by_1 + cz_1 + d|}{\sqrt{a^2 + b^2 + c^2}}$$

Parallel lines

- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$.
- Goal: distance between lines.

Parallel lines




Parallel lines


- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$.
- Goal: distance between lines.

Parallel lines $L_1 \parallel L_2 \iff \mathbf{u}_1, \mathbf{u}_2$ collinear

$$\iff \boxed{\mathbf{u}_1 \times \mathbf{u}_2 = \mathbf{0}}$$



Parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$.
- Goal: distance between lines.

Parallel lines $L_1 \parallel L_2 \iff \mathbf{u}_1, \mathbf{u}_2$ collinear

$\iff \mathbf{u}_1 \times \mathbf{u}_2 = \mathbf{0}$ Distance:

$$d = d(L_1, L_2) = d(P_1, L_2) = d(P_2, L_1)$$

Parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- Goal: distance between lines.

Parallel lines $L_1 \parallel L_2 \iff \mathbf{u}_1, \mathbf{u}_2$ collinear
 $\iff \mathbf{u}_1 \times \mathbf{u}_2 = \mathbf{0}$ Distance:

$$d = d(L_1, L_2) = d(P_1, L_2) = d(P_2, L_1)$$


$$d = d(L_1, L_2) = |\text{orth}_{\mathbf{u}_1}(\mathbf{r}_2 - \mathbf{r}_1)|$$

$$d = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \times \mathbf{u}_1|}{|\mathbf{u}_1|} = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \times \mathbf{u}_2|}{|\mathbf{u}_2|}$$

Angle between lines

- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- Goal: find angle between L_1 and L_2 .

Perpendicular lines




Angle between lines

- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$.
- Goal: find angle between L_1 and L_2 .


Perpendicular lines $L_1 \perp L_2 \iff \mathbf{u}_1 \perp \mathbf{u}_2$

\iff

$$\mathbf{u}_1 \cdot \mathbf{u}_2 = 0$$



Angle between lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- Goal: find angle between L_1 and L_2 .


Perpendicular lines $L_1 \perp L_2 \iff \mathbf{u}_1 \perp \mathbf{u}_2$

\iff

$$\mathbf{u}_1 \cdot \mathbf{u}_2 = 0$$

Angle between lines

Angle between lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- Goal: find angle between L_1 and L_2 .


Perpendicular lines $L_1 \perp L_2 \iff \mathbf{u}_1 \perp \mathbf{u}_2$

\iff

$$\mathbf{u}_1 \cdot \mathbf{u}_2 = 0$$

Angle between lines α : angle between $L_1, L_2 \iff \alpha$: acute angle $\mathbf{u}_1, \mathbf{u}_2$

Angle between lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- Goal: find angle between L_1 and L_2 .

Perpendicular lines $L_1 \perp L_2 \iff \mathbf{u}_1 \perp \mathbf{u}_2$


\iff

$$\mathbf{u}_1 \cdot \mathbf{u}_2 = 0$$

Angle between lines α : angle between $L_1, L_2 \iff \alpha$: acute angle $\mathbf{u}_1, \mathbf{u}_2 \iff$



$$\alpha = \arccos \left(\frac{|\mathbf{u}_1 \cdot \mathbf{u}_2|}{|\mathbf{u}_1| |\mathbf{u}_2|} \right)$$

Distance between non-parallel lines





- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

Distance between non-parallel lines





- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .

Distance between non-parallel lines





- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .
- Distance b-n L_2 and points on \mathcal{P} is constant.

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .
- Distance b-n L_2 and points on \mathcal{P} is constant.
- Project L_2 orthogonally on \mathcal{P} ; let the projection be L_2' .

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .
- Distance b-n L_2 and points on \mathcal{P} is constant.
- Project L_2 orthogonally on \mathcal{P} ; let the projection be L'_2 .
- Let L'_2 and L_1 intersect in point Q_1 .

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .
- Distance b-n L_2 and points on \mathcal{P} is constant.
- Project L_2 orthogonally on \mathcal{P} ; let the projection be L'_2 .
- Let L'_2 and L_1 intersect in point Q_1 .
- Let Q_2 be the heel of the perpendicular from Q_1 onto Q_2 .

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .
- Distance b-n L_2 and points on \mathcal{P} is constant.
- Project L_2 orthogonally on \mathcal{P} ; let the projection be L'_2 .
- Let L'_2 and L_1 intersect in point Q_1 .
- Let Q_2 be the heel of the perpendicular from Q_1 onto L_2 .
- $\Rightarrow Q_1Q_2 = d(L_1, L_2)$.

Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- Construct plane \mathcal{P} with directions $\mathbf{u}_1, \mathbf{u}_2$ and passing through L_1 .
- Distance b-n L_2 and points on \mathcal{P} is constant.
- Project L_2 orthogonally on \mathcal{P} ; let the projection be L'_2 .
- Let L'_2 and L_1 intersect in point Q_1 .
- Let Q_2 be the heel of the perpendicular from Q_1 onto L_2 .
- $\Rightarrow Q_1Q_2 = d(L_1, L_2)$.
- $|Q_1Q_2| = d(L_1, L_2)$.


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $|Q_1 Q_2| = d(L_1, L_2)$.


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.


- $|Q_1Q_2| = d(L_1, L_2)$.
- $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2$

Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
 - The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
 - Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.
- $|Q_1Q_2| = d(L_1, L_2)$.
 - $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2 \Rightarrow \mathbf{Q}_1\mathbf{Q}_2$ is proportional to $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2$.


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines
 $= d(L_1, L_2) =$ shortest distance b-n
 points on the two lines.

- $|Q_1Q_2| = d(L_1, L_2)$.
- $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2 \Rightarrow \mathbf{Q}_1\mathbf{Q}_2$ is proportional to $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2$.
- Pick arbitrary points on L_1, L_2 - say, the base points $P_1(\mathbf{r}_1), P_2(\mathbf{r}_2)$.


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $|Q_1Q_2| = d(L_1, L_2)$.
- $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2 \Rightarrow \mathbf{Q}_1\mathbf{Q}_2$ is proportional to $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2$.
- Pick arbitrary points on L_1, L_2 - say, the base points $P_1(\mathbf{r}_1), P_2(\mathbf{r}_2)$.
- Let R be such that $\mathbf{Q}_1\mathbf{R} = \mathbf{P}_1\mathbf{P}_2 = \mathbf{r}_2 - \mathbf{r}_1$.


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $|Q_1Q_2| = d(L_1, L_2)$.
- $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2 \Rightarrow \mathbf{Q}_1\mathbf{Q}_2$ is proportional to $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2$.
- Pick arbitrary points on L_1, L_2 - say, the base points $P_1(\mathbf{r}_1), P_2(\mathbf{r}_2)$.
- Let R be such that $\mathbf{Q}_1\mathbf{R} = \mathbf{P}_1\mathbf{P}_2 = \mathbf{r}_2 - \mathbf{r}_1$.
- Then $\mathbf{P}_2\mathbf{R}$ is proportional to \mathbf{u}_1 .


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $|Q_1Q_2| = d(L_1, L_2)$.
- $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2 \Rightarrow \mathbf{Q}_1\mathbf{Q}_2$ is proportional to $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2$.
- Pick arbitrary points on L_1, L_2 - say, the base points $P_1(\mathbf{r}_1), P_2(\mathbf{r}_2)$.
- Let R be such that $\mathbf{Q}_1\mathbf{R} = \mathbf{P}_1\mathbf{P}_2 = \mathbf{r}_2 - \mathbf{r}_1$.
- Then $\mathbf{P}_2\mathbf{R}$ is proportional to \mathbf{u}_1 .
- $\Rightarrow \mathbf{Q}_2\mathbf{R} = \mathbf{Q}_2\mathbf{P}_2 + \mathbf{P}_2\mathbf{R}$ is perpendicular to \mathbf{n} .


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $|Q_1Q_2| = d(L_1, L_2)$.
- $\mathbf{Q}_1\mathbf{Q}_2 \perp L_1, L_2 \Rightarrow \mathbf{Q}_1\mathbf{Q}_2$ is proportional to $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2$.
- Pick arbitrary points on L_1, L_2 - say, the base points $P_1(\mathbf{r}_1), P_2(\mathbf{r}_2)$.
- Let R be such that $\mathbf{Q}_1\mathbf{R} = \mathbf{P}_1\mathbf{P}_2 = \mathbf{r}_2 - \mathbf{r}_1$.
- Then $\mathbf{P}_2\mathbf{R}$ is proportional to \mathbf{u}_1 .
- $\Rightarrow \mathbf{Q}_2\mathbf{R} = \mathbf{Q}_2\mathbf{P}_2 + \mathbf{P}_2\mathbf{R}$ is perpendicular to \mathbf{n} .
- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)$.


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

$\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$


Distance between non-parallel lines



- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)$.
- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)|$

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$

- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)| = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}|}{|\mathbf{n}|}$

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$

- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)| = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2)|}{|\mathbf{u}_1 \times \mathbf{u}_2|}$

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$

- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)| = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2)|}{|\mathbf{u}_1 \times \mathbf{u}_2|}$

- If lines are intersecting we know $d(L_1, L_2) = 0$.

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$

- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)| = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2)|}{|\mathbf{u}_1 \times \mathbf{u}_2|}$

- If lines are intersecting we know $d(L_1, L_2) = 0$. Since the lines intersect L_2 and L'_2 coincide.

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$

- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)| = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2)|}{|\mathbf{u}_1 \times \mathbf{u}_2|}$

- If lines are intersecting we know $d(L_1, L_2) = 0$. Since the lines intersect L_2 and L_2' coincide. $\Rightarrow (\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2) = 0$

Distance between non-parallel lines




- Given: lines $L_1 : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}_1$
 $L_2 : \mathbf{r} = \mathbf{r}_2 + s\mathbf{u}_2$
- The lines are skew or intersecting, i.e., $\mathbf{n} = \mathbf{u}_1 \times \mathbf{u}_2 \neq \mathbf{0}$.
- Goal: find distance between the lines $= d(L_1, L_2) =$ shortest distance b-n points on the two lines.

- $\Rightarrow \mathbf{Q}_1\mathbf{Q}_2 = \text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1).$

- $d(L_1, L_2) = |\text{proj}_{\mathbf{n}}(\mathbf{r}_2 - \mathbf{r}_1)| = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}|}{|\mathbf{n}|} = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2)|}{|\mathbf{u}_1 \times \mathbf{u}_2|}$


- If lines are intersecting we know $d(L_1, L_2) = 0$. Since the lines intersect L_2 and L'_2 coincide. $\Rightarrow (\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2) = 0 \Rightarrow$ the formula $d(L_1, L_2) = \frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot (\mathbf{u}_1 \times \mathbf{u}_2)|}{|\mathbf{u}_1 \times \mathbf{u}_2|} = 0$ produces the expected result.

Distance between parallel line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- The plane and the line are parallel,
- Goal: find distance between the the two.

Distance between parallel line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- The plane and the line are parallel, i.e. $\mathbf{u} \cdot \mathbf{n} = 0$.
- Goal: find distance between the two.


Distance between parallel line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- The plane and the line are parallel, i.e. $\mathbf{u} \cdot \mathbf{n} = 0$.
- Goal: find distance between the the two.

Distance from L to \mathcal{P} : $d(L, \mathcal{P}) = d(P_1, \mathcal{P})$

Distance between parallel line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- The plane and the line are parallel, i.e. $\mathbf{u} \cdot \mathbf{n} = 0$.
- Goal: find distance between the two.

Distance from L to \mathcal{P} : $d(L, \mathcal{P}) = d(P_1, \mathcal{P})$

$$d(L, \mathcal{P}) = |\text{orth}_{\mathbf{u}}(\mathbf{r}_1 - \mathbf{r}_0)| = \text{proj}_{\mathbf{n}}(\mathbf{r}_1 - \mathbf{r}_0)|$$

$$d(L, \mathcal{P}) = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \times \mathbf{u}|}{|\mathbf{u}|} = \frac{|(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}|}{|\mathbf{n}|}$$


Angle between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.

Line **perpendicular** to plane


Angle between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.

Line **perpendicular** to plane $\Leftrightarrow \mathbf{u} \parallel \mathbf{n}$


Angle between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.

Line **perpendicular** to plane $\Leftrightarrow \mathbf{u} \parallel \mathbf{n} \Leftrightarrow \boxed{\mathbf{u} \times \mathbf{n} = \mathbf{0}}$

Angle between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.

Line **perpendicular** to plane $\Leftrightarrow \mathbf{u} \parallel \mathbf{n} \Leftrightarrow \boxed{\mathbf{u} \times \mathbf{n} = \mathbf{0}}$

Angle between line and plane α : angle between L, \mathcal{P} .

$$\sin \alpha =$$

Angle between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.

Line **perpendicular** to plane $\Leftrightarrow \mathbf{u} \parallel \mathbf{n} \Leftrightarrow \boxed{\mathbf{u} \times \mathbf{n} = \mathbf{0}}$

Angle between line and plane α : angle between L, \mathcal{P} .

$$\sin \alpha = \frac{|\text{proj}_n \mathbf{u}|}{|\mathbf{u}|}$$

Angle between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.

Line **perpendicular** to plane $\Leftrightarrow \mathbf{u} \parallel \mathbf{n} \Leftrightarrow \boxed{\mathbf{u} \times \mathbf{n} = \mathbf{0}}$

Angle between line and plane α : angle between L, \mathcal{P} .

$$\sin \alpha = \frac{|\text{proj}_{\mathbf{n}} \mathbf{u}|}{|\mathbf{u}|} = \frac{|\mathbf{u} \cdot \mathbf{n}|}{|\mathbf{n}| |\mathbf{u}|}$$

Angle between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- plane $\mathcal{P} : (\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.
- Goal: Find/define angle between line and plane.


Line **perpendicular** to plane $\Leftrightarrow \mathbf{u} \parallel \mathbf{n} \Leftrightarrow \boxed{\mathbf{u} \times \mathbf{n} = \mathbf{0}}$

Angle between line and plane α : angle between L, \mathcal{P} .

$$\sin \alpha = \frac{|\text{proj}_{\mathbf{n}} \mathbf{u}|}{|\mathbf{u}|} = \frac{|\mathbf{u} \cdot \mathbf{n}|}{|\mathbf{n}| |\mathbf{u}|}$$


$$\alpha = \arcsin \left(\frac{|\mathbf{u} \cdot \mathbf{n}|}{|\mathbf{u}| |\mathbf{n}|} \right)$$

Intersection between line and plane



- Given: line L : $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane \mathcal{P} : $ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.


Intersection between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane.


Intersection between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.


Intersection between line and plane



- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.


Intersection between line and plane



- Given: line L : $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane \mathcal{P} : $ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$.


Intersection between line and plane



- Given: line L : $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane \mathcal{P} : $ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ **on the line** is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,

Intersection between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ on the line is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$, therefore P lies on both the line and the plane if:

$$(\mathbf{r}_1 + t\mathbf{u} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

Intersection between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ on the line is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$, therefore P lies on both the line and the plane if:

$$(\mathbf{r}_1 + t\mathbf{u} - \mathbf{r}_0) \cdot \mathbf{n} = 0$$

Intersection between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ on the line is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$, therefore P lies on both the line and the plane if:

$$\begin{aligned}(\mathbf{r}_1 + t\mathbf{u} - \mathbf{r}_0) \cdot \mathbf{n} &= 0 \\(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n} + t\mathbf{u} \cdot \mathbf{n} &= 0\end{aligned}$$

Intersection between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ on the line is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$, therefore P lies on both the line and the plane if:

$$\begin{aligned}
 (\mathbf{r}_1 + t\mathbf{u} - \mathbf{r}_0) \cdot \mathbf{n} &= 0 \\
 (\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n} + t\mathbf{u} \cdot \mathbf{n} &= 0 \\
 t &= -\frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}}{\mathbf{u} \cdot \mathbf{n}}
 \end{aligned}$$

Intersection between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ on the line is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$, therefore P lies on both the line and the plane if:

$$\begin{aligned}
 (\mathbf{r}_1 + t\mathbf{u} - \mathbf{r}_0) \cdot \mathbf{n} &= 0 \\
 (\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n} + t\mathbf{u} \cdot \mathbf{n} &= 0 \\
 t &= -\frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}}{\mathbf{u} \cdot \mathbf{n}} \\
 \mathbf{r} &= \mathbf{r}_1 - \frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}}{\mathbf{u} \cdot \mathbf{n}} \mathbf{u}
 \end{aligned}$$

Intersection between line and plane




- Given: line $L : \mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$,
- $\mathbf{r}_1 = (x_1, y_1, z_1)$, $\mathbf{u} = (p, q, r)$,
- plane $\mathcal{P} : ax + by + cz - d = 0$.
- Goal: find the intersection between line and plane.

Let $P_0(\mathbf{r}_0)$ be a point on the plane. Then a point $P(\mathbf{r})$, $\mathbf{r} = (x, y, z)$ is on the plane if $(\mathbf{r} - \mathbf{r}_0) \cdot \mathbf{n} = 0$. A point $P(\mathbf{r})$ on the line is of the form $\mathbf{r} = \mathbf{r}_1 + t\mathbf{u}$, therefore P lies on both the line and the plane if:

$$\begin{aligned}
 (\mathbf{r}_1 + t\mathbf{u} - \mathbf{r}_0) \cdot \mathbf{n} &= 0 \\
 (\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n} + t\mathbf{u} \cdot \mathbf{n} &= 0 \\
 t &= -\frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}}{\mathbf{u} \cdot \mathbf{n}} \\
 \mathbf{r} &= \mathbf{r}_1 - \frac{(\mathbf{r}_1 - \mathbf{r}_0) \cdot \mathbf{n}}{\mathbf{u} \cdot \mathbf{n}} \mathbf{u} \\
 &= (x_1, y_1, z_1) - \frac{ax_1 + by_1 + cz_1 - d}{ap + bq + cr} (p, q, r)
 \end{aligned}$$

Parallel planes



Planes are **parallel**


- Given: planes

$$P_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$$

$$P_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$$

- Goal: Establish whether planes are parallel, find distance b-n planes.

Parallel planes



- Given: planes


$$\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$$

$$\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$$

- Goal: Establish whether planes are parallel, find distance b-n planes.

Planes are **parallel** $\mathcal{P}_1 \parallel \mathcal{P}_2 \Leftrightarrow \mathbf{n}_1, \mathbf{n}_2 \text{ collinear} \Leftrightarrow \boxed{\mathbf{n}_1 \times \mathbf{n}_2 = \mathbf{0}}$.

Parallel planes



- Given: planes


$$\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$$

$$\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$$
- Goal: Establish whether planes are parallel, find distance b-n planes.

Planes are **parallel** $\mathcal{P}_1 \parallel \mathcal{P}_2 \Leftrightarrow \mathbf{n}_1, \mathbf{n}_2$ collinear $\Leftrightarrow \boxed{\mathbf{n}_1 \times \mathbf{n}_2 = \mathbf{0}}$.

Distance: $d(\mathcal{P}_1, \mathcal{P}_2) = |\text{proj}_{\mathbf{n}_1}(\mathbf{r}_2 - \mathbf{r}_1)|$

Parallel planes




- Given: planes
 $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$.
- Goal: Establish whether planes are parallel, find distance b-n planes.

Planes are **parallel** $\mathcal{P}_1 \parallel \mathcal{P}_2 \Leftrightarrow \mathbf{n}_1, \mathbf{n}_2$ collinear $\Leftrightarrow \boxed{\mathbf{n}_1 \times \mathbf{n}_2 = \mathbf{0}}$.

Distance: $d(\mathcal{P}_1, \mathcal{P}_2) = |\text{proj}_{\mathbf{n}_1}(\mathbf{r}_2 - \mathbf{r}_1)| = \boxed{\frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}_1|}{|\mathbf{n}_1|}}$

Parallel planes



- Given: planes

$$\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$$

$$\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$$


- Goal: Establish whether planes are parallel, find distance b-n planes.

Planes are **parallel** $\mathcal{P}_1 \parallel \mathcal{P}_2 \Leftrightarrow \mathbf{n}_1, \mathbf{n}_2$ collinear $\Leftrightarrow \boxed{\mathbf{n}_1 \times \mathbf{n}_2 = \mathbf{0}}$.

Distance: $d(\mathcal{P}_1, \mathcal{P}_2) = |\text{proj}_{\mathbf{n}_1}(\mathbf{r}_2 - \mathbf{r}_1)| = \boxed{\frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}_1|}{|\mathbf{n}_1|}}$

Assume $\mathbf{n}_1 = \mathbf{n}_2 = (a, b, c)$

Parallel planes




- Given: planes
 $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: Establish whether planes are parallel, find distance b-n planes.

Planes are **parallel** $\mathcal{P}_1 \parallel \mathcal{P}_2 \Leftrightarrow \mathbf{n}_1, \mathbf{n}_2$ collinear $\Leftrightarrow \boxed{\mathbf{n}_1 \times \mathbf{n}_2 = \mathbf{0}}$.

Distance: $d(\mathcal{P}_1, \mathcal{P}_2) = |\text{proj}_{\mathbf{n}_1}(\mathbf{r}_2 - \mathbf{r}_1)| = \boxed{\frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}_1|}{|\mathbf{n}_1|}}$

Assume $\mathbf{n}_1 = \mathbf{n}_2 = (a, b, c) \Rightarrow$ plane eq-ns: $\mathcal{P}_1 : ax + by + cz = d_1$
 $\mathcal{P}_2 : ax + by + cz = d_2$

Parallel planes



- Given: planes

$$\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$$

$$\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$$

- Goal: Establish whether planes are parallel, find distance b-n planes.


Planes are **parallel** $\mathcal{P}_1 \parallel \mathcal{P}_2 \Leftrightarrow \mathbf{n}_1, \mathbf{n}_2$ collinear $\Leftrightarrow \boxed{\mathbf{n}_1 \times \mathbf{n}_2 = \mathbf{0}}$.

Distance: $d(\mathcal{P}_1, \mathcal{P}_2) = |\text{proj}_{\mathbf{n}_1}(\mathbf{r}_2 - \mathbf{r}_1)| = \boxed{\frac{|(\mathbf{r}_2 - \mathbf{r}_1) \cdot \mathbf{n}_1|}{|\mathbf{n}_1|}}$

Assume $\mathbf{n}_1 = \mathbf{n}_2 = (a, b, c) \Rightarrow$ plane eq-ns: $\mathcal{P}_1 : ax + by + cz = d_1$
 $\mathcal{P}_2 : ax + by + cz = d_2$


$$\Rightarrow d(\mathcal{P}_1, \mathcal{P}_2) = \frac{|d_2 - d_1|}{\sqrt{a^2 + b^2 + c^2}}$$

Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.

Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Let L - intersection line of two planes.

Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Let L - intersection line of two planes.
- In \mathcal{P}_1 , drop perpendicular from arbitrary point Q_1 to L .

Angle between planes





- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: **define** and find the angle between the two planes.
- Let L - intersection line of two planes.
- In \mathcal{P}_1 , drop perpendicular from arbitrary point Q_1 to L .
- In \mathcal{P}_2 , raise a perpendicular from the perpendicular heel.

Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Let L - intersection line of two planes.
- In \mathcal{P}_1 , drop perpendicular from arbitrary point Q_1 to L .
- In \mathcal{P}_2 , raise a perpendicular from the perpendicular heel.
- Define angle α b-n $\mathcal{P}_1, \mathcal{P}_2$ = acute angle b-n two perpendiculars.

Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Let L - intersection line of two planes.
- In \mathcal{P}_1 , drop perpendicular from arbitrary point Q_1 to L .
- In \mathcal{P}_2 , raise a perpendicular from the perpendicular heel.
- Define angle α b-n $\mathcal{P}_1, \mathcal{P}_2$ = acute angle b-n two perpendiculars.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.


Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.




Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.
- \mathcal{P}_3 is orthogonal to L .




Angle between planes




- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.
- \mathcal{P}_3 is orthogonal to L .
- $\Rightarrow \mathcal{P}_3$ contains the normal vectors $\mathbf{n}_1, \mathbf{n}_2$.





Angle between planes



- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.
- \mathcal{P}_3 is orthogonal to L .
- $\Rightarrow \mathcal{P}_3$ contains the normal vectors $\mathbf{n}_1, \mathbf{n}_2$.
- $\mathbf{n}_1 \perp \mathbf{OQ}_1$ and $\mathbf{n}_2 \perp \mathbf{OQ}_2$.





Angle between planes





- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.
- \mathcal{P}_3 is orthogonal to L .
- $\Rightarrow \mathcal{P}_3$ contains the normal vectors $\mathbf{n}_1, \mathbf{n}_2$.
- $\mathbf{n}_1 \perp \mathbf{OQ}_1$ and $\mathbf{n}_2 \perp \mathbf{OQ}_2$.
- $\alpha = \text{acute} \angle(\mathbf{n}_1, \mathbf{n}_2)$
- $\alpha = \arccos \left(\frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{|\mathbf{n}_1| |\mathbf{n}_2|} \right)$

Angle between planes



- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.
- \mathcal{P}_3 is orthogonal to L .
- $\Rightarrow \mathcal{P}_3$ contains the normal vectors $\mathbf{n}_1, \mathbf{n}_2$.
- $\mathbf{n}_1 \perp \mathbf{OQ}_1$ and $\mathbf{n}_2 \perp \mathbf{OQ}_2$.
- $\alpha = \text{acute} \angle(\mathbf{n}_1, \mathbf{n}_2)$
- $\alpha = \arccos \left(\frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{|\mathbf{n}_1| |\mathbf{n}_2|} \right)$
- \perp planes: $\Rightarrow \alpha = \frac{\pi}{2} \iff \boxed{\mathbf{n}_1 \cdot \mathbf{n}_2 = 0}$.

Angle between planes



- Given: planes $\mathcal{P}_1 : (\mathbf{r} - \mathbf{r}_1) \cdot \mathbf{n}_1 = 0$
 $\mathcal{P}_2 : (\mathbf{r} - \mathbf{r}_2) \cdot \mathbf{n}_2 = 0$
- Goal: define and find the angle between the two planes.
- Consider the plane \mathcal{P}_3 spanned by the two constructed perpendiculars.
- \mathcal{P}_3 is orthogonal to L .
- $\Rightarrow \mathcal{P}_3$ contains the normal vectors $\mathbf{n}_1, \mathbf{n}_2$.
- $\mathbf{n}_1 \perp \mathbf{OQ}_1$ and $\mathbf{n}_2 \perp \mathbf{OQ}_2$.
- $\alpha = \text{acute} \angle(\mathbf{n}_1, \mathbf{n}_2)$
- $\alpha = \arccos \left(\frac{|\mathbf{n}_1 \cdot \mathbf{n}_2|}{|\mathbf{n}_1| |\mathbf{n}_2|} \right)$
- \perp planes: $\Rightarrow \alpha = \frac{\pi}{2} \iff \boxed{\mathbf{n}_1 \cdot \mathbf{n}_2 = 0}$.
- Direction of L is $\mathbf{u} = \mathbf{n}_1 \times \mathbf{n}_2$.