## Calculus I Homework Chain Rule Lecture 11

## 1. Compute the derivative using the chain rule.

(a) 
$$f(x) = \sqrt{1 + x^2}$$

(b) 
$$f(x) = \sqrt{3x^2 - x + 2}$$

(c) 
$$f(x) = \frac{x}{\sqrt{1 + \frac{2}{x^2}}}$$
.

(d) 
$$f(x) = \sqrt{1 - \sqrt{x}}$$
.

(e) 
$$y = (\cos x)^{\frac{1}{2}}$$

(f) 
$$f(x) = \sin^3 x$$
.

(g) 
$$y = (1 + \cos x)^2$$
.

(h) 
$$f(x) = \frac{1}{\sin^3 x}$$
.

(i) 
$$f(x) = \sqrt[3]{4 + 3\tan x}$$
.

(j) 
$$f(x) = (\cos x + 3\sin x)^4$$
.

(k) 
$$y = \sin\left(\sqrt{x}\right)$$

$$(1) \ y = \cos(4x)$$

## 2. Compute the derivative.

(a) 
$$f(x) = (x^4 + 3x^2 - 2)^5$$
.

(b) 
$$f(x) = (4x - x^2)^{100}$$
.

(c) 
$$f(x) = (2x-3)^4(x^2+x+1)^5$$
.

(d) 
$$f(x) = (x^2 + 1)^3 (x^2 + 2)^6$$
.

(e) 
$$f(x) = (3x-1)^4(2x+1)^{-3}$$
.

(f) 
$$f(x) = \frac{1}{1+x^2}$$
.

(g) 
$$f(x) = \left(\frac{x^2+1}{x^2-1}\right)^3$$
.

(h) 
$$f(x) = (x+1)^{\frac{2}{3}} (2x^2-1)^3$$
.

(i) 
$$f(x) = \sqrt{1 - 2x}$$
.

## 3. Differentiate.

(a) 
$$f(x) = \sin(\tan(2x))$$
.

(b) 
$$f(x) = \sec^2(mx)$$
.

(c) 
$$f(x) = \sec^2 x + \tan^2 x$$
.

(d) 
$$f(x) = x \sin\left(\frac{1}{x}\right)$$
.

(m) 
$$\sec^2(3x^2)$$
.

(n) 
$$\csc^2(3x^2)$$
.

(o) 
$$e^{2x}$$
.

(p) 
$$e^{-x^2}$$

(q) 
$$e^{\sqrt{x}}$$

(r) 
$$f(x) = e^{-\frac{1}{x}}$$
.

(s) 
$$5^x$$
.

(t) 
$$e^{2^x}$$
.

(u) 
$$2^{3^x}$$
.

(v) 
$$3^{2^x}$$
.

(w) 
$$y = \sqrt{\sec(4x)}$$

$$(x) y = x^2 \tan(5x)$$

(y) 
$$y = \frac{1 + \sin(x^2)}{1 + \cos(x^2)}$$
.

(j) 
$$f(x) = \sqrt{\frac{x^2 + 1}{x^2 + 4}}$$
.

(k) 
$$f(x) = 3\cot(2x)$$
.

(1) 
$$f(x) = \frac{1}{(1 + \sec x)^2}$$
.

(m) 
$$f(x) = \sqrt[3]{1 + \tan x}$$
.

(n) 
$$f(x) = \cos(2 + x^3)$$
.

(o) 
$$f(x) = \cos\left(\frac{1}{x}\right)\sin(x^2)$$
.

(p) 
$$f(x) = x \sec(kx)$$
.

(e) 
$$f(x) = \left(\frac{1 - \cos(2x)}{1 + \cos(2x)}\right)^4$$
.

(f) 
$$f(x) = \sqrt{\frac{x}{x^2 + 4}}$$
.

(g) 
$$f(t) = \cot^2(\sin t)$$
.

(h) 
$$f(x) = \left(ax + \sqrt{x^2 + b^2}\right)^{-2}$$
.

(i) 
$$f(x) = (x^2 + (1 - 3x)^5)^3$$
.

(j) 
$$f(x) = \sin(\sin(\sin x))$$
.

(k) 
$$f(x) = \sqrt{x + \sqrt{x}}$$
.

(1) 
$$f(x) = \sqrt{x + \sqrt{x + \sqrt{x}}}$$
.

(m) 
$$f(x) = (2r\sin(rx) + n)^p$$
.

(n) 
$$f(x) = \cos^4(\sin^3 x)$$
.

(o) 
$$f(x) = \cos \sqrt{\sin(\tan(\pi x))}$$
.

(p) 
$$f(x) = (x + (x + \sin^2 x)^3)^4$$
.

4. Compute the second derivative.

(a) 
$$f(x) = \sin(-5x)$$
.

(d) 
$$f(x) = e^{\frac{1}{x}}$$
.

(b) 
$$f(x) = \cot(2x)$$
.

(c) 
$$f(x) = e^{-3x}$$
.

(e) 
$$f(x) = e^{\sqrt{x}}$$
.

(f) 
$$f(x) = \frac{e^x - e^{-x}}{e^x + e^{-x}}$$

(g) 
$$f(x) = \frac{1}{2} \ln \left( \frac{1+x}{1-x} \right)$$