

# Precalculus

## Lecture 14

### Graphing Equations

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Graph of an equation

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:

<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

- Let  $F(x, y)$  and  $G(x, y)$  be arbitrary functions in the variables  $x, y$ .

- Let  $F(x, y)$  and  $G(x, y)$  be arbitrary functions in the variables  $x, y$ .
- An equation in  $x, y$  is an expression of the form

$$F(x, y) = G(x, y).$$

- Let  $F(x, y)$  and  $G(x, y)$  be arbitrary functions in the variables  $x, y$ .
- An equation in  $x, y$  is an expression of the form

$$F(x, y) = G(x, y).$$

- An ordered pair of numbers  $(a, b)$  is a solution of the equation if the number  $F(a, b)$  equals the number  $G(a, b)$ .

- Let  $F(x, y)$  and  $G(x, y)$  be arbitrary functions in the variables  $x, y$ .
- An equation in  $x, y$  is an expression of the form

$$F(x, y) = G(x, y).$$

- An ordered pair of numbers  $(a, b)$  is a solution of the equation if the number  $F(a, b)$  equals the number  $G(a, b)$ .

## Definition

The graph of the equation  $F(x, y) = G(x, y)$  is the set of all solutions  $(a, b)$  of the equation.

- Let  $F(x, y)$  and  $G(x, y)$  be arbitrary functions in the variables  $x, y$ .
- An equation in  $x, y$  is an expression of the form

$$F(x, y) = G(x, y).$$

- An ordered pair of numbers  $(a, b)$  is a solution of the equation if the number  $F(a, b)$  equals the number  $G(a, b)$ .

## Definition

The graph of the equation  $F(x, y) = G(x, y)$  is the set of all solutions  $(a, b)$  of the equation.

- Two equations are equivalent if they have the same graphs (set of solutions).



- Let  $F(x, y)$  and  $G(x, y)$  be arbitrary functions in the variables  $x, y$ .
- An equation in  $x, y$  is an expression of the form

$$F(x, y) = G(x, y).$$

- An ordered pair of numbers  $(a, b)$  is a solution of the equation if the number  $F(a, b)$  equals the number  $G(a, b)$ .

## Definition

The graph of the equation  $F(x, y) = G(x, y)$  is the set of all solutions  $(a, b)$  of the equation.

- Two equations are equivalent if they have the same graphs (set of solutions).
- If we set  $H(x, y) = F(x, y) - G(x, y)$ , we transform an arbitrary equation to an equivalent equation of the form:

$$H(x, y) = 0.$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5)$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5)$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5) = -21 + 20 = -1,$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5) = -21 + 20 = -1,$$

so  $(-3, -5)$  is a solution.

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5) = -21 + 20 = -1,$$

so  $(-3, -5)$  is a solution.

- For  $x = 3$ ,  $y = 5$ , we have:



## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5) = -21 + 20 = -1,$$

so  $(-3, -5)$  is a solution.

- For  $x = 3$ ,  $y = 5$ , we have:

$$(7x - 4y) = 7(3) - 4(5)$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5) = -21 + 20 = -1,$$

so  $(-3, -5)$  is a solution.

- For  $x = 3$ ,  $y = 5$ , we have:

$$(7x - 4y) = 7(3) - 4(5)$$

## Example

Determine which of the following points

- $(-3, -5)$
- $(3, 5)$

is a solution to the equation

$$7x - 4y = -1.$$

- For  $x = -3$ ,  $y = -5$ , we have:

$$7x - 4y = 7(-3) - 4(-5) = -21 + 20 = -1,$$

so  $(-3, -5)$  is a solution.

- For  $x = 3$ ,  $y = 5$ , we have:

$$(7x - 4y) = 7(3) - 4(5) = 21 - 20 = 1 \neq -1,$$

so  $(3, 5)$  is **not** a solution.

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4}$$



## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4}$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4}$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{\sqrt{2}}{4}\right)^2$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{\sqrt{2}}{4}\right)^2$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{\sqrt{2}}{4}\right)^2 = \frac{3}{4} + 2 \cdot \frac{2}{4^2}$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{\sqrt{2}}{4}\right)^2 = \frac{3}{4} + 2 \cdot \frac{2}{4^2}$$



## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{\sqrt{2}}{4}\right)^2 = \frac{3}{4} + \cancel{2} \cdot \frac{\cancel{2}}{\cancel{4}^2} = 1,$$

## Example

Determine which of the points  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$ ,  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution to the equation

$$x^2 + 2y^2 = 1.$$

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{1}{2}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{1}{2}\right)^2 = \frac{3}{4} + 2 \cdot \frac{1}{4} = \frac{5}{4} \neq 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$  is **not** a solution.

- For  $x = \frac{\sqrt{3}}{2}$ ,  $y = \frac{\sqrt{2}}{4}$ , we have:

$$x^2 + 2y^2 = \left(\frac{\sqrt{3}}{2}\right)^2 + 2\left(\frac{\sqrt{2}}{4}\right)^2 = \frac{3}{4} + \cancel{2} \cdot \frac{\cancel{2}}{4^{\cancel{2}}} = 1,$$

so  $\left(\frac{\sqrt{3}}{2}, \frac{\sqrt{2}}{4}\right)$  is a solution.

## Question

*Can we plot the graph of an arbitrary equation,  $F(x, y) = G(x, y)$ ?*

## Question

*Can we plot the graph of an arbitrary equation,  $F(x, y) = G(x, y)$ ?*

- For sufficiently well behaved equations the answer is yes.

## Question

*Can we plot the graph of an arbitrary equation,  $F(x, y) = G(x, y)$ ?*

- For sufficiently well behaved equations the answer is yes.
- We illustrate one computer algorithm for doing this.

## Question

*Can we plot the graph of an arbitrary equation,  $F(x, y) = G(x, y)$ ?*

- For sufficiently well behaved equations the answer is yes.
- We illustrate one computer algorithm for doing this.
- The theory behind plotting arbitrary equations, even when they are well behaved, is well beyond the scope of the present course.

## Question

*Can we plot the graph of an arbitrary equation,  $F(x, y) = G(x, y)$ ?*

- For sufficiently well behaved equations the answer is yes.
- We illustrate one computer algorithm for doing this.
- The theory behind plotting arbitrary equations, even when they are well behaved, is well beyond the scope of the present course.
- In particular, while computer algorithms plot graphs of well-behaved equations relatively well, it is not clear why those algorithms work.

## Question


*Can we plot the graph of an arbitrary equation,  $F(x, y) = G(x, y)$ ?*

- For sufficiently well behaved equations the answer is yes.
- We illustrate one computer algorithm for doing this.
- The theory behind plotting arbitrary equations, even when they are well behaved, is well beyond the scope of the present course.
- In particular, while computer algorithms plot graphs of well-behaved equations relatively well, it is not clear why those algorithms work.
- When, using algebra, we can express one variable in terms of the other, it is easy to produce the graph of the equation.




## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*



## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*

- Suppose  $y = f(x)$ .




Demonstration of the  
algorithm for

$$y = -\frac{1}{3}x + 1$$

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .

Demonstration of the  
algorithm for  
 $y = -\frac{1}{3}x + 1$   
from  $x = -9$  to  
 $x = 9$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .

Demonstration of the  
algorithm for  
 $y = -\frac{1}{3}x + 1$   
from  $x = -9$  to  
 $x = 9$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .

Demonstration of the  
algorithm for  
 $y = -\frac{1}{3}x + 1$   
from  $x = -9$  to  
 $x = 9$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.

Demonstration of the  
algorithm for  
 $y = -\frac{1}{3}x + 1$   
from  $x = -9$  to  
 $x = 9$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .

Demonstration of the algorithm for  
 $y = -\frac{1}{3}x + 1$   
 from  $x = -9$  to  
 $x = 9$ .



## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




Demonstration of the  
algorithm for  
 $y = -\frac{1}{3}x + 1$   
from  $x = -9$  to  
 $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .

Demonstration of the algorithm for  
 $y = -\frac{1}{3}x + 1$   
 from  $x = -9$  to  
 $x = 9$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.




Demonstration of the algorithm for  
 $y = -\frac{1}{3}x + 1$   
 from  $x = -9$  to  
 $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.




Demonstration of the algorithm for  $y = -\frac{1}{3}x + 1$  from  $x = -9$  to  $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.




Demonstration of the algorithm for  $y = -\frac{1}{3}x + 1$  from  $x = -9$  to  $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.




Demonstration of the algorithm for  $y = -\frac{1}{3}x + 1$  from  $x = -9$  to  $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.




Demonstration of the algorithm for  $y = -\frac{1}{3}x + 1$  from  $x = -9$  to  $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.



Demonstration of the algorithm for

$$y = \frac{1}{9}x^2 - \frac{1}{2}x + 4$$


from  $x = -9$  to  
 $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .



## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.



Demonstration of the algorithm for


$$y = \frac{1}{9}x^2 - \frac{1}{2}x + 4$$

from  $x = -9$  to  $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.



Demonstration of the algorithm for


$$y = \frac{1}{9}x^2 - \frac{1}{2}x + 4$$

from  $x = -9$  to  $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.



Demonstration of the algorithm for


$$y = -\frac{1}{9}x^2 - \frac{1}{2}x + 6$$

from  $x = -9$  to  
 $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.




Demonstration of the algorithm for  
 $y = -\frac{1}{9}x^2 - \frac{1}{2}x + 6$   
 from  $x = -9$  to  
 $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

## Question

*Can we plot the graph of an equation of the form  $y = f(x)$  or  $x = h(y)$  (for continuous  $h, f$ )? Yes.*




Demonstration of the algorithm for  
 $y = -\frac{1}{9}x^2 - \frac{1}{2}x + 6$   
 from  $x = -9$  to  
 $x = 9$ .

- Suppose  $y = f(x)$ .
- To plot the graph from  $x = a$  to  $x = b$ , select  $n + 1$  points  $x_0, x_1, \dots, x_n$  on  $[a, b]$ .
  - Usually we choose the points to be evenly spaced with  $x_0 = a$  and  $x_n = b$ .
  - $n + 1$  points split  $[a, b]$  into  $n$  intervals.
  - Each interval has length  $\Delta = \frac{b-a}{n}$ .
  - The formula for the  $j^{\text{th}}$  point is then  $x_j = a + \Delta \cdot j$ .
- The points  $(x_0, f(x_0)), \dots, (x_n, f(x_n))$  all lie on the graph.
- Connect them with straight lines.
- Repeat for increasing number of segments  $n$ .

You will not be tested on the material in the following slide.


## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ?*



## (Elementary Computer algorithm for sketching graphs)


Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? **Yes.**





## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*




We illustrate the  
algorithm for:

$$x^2 + 2y^2 = 1$$

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*




We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*

- Elementary algorithm: fix large rectangle.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.



We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .

We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$



## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  **have different sign** then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign **then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .**

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.

We illustrate the algorithm for:


$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$


Set  $H(x, y) = x^2 + 2y^2 - 1$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.



## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- **For each triangle:**
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$

$$x^2 + 2y^2 - 1 = 0$$


$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.



## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



We illustrate the algorithm for:

$$x^2 + 2y^2 = 1$$


$$x^2 + 2y^2 - 1 = 0$$

$$\text{Set } H(x, y) = x^2 + 2y^2 - 1$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:

$$y^2(y^2 - 3) = x^2(x^2 - 5)$$


$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.



## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:

$$y^2(y^2 - 3) = x^2(x^2 - 5)$$


$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.



## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.



Illustrate the algorithm for:


$$y^2(y^2 - 3) = x^2(x^2 - 5)$$

$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## (Elementary Computer algorithm for sketching graphs)

*Let  $H$ -continuous; is there simple algorithm to sketch  $H(x, y) = 0$ ? Yes.*



Illustrate the algorithm for:

$$y^2(y^2 - 3) = x^2(x^2 - 5)$$


$$H(x, y) = y^2(y^2 - 3) - x^2(x^2 - 5)$$

- Elementary algorithm: fix large rectangle.
- Split the grid in triangular mesh. One strategy to do that is shown.
- For each triangle:
  - Fix two corners  $P(x_P, y_P)$  and  $Q(x_Q, y_Q)$ .
  - If  $H(x_P, y_P)$  and  $H(x_Q, y_Q)$  have different sign then  $H$  must become zero somewhere on the segment between  $P$  and  $Q$ .
  - Select a point between  $P$  and  $Q$  and “guess” that  $H$  is zero there.
    - In our implementation, we select the midpoint (i.e.,  $\frac{1}{2}P + \frac{1}{2}Q$ ).
    - Connect the selected pts. for each triangle.
  - Repeat for ever finer grid.

## Definition

Points on the graph of  $F(x, y) = G(x, y)$  for which


- $x = 0$  are called  $y$ -intercepts (those lie on the  $y$  axis)



## Definition

Points on the graph of  $F(x, y) = G(x, y)$  for which


- $x = 0$  are called  $y$ -intercepts (those lie on the  $y$  axis)
- $y = 0$  are called  $x$ -intercept (those lie on the  $x$  axis).



## Definition

Points on the graph of  $F(x, y) = G(x, y)$  for which

- $x = 0$  are called  $y$ -intercepts (those lie on the  $y$  axis)
- $y = 0$  are called  $x$ -intercept (those lie on the  $x$  axis).




- To find the  $x$  intercepts set  $y = 0$  and solve for  $x$ .



## Definition

Points on the graph of  $F(x, y) = G(x, y)$  for which


- $x = 0$  are called  $y$ -intercepts (those lie on the  $y$  axis)
- $y = 0$  are called  $x$ -intercept (those lie on the  $x$  axis).




- To find the  $x$  intercepts set  $y = 0$  and solve for  $x$ .
- To find the  $y$  intercepts set  $x = 0$  and solve for  $y$ .

## Example

Find the  $x$  and  $y$  intercepts of the graph of the equation  $y = x^3 - x$ .



## Example



Find the  $x$  and  $y$  intercepts of the graph of the equation  $y = x^3 - x$ .

To find the  $y$  intercept, set  $x = 0$  to get  $y = 0$ . To find the  $x$  intercepts, set  $y = 0$  and solve

$$x^3 - x = 0$$

$$x(x^2 - 1) = 0$$


$$x(x - 1)(x + 1) = 0$$

The  $x$ -intercepts

$$x = 0 \text{ or } x = 1 \quad \text{or } x = -1.$$


are:  $(-1, 0)$ ,  $(0, 0)$ ,  $(1, 0)$ , the  $y$ -intercept is  $(0, 0)$ .

## Example



Find the  $x$  and  $y$  intercepts of the graph of the equation  $x^2 + 3x + y^2 = \frac{7}{4}$ .

## Example



Answer: the  
y-intercepts are:

$\left(0, \sqrt{\frac{7}{4}}\right),$   
 $\left(0, -\sqrt{\frac{7}{4}}\right)$ ; the x  
 intercepts are:  
 $\left(\frac{1}{2}, 0\right)$  and  $\left(-\frac{7}{2}, 0\right)$

Find the x and y intercepts of the graph of the equation  $x^2 + 3x + y^2 = \frac{7}{4}$ .

To find the y intercept, set  $x = 0$  and solve:

$$y^2 = \frac{7}{4} \Rightarrow y = \pm \sqrt{\frac{7}{4}}$$

To find the x intercepts, set  $y = 0$  and solve:

$$x^2 + 3x = \frac{7}{4}$$

$$x^2 + 3x - \frac{7}{4} = 0$$


$$4x^2 + 12x - 7 = 0$$

$$(2x - 1)(2x + 7) = 0$$

$$2x - 1 = 0 \quad \text{or} \quad 2x + 7 = 0$$

$$x = \frac{1}{2} \quad \text{or} \quad x = -\frac{7}{2}$$

- A graph is symmetric with respect to the  $x$  axis for each  $(x, y)$  lying on the graph  $(x, -y)$  also lies on the graph.
- A graph is symmetric with respect to the  $y$  axis for each  $(x, y)$  lying on the graph  $(-x, y)$  also lies on the graph.
- A graph is symmetric with respect to the origin if for each  $(x, y)$  lying on the graph  $(-x, -y)$  also lies on the graph.



### Observation

*The graph of the equation*

$$(x - h)^2 + (y - k)^2 = r^2$$

*is a circle with radius  $r$  and center  $(h, k)$ .*

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$ax^2 + bx + c$$



## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$ax^2 + bx + c = a \left( x^2 + \frac{b}{a}x \right) + c$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$ax^2 + bx + c = a \left( x^2 + \frac{b}{a}x \right) + c$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned} ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\ &= a \left( x^2 + \color{red}{2} \cdot \frac{b}{\color{red}{2}a}x \right) + c \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \quad \right)^2 - \left( \quad \right)^2 \right) + c \quad \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \quad \right)^2 \end{array} \right.
 \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned} ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\ &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\ &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \quad \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \end{array} \right. \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2\frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( \color{red}{x}^2 + 2\frac{b}{2a}\color{red}{x} + \left(\frac{b}{2a}\right)^2 - \left(\frac{b}{2a}\right)^2 \right) + c \\
 &= a \left( \left( \color{red}{x} + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left(\frac{b}{2a}\right)^2 \\ \text{use} \\ (\color{red}{A} + B)^2 = \\ \color{red}{A}^2 + 2\color{red}{A}B + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A + B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right.$$



## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c
 \end{aligned}
 \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A + B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right.$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c && \left. \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right\} \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 - a \cdot \frac{b^2}{4a^2} + c
 \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . To *complete the square* means to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2 \frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c \quad \left| \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right. \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 - \cancel{a} \cdot \frac{b^2}{\cancel{4a^2}} + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a}.
 \end{aligned}$$

## Definition (Completing the square)

Let  $a \neq 0$ . **To complete the square means** to carry out the following algebraic manipulation.

$$\begin{aligned}
 ax^2 + bx + c &= a \left( x^2 + \frac{b}{a}x \right) + c \\
 &= a \left( x^2 + 2 \cdot \frac{b}{2a}x \right) + c \\
 &= a \left( x^2 + 2\frac{b}{2a}x + \left( \frac{b}{2a} \right)^2 - \left( \frac{b}{2a} \right)^2 \right) + c && \left. \begin{array}{l} \text{Add \& subtract} \\ \left( \frac{b}{2a} \right)^2 \\ \text{use} \\ (A+B)^2 = \\ A^2 + 2AB + B^2 \end{array} \right\} \\
 &= a \left( \left( x + \frac{b}{2a} \right)^2 - \frac{b^2}{4a^2} \right) + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 - a \cdot \frac{b^2}{4a^2} + c \\
 &= a \left( x + \frac{b}{2a} \right)^2 + c - \frac{b^2}{4a}.
 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$3x^2 - 5x + 1$$

## Example (Completing the square)

Complete the square.

$$3x^2 - 5x + 1 = 3 \left( x^2 - ? x \right) + 1$$

## Example (Completing the square)

Complete the square.

$$3x^2 - 5x + 1 = 3 \left( x^2 - \frac{5}{3}x \right) + 1$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - \color{red}{2} \cdot \frac{5}{\color{red}{2} \cdot 3} x \right) + 1 \end{aligned}$$



## Example (Completing the square)

Complete the square.

$$\begin{aligned}
 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\
 &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\
 &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + ? \quad - ? \right) + 1
 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \textcolor{red}{?} - \textcolor{red}{?} \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \text{?} \quad - \text{?} \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - ? \right) + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned}3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\&= 3 \left( \left( x - \frac{5}{6} \right)^2 - ? \right) + 1\end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \end{aligned}$$



## Example (Completing the square)

Complete the square.

$$\begin{aligned}3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\&= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1\end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned} 3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\ &= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\ &= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\ &= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned}3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\&= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 + ?\end{aligned}$$

## Example (Completing the square)

Complete the square.

$$\begin{aligned}3x^2 - 5x + 1 &= 3 \left( x^2 - \frac{5}{3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{2 \cdot 3}x \right) + 1 \\&= 3 \left( x^2 - 2 \cdot \frac{5}{6}x + \left( \frac{5}{6} \right)^2 - \left( \frac{5}{6} \right)^2 \right) + 1 \\&= 3 \left( \left( x - \frac{5}{6} \right)^2 - \frac{25}{36} \right) + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 - \frac{25}{12} + 1 \\&= 3 \left( x - \frac{5}{6} \right)^2 - \frac{13}{12}.\end{aligned}$$

## Example

Show that the graph of the given equation is a circle. Find the center and radius of the circle.

- $x^2 + 2x + y^2 = 1.$
- $x^2 + x + 2y^2 + y = 1 + y^2.$
- $x^2 = 3x - y^2 - 2y.$
- $3x^2 + y = -3y^2.$
- $2x^2 + y = \frac{1}{2}x - 2y^2.$

## Example

Find an equation of a circle with center  $(2, 3)$  and passing through the point  $(-1, 1)$ .