

# Precalculus

## Lecture 21

Todor Milev

`https://github.com/tmilev/freecalc`

2020

# Outline

## 1 New Functions from Old Functions

# Outline

- 1 New Functions from Old Functions
- 2 Composing Functions with Linear Transformations

# Outline

- 1 New Functions from Old Functions
- 2 Composing Functions with Linear Transformations
- 3 Graphing Absolute Value of a Function

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) =$$

$$(f - g)(x) =$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) =$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$



# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = ?$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) =$$

$$\left(\frac{f}{g}\right)(x) =$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = ?$$

$$\left(\frac{f}{g}\right)(x) =$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) =$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = ?$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

for  $g(x) \neq 0$  .

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$(f + g)(x) = f(x) + g(x)$$

$$(f - g)(x) = f(x) - g(x)$$

$$(f \cdot g)(x) = f(x) \cdot g(x)$$

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)}$$

for  $g(x) \neq 0$  .

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ .



# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ .

$$\text{Dom}(f + g) = ?$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others.

$$\text{Dom}(f + g) = ?$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned}(f + g)(x) &= f(x) + g(x) \\(f - g)(x) &= f(x) - g(x) \\(f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 .\end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\text{Dom}(f + g) = ?$$

$$\text{Dom}(f - g) = ?$$

$$\text{Dom}(f \cdot g) = ?$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= ? \\ \text{Dom}(f \cdot g) &= ? \\ \text{Dom}\left(\frac{f}{g}\right) &= ? \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= ? \\ \text{Dom}(f \cdot g) &= ? \\ \text{Dom}\left(\frac{f}{g}\right) &= ? \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= ? \\ \text{Dom}(f \cdot g) &= ? \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = ?$$



# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \end{aligned} \quad \Bigg| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \end{array}$$

$$\text{Dom}\left(\frac{f}{g}\right) = \text{Dom}(f) \cap \text{Dom}(g) \cap \{x | g(x) \neq 0\}$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \left| \text{for } g(x) \neq 0 \right. \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x | g(x) \neq 0\} \end{aligned} \quad \left| \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \\ \\ \text{right expr.} \\ \text{stands for set} \\ \text{where } g(x) \neq 0 \end{array} \right.$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x | g(x) \neq 0\} \end{aligned} \quad \begin{array}{l} \Bigg| \text{ } \cap \text{ stands for} \\ \text{set intersection} \\ \\ \Bigg| \text{ right expr.} \\ \text{stands for set} \\ \text{where } g(x) \neq 0 \end{array}$$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore

$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \mid g(x) \neq 0\} \end{aligned}$$

$\cap$  stands for  
set intersection  
  
 right expr.  
stands for set  
where  $g(x) \neq 0$

# Combinations of Functions

Two functions  $f$  and  $g$  can be combined to form new functions  $f + g$ ,  $f - g$ ,  $f \cdot g$ , and  $\frac{f}{g}$ :

$$\begin{aligned} (f + g)(x) &= f(x) + g(x) \\ (f - g)(x) &= f(x) - g(x) \\ (f \cdot g)(x) &= f(x) \cdot g(x) \\ \left(\frac{f}{g}\right)(x) &= \frac{f(x)}{g(x)} \quad \Bigg| \text{ for } g(x) \neq 0 \end{aligned}$$

Let  $\text{Dom}(f)$  denote the domain of  $f$ . The function  $f + g$  is defined only if both  $f$  and  $g$  are defined, and similarly for the others. Therefore


$$\begin{aligned} \text{Dom}(f + g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f - g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}(f \cdot g) &= \text{Dom}(f) \cap \text{Dom}(g) \\ \text{Dom}\left(\frac{f}{g}\right) &= \text{Dom}(f) \cap \text{Dom}(g) \cap \{x \mid g(x) \neq 0\} \end{aligned} \quad \begin{array}{l} \cap \text{ stands for} \\ \text{set intersection} \\ \\ \text{right expr.} \\ \text{stands for set} \\ \text{where } g(x) \neq 0 \end{array}$$

## Definition (Composition of $f$ and $g$ )

If  $f$  and  $g$  are two functions, then the composition of  $f$  and  $g$  is written  $f \circ g$  and is defined by the formula

$$(f \circ g)(x) = f(g(x)).$$

Imagine  $f$  and  $g$  as machines taking some input and producing some output. Then  $f \circ g$  corresponds to attaching both machines end-to-end so that the output of  $g$  becomes the input of  $f$ .




## Definition (Composition of $f$ and $g$ )

If  $f$  and  $g$  are two functions, then the composition of  $f$  and  $g$  is written  $f \circ g$  and is defined by the formula

$$(f \circ g)(x) = f(g(x)).$$

Imagine  $f$  and  $g$  as machines taking some input and producing some output. Then  $f \circ g$  corresponds to attaching both machines end-to-end so that the output of  $g$  becomes the input of  $f$ .



The domain of  $f \circ g$  is the set of all numbers  $x$  in the domain of  $g$  such that  $g(x)$  is in the domain of  $f$ . If the domain of  $f$  is  $A$  and the domain of  $g$  is  $B$ , we write this as

$$\{x \in B \mid g(x) \in A\}.$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .



## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x)$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x))$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x})$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$



## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in \mathbb{R}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x)$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x))$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x})$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = \sqrt{3-\sqrt{x}}$$



## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3 - \sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3 - \sqrt{x} \geq 0$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3 - \sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3 - \sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3 - \sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3 - \sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3-x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3-\sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3-\sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

$$x \leq 9$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3 - \sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3 - \sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

$$x \leq 9$$

$$x \in ?$$



## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(f \circ g)(x) = f(g(x)) = f(\sqrt{3-x}) = \sqrt{\sqrt{3-x}} = \sqrt[4]{3-x}$$

Domain:

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$x \in (-\infty, 3].$$

$$(g \circ f)(x) = g(f(x)) = g(\sqrt{x}) = \sqrt{3 - \sqrt{x}}$$

Domain :

$$x \geq 0$$

$$3 - \sqrt{x} \geq 0$$

$$-\sqrt{x} \geq -3$$

$$\sqrt{x} \leq 3$$

$$x \leq 9$$

$$x \in [0, 9]$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x)$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x))$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x})$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$



## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

$$x \geq -6$$



## Example

Find  $f \circ g, g \circ f, g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

$$x \geq -6$$

$$x \in ?$$

## Example

Find  $f \circ g$ ,  $g \circ f$ ,  $g \circ g$  and their domains, where  $f(x) = \sqrt{x}$  and  $g(x) = \sqrt{3-x}$ .

$$(g \circ g)(x) = g(g(x)) = g(\sqrt{3-x}) = \sqrt{3 - \sqrt{3-x}}$$

Domain :

$$3 - x \geq 0$$

$$-x \geq -3$$

$$x \leq 3$$

$$3 - \sqrt{3-x} \geq 0$$

$$-\sqrt{3-x} \geq -3$$

$$\sqrt{3-x} \leq 3$$

$$3 - x \leq 9$$

$$-x \leq 6$$

$$x \geq -6$$

$$x \in [-6, 3].$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$x \neq ?$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$\begin{array}{l} f(x) = \frac{2x - 1}{x + 2} \\ g(x) = \frac{2x + 3}{5x - 7} \end{array} \quad \left| \quad x \neq -2 \right.$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq ?$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x))$$



## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right)$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right) = \frac{2\left(\frac{2x + 3}{5x - 7}\right) - 1}{\frac{2x + 3}{5x - 7} + 2}$$

## Example

Give simplified formulas for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right) = \frac{2\left(\frac{2x + 3}{5x - 7}\right) - 1}{\frac{2x + 3}{5x - 7} + 2}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} \end{aligned}$$



## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{\cancel{5x-7}}}{\frac{2x+3+(10x-14)}{\cancel{5x-7}}} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

$$x \neq ?$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$



## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x))$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right)$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq -2$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

# Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x - 1}{x + 2}$$

$$x \neq -2$$

$$g(x) = \frac{2x + 3}{5x - 7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x + 3}{5x - 7}\right) = \frac{2\left(\frac{2x + 3}{5x - 7}\right) - 1}{\frac{2x + 3}{5x - 7} + 2}$$

$$= \frac{\frac{2(2x + 3)}{5x - 7} - \frac{5x - 7}{5x - 7}}{\frac{2x + 3}{5x - 7} + \frac{2(5x - 7)}{5x - 7}} = \frac{\frac{4x + 6 - (5x - 7)}{5x - 7}}{\frac{2x + 3 + (10x - 14)}{5x - 7}} = \frac{-x + 13}{12x - 11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x - 1}{x + 2}\right) = \frac{2\left(\frac{2x - 1}{x + 2}\right) - 1}{\frac{2x - 1}{x + 2} + 2}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

**=?**

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$



# Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq ?$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

## Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$\begin{aligned} (f \circ g)(x) &= f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2} \\ &= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11} \end{aligned}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$\begin{aligned} (f \circ f)(x) &= f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2} \\ &= \frac{3x-4}{4x+3} \end{aligned}$$

$$x \neq -2, -\frac{3}{4}$$

# Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$

$$(g \circ f)(x) = ?$$

$$x \neq ?$$

$$(g \circ g)(x) = ?$$

$$x \neq ?$$

# Example

Give simplified f-las for  $f \circ g$ ,  $f \circ f$ ,  $g \circ f$ ,  $g \circ g$ . Find the implied domains.

$$f(x) = \frac{2x-1}{x+2}$$

$$x \neq -2$$

$$g(x) = \frac{2x+3}{5x-7}$$

$$x \neq \frac{7}{5}$$

$$(f \circ g)(x) = f(g(x)) = f\left(\frac{2x+3}{5x-7}\right) = \frac{2\left(\frac{2x+3}{5x-7}\right) - 1}{\frac{2x+3}{5x-7} + 2}$$

$$= \frac{\frac{2(2x+3)}{5x-7} - \frac{5x-7}{5x-7}}{\frac{2x+3}{5x-7} + \frac{2(5x-7)}{5x-7}} = \frac{\frac{4x+6-(5x-7)}{5x-7}}{\frac{2x+3+(10x-14)}{5x-7}} = \frac{-x+13}{12x-11}$$

$$x \neq \frac{11}{12}, \frac{7}{5}$$

$$(f \circ f)(x) = f(f(x)) = f\left(\frac{2x-1}{x+2}\right) = \frac{2\left(\frac{2x-1}{x+2}\right) - 1}{\frac{2x-1}{x+2} + 2}$$

$$= \frac{3x-4}{4x+3}$$

$$x \neq -2, -\frac{3}{4}$$


$$(g \circ f)(x) = \frac{7x+4}{3x-19}$$

$$x \neq -2, \frac{19}{3}$$

$$(g \circ g)(x) = \frac{19x-15}{-25x+64}$$


$$x \neq \frac{7}{5}, \frac{64}{25}$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

# Transformations of Functions




- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$f(x) + c$	
$f(x) - c$	
$f(x - c)$	
$f(x + c)$	



# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$


$$f(x) - c$$

$$f(x - c)$$

$$f(x + c)$$

Shift the graph of  $f(x)$   $c$  units up.

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

$$f(x) - c$$


$$f(x - c)$$

$$f(x + c)$$

Shift the graph of  $f(x)$   $c$  units up.

Shift the graph of  $f(x)$   $c$  units down.

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.


$$f(x) - c$$

Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$

Shift the graph of  $f(x)$   $c$  units down.


$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$



# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$

Shift the graph of  $f(x)$   $c$  units down.


$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$



# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$  ?

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$


Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$

Shift the graph of  $f(x)$   $c$  units right .

$$f(x + c)$$

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$

Shift the graph of  $f(x)$   $c$  units down.


$$f(x - c)$$

Shift the graph of  $f(x)$   $c$  units right .

$$f(x + c)$$

Shift the graph of  $f(x)$   $c$  units left.

# Transformations of Functions



- What happens to the graph if we add/subtract a positive constant  $c$  in the equation of a function  $f$ ?
- What happens if we add or subtract  $c$  from  $x$  before applying the function  $f$ ?

$$f(x) + c$$

Shift the graph of  $f(x)$   $c$  units up.

$$f(x) - c$$

Shift the graph of  $f(x)$   $c$  units down.

$$f(x - c)$$


Shift the graph of  $f(x)$   $c$  units right .

$$f(x + c)$$

Shift the graph of  $f(x)$   $c$  units left.


## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.



## Example


Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.



Complete the square:

## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$f(x) = x^2 + 6x + 10$$

## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + ?) + 10 - ? \end{aligned}$$



## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + 9) + 10 - 9 \end{aligned}$$

## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + 9) + 10 - 9 \\ &= (x + 3)^2 + 1 \end{aligned}$$

## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + 9) + 10 - 9 \\ &= (x + 3)^2 + 1 \\ &= (x - (-3))^2 + 1 \end{aligned}$$

## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + 9) + 10 - 9 \\ &= (x + 3)^2 + 1 \\ &= (x - (-3))^2 + 1 \end{aligned}$$

## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + 9) + 10 - 9 \\ &= (x + 3)^2 + 1 \\ &= (x - (-3))^2 + 1 \end{aligned}$$


## Example

Relative to the graph of  $f(x) = x^2$ , draw a graph of  $f(x) = x^2 + 6x + 10$ . Assume the graph of  $f(x) = x^2$  given.




Complete the square:

$$\begin{aligned} f(x) &= x^2 + 6x + 10 \\ &= (x^2 + 6x + 9) + 10 - 9 \\ &= (x + 3)^2 + 1 \\ &= (x - (-3))^2 + 1 \end{aligned}$$



- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?


$cf(x)$	
$\frac{1}{c}f(x)$	
$-f(x)$	
$f(-x)$	



- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?

$cf(x)$	
$\frac{1}{c}f(x)$	
$-f(x)$	
$f(-x)$	






- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?

 $cf(x)$  $\frac{1}{c}f(x)$  $-f(x)$  $f(-x)$ 

Stretch the graph of  $f(x)$  vertically by a factor of  $c$ .



- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?

$$cf(x)$$


$$\frac{1}{c}f(x)$$

$$-f(x)$$

$$f(-x)$$

Stretch the graph of  $f(x)$  vertically by a factor of  $c$ .

Compress the graph of  $f(x)$  vertically by a factor of  $c$ .



- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?

$$cf(x)$$

$$\frac{1}{c}f(x)$$


$$-f(x)$$

$$f(-x)$$

Stretch the graph of  $f(x)$  vertically by a factor of  $c$ .

Compress the graph of  $f(x)$  vertically by a factor of  $c$ .

Reflect the graph of  $f(x)$  in the  $x$ -axis.



- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?

$$cf(x)$$

Stretch the graph of  $f(x)$  vertically by a factor of  $c$ .

$$\frac{1}{c}f(x)$$


Compress the graph of  $f(x)$  vertically by a factor of  $c$ .

$$-f(x)$$

Reflect the graph of  $f(x)$  in the  $x$ -axis.


$$f(-x)$$

Reflect the graph of  $f(x)$  in the  $y$ -axis.




- What happens if we multiply or divide by a constant  $c > 1$  in the equation of a function  $f$ ?
- What happens if we multiply  $f$  by  $-1$ ?
- What happens if we multiply  $x$  by  $-1$  before applying  $f$ ?

$cf(x)$	Stretch the graph of $f(x)$ vertically by a factor of $c$ .
$\frac{1}{c}f(x)$	Compress the graph of $f(x)$ vertically by a factor of $c$ .
$-f(x)$	Reflect the graph of $f(x)$ in the $x$ -axis.
$f(-x)$	Reflect the graph of $f(x)$ in the $y$ -axis.




- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

$$f(cx)$$

$$f\left(\frac{1}{c}x\right)$$




- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 

?






- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 

?




- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 

?






- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f(\frac{1}{c}x)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$ 
 $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?


$f(cx)$	?
$f\left(\frac{1}{c}x\right)$	





- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?


$f(cx)$	?
$f\left(\frac{1}{c}x\right)$	



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

 $f(cx)$  $f\left(\frac{1}{c}x\right)$ 


?



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

$$f(cx)$$
$$f\left(\frac{1}{c}x\right)$$

Compress the graph of  $f(x)$  horizontally by a factor of  $c$ .



- What happens if we multiply  $x$  by const.  $c > 1$  before applying  $f$ ?
- What happens if we divide  $x$  by const.  $c > 1$  before applying  $f$ ?

$$f(cx)$$

$$f\left(\frac{1}{c}x\right)$$

Compress the graph of  $f(x)$  horizontally by a factor of  $c$ .

**Stretch the graph of  $f(x)$  horizontally by a factor of  $c$ .**

What happens when we take the absolute value of a function?

What happens when we take the absolute value of a function?

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

What happens when we take the absolute value of a function?

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

This tells us how to draw the graph of  $y = |f(x)|$ : the part of the graph above the  $x$ -axis remains the same; the part below the  $x$ -axis is reflected about the  $x$ -axis.


What happens when we take the absolute value of a function?

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

This tells us how to draw the graph of  $y = |f(x)|$ : the part of the graph above the  $x$ -axis remains the same; the part below the  $x$ -axis is reflected about the  $x$ -axis.

### Example

Draw the graph of the function  $f(x) = |x^2 - 1|$ .






What happens when we take the absolute value of a function?

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

This tells us how to draw the graph of  $y = |f(x)|$ : the part of the graph above the  $x$ -axis remains the same; the part below the  $x$ -axis is reflected about the  $x$ -axis.

### Example

Draw the graph of the function  $f(x) = |x^2 - 1|$ .



- Draw the graph of  $f(x) = x^2 - 1$ .


What happens when we take the absolute value of a function?

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

This tells us how to draw the graph of  $y = |f(x)|$ : the part of the graph above the  $x$ -axis remains the same; the part below the  $x$ -axis is reflected about the  $x$ -axis.

### Example

Draw the graph of the function  $f(x) = |x^2 - 1|$ .



- Draw the graph of  $f(x) = x^2 - 1$ .
- Identify the part(s) below the  $x$ -axis.


What happens when we take the absolute value of a function?

$$|f(x)| = \begin{cases} f(x) & \text{if } f(x) \geq 0 \\ -f(x) & \text{if } f(x) < 0 \end{cases}$$

This tells us how to draw the graph of  $y = |f(x)|$ : the part of the graph above the  $x$ -axis remains the same; the part below the  $x$ -axis is reflected about the  $x$ -axis.

### Example

Draw the graph of the function  $f(x) = |x^2 - 1|$ .



- Draw the graph of  $f(x) = x^2 - 1$ .
- Identify the part(s) below the  $x$ -axis.
- Flip those parts over the  $x$ -axis.