

# Precalculus

## Lecture 17

Todor Milev

`https://github.com/tmilev/freecalc`

2020

# Outline

- 1 Cartesian coordinate system
  - The Pythagorean Theorem, Euclidean Distance
  - Vectors
  - Segments, Midpoints

# License to use and redistribute


These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.


- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# Rectangular/Cartesian Coordinates




- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:

# Rectangular/Cartesian Coordinates




- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),

# Rectangular/Cartesian Coordinates




- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),
  - 2 pairwise perpendicular lines intersecting at the origin, called axes,

# Rectangular/Cartesian Coordinates



- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),
  - 2 pairwise perpendicular lines intersecting at the origin, called axes,
  - a direction in each of the coordinate axis.


# Rectangular/Cartesian Coordinates



- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),
  - 2 pairwise perpendicular lines intersecting at the origin, called axes,
  - a direction in each of the coordinate axis.
- The axes are labeled as  $x$ -axis and  $y$ -axis.




# Rectangular/Cartesian Coordinates




- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),
  - 2 pairwise perpendicular lines intersecting at the origin, called axes,
  - a direction in each of the coordinate axis.
- The axes are labeled as  $x$ -axis and  $y$ -axis.
- The  $x$  axis is drawn horizontal with direction pointing from left to right.

# Rectangular/Cartesian Coordinates




- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),
  - 2 pairwise perpendicular lines intersecting at the origin, called axes,
  - a direction in each of the coordinate axis.
- The axes are labeled as  $x$ -axis and  $y$ -axis.
- The  $x$  axis is drawn horizontal with direction pointing from left to right.
- The  $y$  axis is drawn vertical, pointing up.

# Rectangular/Cartesian Coordinates




- The Cartesian (rectangular) coordinate system is a way to represent points on the plane.
- To introduce Cartesian coordinates, fix:
  - a point  $O$  (called the origin),
  - 2 pairwise perpendicular lines intersecting at the origin, called axes,
  - a direction in each of the coordinate axis.
- The axes are labeled as  $x$ -axis and  $y$ -axis.
- The  $x$  axis is drawn horizontal with direction pointing from left to right.
- The  $y$  axis is drawn vertical, pointing up.
- The Cartesian coordinate system is named after René Descartes (1596-1650) (Latinized name: Cartesius).

# Rectangular/Cartesian Coordinates




- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .

# Rectangular/Cartesian Coordinates




- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.

# Rectangular/Cartesian Coordinates




- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q$  = base of perpendicular from  $P$  to  $x$ -axis.

# Rectangular/Cartesian Coordinates



- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q =$  base of perpendicular from  $P$  to  $x$ -axis.
- Define  $x$  as signed distance b-n  $O$  and  $Q$ .


# Rectangular/Cartesian Coordinates



- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q$  = base of perpendicular from  $P$  to  $x$ -axis.
- Define  $x$  as **signed distance b-n  $O$  and  $Q$** .
- Take distance with  **$+$  sign if  $OQ$  points in direction of  $x$ -axis,  $-$  sign else.**




# Rectangular/Cartesian Coordinates




- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q$  = base of perpendicular from  $P$  to  $x$ -axis.
- Define  $x$  as signed distance b-n  $O$  and  $Q$ .
- Take distance with  $+$  sign if  $OQ$  points in direction of  $x$ -axis,  $-$  sign else.
- To define  $y$ , do the same with the  $y$  axis.

# Rectangular/Cartesian Coordinates




- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q$  = base of perpendicular from  $P$  to  $x$ -axis.
- Define  $x$  as signed distance b-n  $O$  and  $Q$ .
- Take distance with  $+$  sign if  $OQ$  points in direction of  $x$ -axis,  $-$  sign else.
- To define  $y$ , do the same with the  $y$  axis.
- $(x, y)$  = Cartesian coordinates of  $P$ .

# Rectangular/Cartesian Coordinates




- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q$  = base of perpendicular from  $P$  to  $x$ -axis.
- Define  $x$  as signed distance b-n  $O$  and  $Q$ .
- Take distance with  $+$  sign if  $OQ$  points in direction of  $x$ -axis,  $-$  sign else.
- To define  $y$ , do the same with the  $y$  axis.
- $(x, y)$  = Cartesian coordinates of  $P$ .
- $x$  is called the  $x$ -coordinate of  $P$ ,  $y$ - the  $y$  coordinate.

# Rectangular/Cartesian Coordinates



- Let  $P$  -point. We assign to it a pair of numbers  $(x, y)$ .
- Distinct points are assigned distinct pairs.
- $Q$  = base of perpendicular from  $P$  to  $x$ -axis.
- Define  $x$  as signed distance b-n  $O$  and  $Q$ .
- Take distance with  $+$  sign if  $OQ$  points in direction of  $x$ -axis,  $-$  sign else.
- To define  $y$ , do the same with the  $y$  axis.
- $(x, y)$  = Cartesian coordinates of  $P$ .
- $x$  is called the  $x$ -coordinate of  $P$ ,  $y$ - the  $y$  coordinate.
- $(x, y)$  = signed lengths of sides of the rectangle indicated in the picture.


# Rectangular/Cartesian Coordinates



- The coordinate axes split the plane in 4 regions, called quadrants.
- The quadrants are labeled as indicated.
- For a point has coordinates  $(x, y)$ ,  $x \neq 0$ ,  $y \neq 0$ , the signs of  $x$  and  $y$  are determined by the quadrant that contains the point.

Quadrant	$(x, y)$
I	$(+, +)$
II	$(-, +)$
III	$(-, -)$
IV	$(+, -)$


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ .
- $(2, -3)$ .
- $(-3, 2)$ .
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ .
- $(2, -3)$ .
- $(-3, 2)$ .
- $(-1, -1)$ .

## Example




Plot the points and name the Quadrant that contains them

- (1, 2).
- (2, -3).
- (-3, 2).
- (-1, -1).




## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant ?
- $(2, -3)$ .
- $(-3, 2)$ .
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- (1, 2). Quadrant I
- (2, -3).
- (-3, 2).
- (-1, -1).


## Example



Plot the points and name the Quadrant that contains them

- (1, 2). Quadrant I
- (2, -3).
- (-3, 2).
- (-1, -1).


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ .
- $(-3, 2)$ .
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant ?
- $(-3, 2)$ .
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ .
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ .
- $(-1, -1)$ .

## Example




Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ .
- $(-1, -1)$ .




## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ . Quadrant ?
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ . Quadrant II
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ . Quadrant II
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ . Quadrant II
- $(-1, -1)$ .


## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ . Quadrant II
- $(-1, -1)$ . Quadrant ?

## Example



Plot the points and name the Quadrant that contains them

- $(1, 2)$ . Quadrant I
- $(2, -3)$ . Quadrant IV
- $(-3, 2)$ . Quadrant II
- $(-1, -1)$ . Quadrant III

- A triangle is a right-angled triangle if two of its sides are perpendicular.
- The two sides perpendicular to one another are called legs.
- The two legs form a right angle ( $90^\circ$ ).
- The side opposite to the right angle is called the hypotenuse.

## Theorem

*Let  $a, b$  be the lengths of the legs of a right-angled triangle and  $c$  the length of its hypotenuse. Then*

$$a^2 + b^2 = c^2.$$

## Theorem

*Let  $(x_1, y_1)$  and  $(x_2, y_2)$  be two points in the plane. Then the distance  $d$  between the two points is given by*

$$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$



## Example

Find the distance between  $(-2, 3)$  and  $(3, 5)$ .

$$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2} = \sqrt{(-2 - 3)^2 + (3 - 5)^2} = \sqrt{(-5)^2 + (-2)^2} = \sqrt{25 + 4} = \sqrt{29} \approx 5.385.$$

## Example

Find the distances between the indicated points.

$P$	$Q$	distance
$(2, 3)$	$(3, 5)$	?
$(-2, -3)$	$(3, 5)$	?
$(-2, -3)$	$(3, -5)$	?
$(-2, 3)$	$(3, -5)$	?

### Example

Do the points  $(1, 2)$ ,  $(2, 3)$ ,  $(4, -1)$  form a right-angled triangle?

### Example

Do the points  $(1, 2)$ ,  $(2, 4)$ ,  $(3, 1)$  form a right-angled triangle?

### Example


Do the indicated points form a right-angled triangle?

$(-1, -2)$     $(3, 5)$     $(6, -6)$    ?

$(1, 2)$     $(3, 5)$     $(6, 6)$    ?

$(0, 0)$     $(2, 3)$     $(3, -2)$    ?

$(0, 0)$     $(2, 3)$     $(-2, 3)$    ?




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v}$$


$$c \cdot \mathbf{u}$$



### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define


$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) \\ c \cdot \mathbf{u}\end{aligned}$$



### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define


$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) \\ c \cdot \mathbf{u} &\end{aligned}$$



### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. **Define**


$$\begin{array}{l} \mathbf{u} + \mathbf{v} \\ c \cdot \mathbf{u} \end{array} = \begin{array}{l} (x_1, y_1) + (x_2, y_2) \\ (x_1 + x_2, y_1 + y_2) \end{array}$$



### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. **Define**

$$\begin{array}{l} \mathbf{u} + \mathbf{v} \\ c \cdot \mathbf{u} \end{array} = \begin{array}{l} (x_1, y_1) + (x_2, y_2) \\ (x_1 + x_2, y_1 + y_2) \end{array}$$




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1)\end{aligned}$$






### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. **Define**

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. **Define**

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$  corresponds to a new point which we denote by  $P + Q$ .



### Definition ( $+$ , $\cdot$ in $\mathbb{R}^2$ )


Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$  corresponds to a new point which we denote by  $P + Q$ .
- One can show: the line through  $O$  and  $P$  is parallel to the line through  $Q$  and  $P + Q$ .






### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$  corresponds to a new point which we denote by  $P + Q$ .
- One can show: the line through  $O$  and  $P$  is parallel to the line through  $Q$  and  $P + Q$ .
- One can show: the line through  $O$  and  $Q$  is parallel to the line through  $P$  and  $P + Q$ .




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $(x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$  corresponds to a new point which we denote by  $P + Q$ .
- One can show: the line through  $O$  and  $P$  is parallel to the line through  $Q$  and  $P + Q$ .
- One can show: the line through  $O$  and  $Q$  is parallel to the line through  $P$  and  $P + Q$ .
- The points  $O, P, P + Q$  and  $Q$  form a parallelogram.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $c(x_1, y_1) = (cx_1, cy_1)$




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $c(x_1, y_1) = (cx_1, cy_1)$  corresponds to a new point which we denote by  $cP$ .




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.
- Let  $P, Q$  be points with respective coordinates  $(x_1, y_1)$  and  $(x_2, y_2)$ ; let  $O$  be the origin.
- $c(x_1, y_1) = (cx_1, cy_1)$  corresponds to a new point which we denote by  $cP$ .
- One can show  $O, P$  and  $cP$  lie on the same line.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.



### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$


Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.
- The correspondence between points in the plane and pairs of numbers depends on the choice of Cartesian coordinate system.






### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\mathbf{u} + \mathbf{v} = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2)$$

$$c \cdot \mathbf{u} = c \cdot (x_1, y_1) = (cx_1, cy_1).$$

- Fix a Cartesian coordinate system in the plane.
- The correspondence between points in the plane and pairs of numbers depends on the choice of Cartesian coordinate system.
- If we change the coordinate system we change  $+, \cdot$ .




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

- Fix a Cartesian coordinate system in the plane.
- The correspondence between points in the plane and pairs of numbers depends on the choice of Cartesian coordinate system.
- If we change the coordinate system we change  $+, \cdot$ .
- The points in the plane, equipped with the operations  $+, \cdot$  form a mathematical object which we call a vector space.




### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

### Definition (Translation)

Let  $P$  with coordinates  $(x, y)$  be a point and let  $(a, b)$  be a pair of numbers. The point  $P' = (x, y) + (a, b) = (x + a, y + b)$  is called the translation (shift) of  $P$   $a$  units right and  $b$  units up.



### Definition $(+, \cdot \text{ in } \mathbb{R}^2)$


Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

### Definition (Translation)

Let  $P$  with coordinates  $(x, y)$  be a point and let  $(a, b)$  be a pair of numbers. The point  $P' = (x, y) + (a, b) = (x + a, y + b)$  is called the translation (shift) of  $P$   $a$  units right and  $b$  units up.

- We allow shifts by negative units.



### Definition ( $+$ , $\cdot$ in $\mathbb{R}^2$ )


Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define

$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

### Definition (Translation)

Let  $P$  with coordinates  $(x, y)$  be a point and let  $(a, b)$  be a pair of numbers. The point  $P' = (x, y) + (a, b) = (x + a, y + b)$  is called the translation (shift) of  $P$   $a$  units right and  $b$  units up.

- We allow shifts by negative units.
- Translation down by  $b$  units we define to be translation up by  $-b$  units.



### Definition ( $+$ , $\cdot$ in $\mathbb{R}^2$ )

Let  $\mathbf{u} = (x_1, y_1)$  and  $\mathbf{v} = (x_2, y_2)$  be pairs of numbers and let  $c$  be a number. Define


$$\begin{aligned}\mathbf{u} + \mathbf{v} &= (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2) \\ c \cdot \mathbf{u} &= c \cdot (x_1, y_1) = (cx_1, cy_1).\end{aligned}$$

### Definition (Translation)

Let  $P$  with coordinates  $(x, y)$  be a point and let  $(a, b)$  be a pair of numbers. The point  $P' = (x, y) + (a, b) = (x + a, y + b)$  is called the translation (shift) of  $P$   $a$  units right and  $b$  units up.


- We allow shifts by negative units.
- Translation down by  $b$  units we define to be translation up by  $-b$  units.
- Translation left by  $a$  units we define to be translation right by  $-a$  units.

## Example



Translate  $(-3, 2)$  4 units right and 2 units down.


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.




## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (? , ? )$$


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2)$$


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2) = (-3 + 4, 2 + (-2))$$


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2) = (-3 + 4, 2 + (-2)) = (? , ?).$$


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2) = (-3 + 4, 2 + (-2)) = (\mathbf{1}, \mathbf{0}).$$


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2) = (-3 + 4, 2 + (-2)) = (1, 0).$$


## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2) = (-3 + 4, 2 + (-2)) = (1, 0).$$

## Example



Translate  $(-3, 2)$  4 units right and 2 units down.

$$(-3, 2) + (4, -2) = (-3 + 4, 2 + (-2)) = (1, 0).$$

## Example

Translate the point in the indicated way.

Point	Translation	result
$(2, 3)$	2 units left 3 units up	?
$(2, 1)$	2 units left $-2$ units down	?
$(-2, 1)$	$-1$ units right 2 units down	?
$(-2, 3)$	$-1$ units left 2 units up	?



## Observation

*The segment connecting  $P$  and  $Q$  consists of all points of the form*

$$tP + (1 - t)Q,$$

*where  $t$  runs over all numbers in the interval  $[0, 1]$ .*

- Let  $P$  have coordinates  $(x_1, y_1)$  and  $Q$  have coordinates  $(x_2, y_2)$ .
- Then the segment between  $P$  and  $Q$  consists of the points with coordinates

$$t(x_1, y_1) + (1 - t)(x_2, y_2).$$

## Observation

*The midpoint of the segment between  $P$  and  $Q$  is the point with  $t = \frac{1}{2}$ .*

$$\text{Midpoint}(P, Q) = \frac{1}{2}P + \frac{1}{2}Q = \left( \frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right).$$

## Example

Let  $P$  have coordinates  $(x_1, y_1)$  and  $Q$  have coordinates  $(x_2, y_2)$ . Let the midpoint of  $P$  and  $Q$  be  $R$ . Write the formula for the distance  $a$  between  $P$  and  $Q$ , and for the distance  $b$  between  $Q$  and  $R$ . Show that  $b = \frac{1}{2}a$ .

## Example

Find the midpoint of the indicated pairs of points.

$P$	$Q$	midpoint
$(1, 2)$	$(-1, -2)$	?
$(1, 2)$	$(1, -2)$	?
$(-1, 2)$	$(1, -2)$	?
$(-2, -3)$	$(3, 2)$	?