

# Calculus III

## Lecture 3

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

- 1 Cross product of vectors
  - Determinants
  - Cross product in coordinates

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.


- Latest version of the .tex sources of the slides:  
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
`https://creativecommons.org/licenses/by/3.0/us/`  
and the links therein.

# Torque




- If we tighten a bolt

# Torque




- If we tighten a bolt using a wrench,

# Torque




- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.

# Torque



- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:


# Torque



- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:
  - component  $\mathbf{F}_o$  orthogonal to the plane of rotation




# Torque




- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:
  - component  $\mathbf{F}_o$  orthogonal to the plane of rotation
  - component  $\mathbf{F}_\rho$  in the plane of rotation towards/away from the center

# Torque




- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:
  - component  $\mathbf{F}_o$  orthogonal to the plane of rotation
  - component  $\mathbf{F}_\rho$  in the plane of rotation towards/away from the center
  - component  $\mathbf{F}_\theta$  tangent to the motion of the wrench.

# Torque




- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:
  - component  $\mathbf{F}_o$  orthogonal to the plane of rotation
  - component  $\mathbf{F}_\rho$  in the plane of rotation towards/away from the center
  - component  $\mathbf{F}_\theta$  tangent to the motion of the wrench.
- Only  $\mathbf{F}_\theta$  contributes to the bolt motion.

# Torque




- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:
  - component  $\mathbf{F}_o$  orthogonal to the plane of rotation
  - component  $\mathbf{F}_\rho$  in the plane of rotation towards/away from the center
  - component  $\mathbf{F}_\theta$  tangent to the motion of the wrench.
- Only  $\mathbf{F}_\theta$  contributes to the bolt motion.
- The force of bolt motion  $\tau$  is proportional to length of wrench.

# Torque



- If we tighten a bolt using a wrench, it moves in direction perpendicular to the motion of the wrench.
- Let arm of the wrench: given by vector  $\mathbf{r}$ .
- Suppose we are applying a force  $\mathbf{F}$  at arm of the wrench. The force has three components:
  - component  $\mathbf{F}_o$  orthogonal to the plane of rotation
  - component  $\mathbf{F}_\rho$  in the plane of rotation towards/away from the center
  - component  $\mathbf{F}_\theta$  tangent to the motion of the wrench.
- Only  $\mathbf{F}_\theta$  contributes to the bolt motion.
- The force of bolt motion  $\tau$  is proportional to length of wrench.
- It turns out  $\tau = \mathbf{r} \times (\mathbf{F}_\rho + \mathbf{F}_\theta)$ , where  $\times$  is the vector cross product.

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are colinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}| |\text{orth}_{\mathbf{u}} \mathbf{v}| = |\mathbf{u}| |\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .



# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_u \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}} \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}} \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}} \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\mathbf{orth}_u \mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

# The Cross Product $\times$




## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\mathbf{orth}_\mathbf{u}\mathbf{v}| = |\mathbf{u}||\mathbf{v}|\sin\alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .



# The Cross Product $\times$



## Definition (Cross product)

$\mathbf{u} \times \mathbf{v}$  is the vector uniquely determined by the following.

- If  $\mathbf{u}, \mathbf{v}$  are non-zero and non-collinear.
  - $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .
  - The magnitude of  $\mathbf{u} \times \mathbf{v}$  equals  $|\mathbf{u}||\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{u}||\mathbf{v}| \sin \alpha$ .
  - The direction of  $\mathbf{u} \times \mathbf{v}$  is such that when viewed from the tip of  $\mathbf{u} \times \mathbf{v}$ ,  $\mathbf{v}$  is counter-clockwise from  $\mathbf{u}$ .
- If  $\mathbf{u}, \mathbf{v}$  are collinear or zero then  $\mathbf{u} \times \mathbf{v} = \mathbf{0}$ .

There are a couple of hand rules to help figure out the direction of the cross product.

# Properties of Cross Product

Let  $\mathbf{u}$ ,  $\mathbf{v}$  non-zero vectors,  $\alpha = \angle(\mathbf{u}, \mathbf{v})$ .

# Properties of Cross Product

Let  $\mathbf{u}$ ,  $\mathbf{v}$  non-zero vectors,  $\alpha = \angle(\mathbf{u}, \mathbf{v})$ .

- $|\mathbf{v} \times \mathbf{u}| = |\mathbf{u} \times \mathbf{v}|$ .

# Properties of Cross Product

Let  $\mathbf{u}$ ,  $\mathbf{v}$  non-zero vectors,  $\alpha = \angle(\mathbf{u}, \mathbf{v})$ .

- $|\mathbf{v} \times \mathbf{u}| = |\mathbf{u} \times \mathbf{v}|$ .

Indeed, that is because

$$|\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{v}| \sin \alpha \implies |\mathbf{u} \times \mathbf{v}| = |\mathbf{u}| |\mathbf{v}| \sin \alpha$$

# Properties of Cross Product

Let  $\mathbf{u}$ ,  $\mathbf{v}$  non-zero vectors,  $\alpha = \angle(\mathbf{u}, \mathbf{v})$ .

- $|\mathbf{v} \times \mathbf{u}| = |\mathbf{u} \times \mathbf{v}|$ .

Indeed, that is because

$$|\text{orth}_{\mathbf{u}} \mathbf{v}| = |\mathbf{v}| \sin \alpha \implies |\mathbf{u} \times \mathbf{v}| = |\mathbf{u}| |\mathbf{v}| \sin \alpha$$

- Cross product is anti-symmetric:

$$\mathbf{v} \times \mathbf{u} = -\mathbf{u} \times \mathbf{v}.$$

# Properties of Cross Product

Let  $\mathbf{u}, \mathbf{v}$  non-zero vectors,  $\alpha = \angle(\mathbf{u}, \mathbf{v})$ .

- $|\mathbf{v} \times \mathbf{u}| = |\mathbf{u} \times \mathbf{v}|$ .

Indeed, that is because

$$|\text{orth}_{\mathbf{u}}\mathbf{v}| = |\mathbf{v}| \sin \alpha \implies |\mathbf{u} \times \mathbf{v}| = |\mathbf{u}| |\mathbf{v}| \sin \alpha$$

- Cross product is anti-symmetric:

$$\mathbf{v} \times \mathbf{u} = -\mathbf{u} \times \mathbf{v}.$$

- Cross product is linear in each argument:

$$\mathbf{u} \times (a\mathbf{v} + b\mathbf{w}) = a\mathbf{u} \times \mathbf{v} + b\mathbf{u} \times \mathbf{w}$$

$$(a\mathbf{u} + b\mathbf{w}) \times \mathbf{v} = a\mathbf{u} \times \mathbf{v} + b\mathbf{w} \times \mathbf{v}$$

# $\text{orth}_u$ is a linear operator

## Theorem

$$\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_u\mathbf{v}_1 + \text{orth}_u\mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2)$$



# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

$$\begin{aligned}\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2))\end{aligned}$$

# $\text{orth}_u$ is a linear operator

## Theorem

$$\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

$$\begin{aligned} \text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_u(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_u(\mathbf{v}_1) + \text{proj}_u(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_u(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_u(\mathbf{v}_2)) \end{aligned}$$

# $\text{orth}_u$ is a linear operator

## Theorem

$$\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

$$\begin{aligned}\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_u(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_u(\mathbf{v}_1) + \text{proj}_u(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_u(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_u(\mathbf{v}_2))\end{aligned}$$

# $\text{orth}_u$ is a linear operator

## Theorem

$$\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

$$\begin{aligned} \text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_u(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_u(\mathbf{v}_1) + \text{proj}_u(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_u(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_u(\mathbf{v}_2)) \\ &= \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2 \end{aligned}$$

# $\text{orth}_u$ is a linear operator

## Theorem

$$\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2$$

## Proof.

Geometric proof:

Algebraic proof:

$$\begin{aligned} \text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_u(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_u(\mathbf{v}_1) + \text{proj}_u(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_u(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_u(\mathbf{v}_2)) \\ &= \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2 \end{aligned}$$


# $\text{orth}_u$ is a linear operator

## Theorem

$$\text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned} \text{orth}_u(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_u(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_u(\mathbf{v}_1) + \text{proj}_u(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_u(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_u(\mathbf{v}_2)) \\ &= \text{orth}_u \mathbf{v}_1 + \text{orth}_u \mathbf{v}_2 \end{aligned}$$


# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned}\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_{\mathbf{u}}(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2\end{aligned}$$

Let  $\mathcal{P}$ : plane  $\perp \mathbf{u}$ .


# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned}\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_{\mathbf{u}}(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2\end{aligned}$$

Let  $\mathcal{P}$ : plane  $\perp \mathbf{u}$ .




# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned} \text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_{\mathbf{u}}(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2 \end{aligned}$$

Let  $\mathcal{P}$ : plane  $\perp \mathbf{u}$ .


# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned} \text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_{\mathbf{u}}(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2 \end{aligned}$$

Let  $\mathcal{P}$ : plane  $\perp \mathbf{u}$ .


# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned} \text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_{\mathbf{u}}(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2 \end{aligned}$$

Let  $\mathcal{P}$ : plane  $\perp \mathbf{u}$ .


# $\text{orth}_{\mathbf{u}}$ is a linear operator

## Theorem

$$\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) = \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2$$

## Proof.

Geometric proof:



Algebraic proof:

$$\begin{aligned}\text{orth}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) &= (\mathbf{v}_1 + \mathbf{v}_2) - \text{proj}_{\mathbf{u}}(\mathbf{v}_1 + \mathbf{v}_2) \\ &= (\mathbf{v}_1 + \mathbf{v}_2) - (\text{proj}_{\mathbf{u}}(\mathbf{v}_1) + \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= (\mathbf{v}_1 - \text{proj}_{\mathbf{u}}(\mathbf{v}_1)) + (\mathbf{v}_2 - \text{proj}_{\mathbf{u}}(\mathbf{v}_2)) \\ &= \text{orth}_{\mathbf{u}}\mathbf{v}_1 + \text{orth}_{\mathbf{u}}\mathbf{v}_2\end{aligned}$$

Let  $\mathcal{P}$ : plane  $\perp \mathbf{u}$ .

# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

Geometric justification.




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\begin{aligned}\mathbf{u} \times \mathbf{v}_1 &= \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1 \\ \mathbf{u} \times \mathbf{v}_2 &= \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2 \\ \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2))\end{aligned}$$




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\begin{aligned} \mathbf{u} \times \mathbf{v}_1 &= \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1 \\ \mathbf{u} \times \mathbf{v}_2 &= \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2 \\ \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\begin{aligned} \mathbf{u} \times \mathbf{v}_1 &= \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1 \\ \mathbf{u} \times \mathbf{v}_2 &= \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2 \\ \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \\ &\Rightarrow \text{suffices to prove theorem when } \mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}. \end{aligned}$$






# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .

Since  $(a\mathbf{u}) \times \mathbf{v} = a(\mathbf{u} \times \mathbf{v}) \Rightarrow$  suffices to prove theorem when  $|\mathbf{u}| = 1$ .




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .

Since  $(a\mathbf{u}) \times \mathbf{v} = a(\mathbf{u} \times \mathbf{v}) \Rightarrow$  suffices to prove theorem when  $|\mathbf{u}| = 1$ .

When  $|\mathbf{u}| = 1$ , applying  $\mathbf{u} \times$  rotates all vectors in the plane  $\mathcal{P}$  at angle  $\frac{\pi}{2}$ .




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .

Since  $(a\mathbf{u}) \times \mathbf{v} = a(\mathbf{u} \times \mathbf{v}) \Rightarrow$  suffices to prove theorem when  $|\mathbf{u}| = 1$ .

When  $|\mathbf{u}| = 1$ , applying  $\mathbf{u} \times$  rotates all vectors in the plane  $\mathcal{P}$  at angle  $\frac{\pi}{2}$ .




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .

Since  $(a\mathbf{u}) \times \mathbf{v} = a(\mathbf{u} \times \mathbf{v}) \Rightarrow$  suffices to prove theorem when  $|\mathbf{u}| = 1$ .

When  $|\mathbf{u}| = 1$ , applying  $\mathbf{u} \times$  rotates all vectors in the plane  $\mathcal{P}$  at angle  $\frac{\pi}{2}$ .




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .

Since  $(a\mathbf{u}) \times \mathbf{v} = a(\mathbf{u} \times \mathbf{v}) \Rightarrow$  suffices to prove theorem when  $|\mathbf{u}| = 1$ .

When  $|\mathbf{u}| = 1$ , applying  $\mathbf{u} \times$  rotates all vectors in the plane  $\mathcal{P}$  at angle  $\frac{\pi}{2}$ .




# Justification of Linearity of $\times$ Product

## Theorem

$$\mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) = \mathbf{u} \times \mathbf{v}_1 + \mathbf{u} \times \mathbf{v}_2.$$

## Geometric justification.



$$\mathbf{u} \times \mathbf{v}_1 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_1$$

$$\mathbf{u} \times \mathbf{v}_2 = \mathbf{u} \times \text{orth}_{\mathbf{u}} \mathbf{v}_2$$

$$\begin{aligned} \mathbf{u} \times (\mathbf{v}_1 + \mathbf{v}_2) &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1 + \mathbf{v}_2)) \\ &= \mathbf{u} \times (\text{orth}_{\mathbf{u}} (\mathbf{v}_1) + \text{orth}_{\mathbf{u}} (\mathbf{v}_2)) \end{aligned}$$

$\Rightarrow$  suffices to prove theorem when  $\mathbf{v}_1, \mathbf{v}_2 \perp \mathbf{u}$ .

Since  $(a\mathbf{u}) \times \mathbf{v} = a(\mathbf{u} \times \mathbf{v}) \Rightarrow$  suffices to prove theorem when  $|\mathbf{u}| = 1$ .

When  $|\mathbf{u}| = 1$ , applying  $\mathbf{u} \times$  rotates all vectors in the plane  $\mathcal{P}$  at angle  $\frac{\pi}{2}$ . The statement of the theorem now follows from the fact that rotation preserves sums of vectors.



# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.



# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.
- Since  $\sigma$  - one to one,  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  have no repetition.

# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.
- Since  $\sigma$  - one to one,  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  have no repetition.

## Definition

A one-to-one function from the set  $\{1, 2, \dots, n\}$  to itself is called a permutation (“shuffling”).

# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.
- Since  $\sigma$  - one to one,  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  have no repetition.

## Definition

A one-to-one function from the set  $\{1, 2, \dots, n\}$  to itself is called a **permutation** (“shuffling”).

- There are  $n!$  different permutations:

# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.
- Since  $\sigma$  - one to one,  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  have no repetition.

## Definition

A one-to-one function from the set  $\{1, 2, \dots, n\}$  to itself is called a permutation (“shuffling”).

- There are  $n!$  different permutations:
  - there are  $n$  ways to select  $\sigma(1)$ ,

# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.
- Since  $\sigma$  - one to one,  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  have no repetition.

## Definition

A one-to-one function from the set  $\{1, 2, \dots, n\}$  to itself is called a permutation (“shuffling”).

- There are  $n!$  different permutations:
  - there are  $n$  ways to select  $\sigma(1)$ ,
  - $n - 1$  ways to select  $\sigma(2)$  (one number is already taken),

# Permutations and permutation signs

- Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be one to one function.
- Since  $\sigma$  - one to one,  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  have no repetition.

## Definition

A one-to-one function from the set  $\{1, 2, \dots, n\}$  to itself is called a permutation (“shuffling”).

- There are  $n!$  different permutations:
  - there are  $n$  ways to select  $\sigma(1)$ ,
  - $n - 1$  ways to select  $\sigma(2)$  (one number is already taken),
  - and so on, total:  $n \cdot (n - 1) \cdots 1 = n!$  ways to make a permutation.

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a **single swap of neighboring numbers**.

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are



# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are **transpositions** of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of **neighboring** numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are **not transpositions** of one another.

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are **not** transpositions of one another.
- Write the numbers  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  in a sequence.

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are **not** transpositions of one another.
- Write the numbers  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  in a sequence.
- Using transpositions, get from  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  to the properly ordered sequence  $1, 2, \dots, n$ .

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are **not** transpositions of one another.
- Write the numbers  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  in a sequence.
- Using transpositions, get from  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  to the properly ordered sequence  $1, 2, \dots, n$ .
- Number of transpositions used varies depending how we do it, but parity (even-ness) of # of transpositions is always the same.

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are **not** transpositions of one another.
- Write the numbers  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  in a sequence.
- Using transpositions, get from  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  to the properly ordered sequence  $1, 2, \dots, n$ .
- Number of transpositions used varies depending how we do it, but parity (even-ness) of # of transpositions is always the same.

## Definition

If we can get from  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  to  $(1, 2, \dots, n)$  with even # of transpositions, define  $\text{sign}(\sigma)$  to be 1, else define  $\text{sign}(\sigma)$  to be  $-1$ .

# Sign of permutation

- Given two sequences of numbers, define them to be transpositions of one another if one is obtained from the other with a single swap of neighboring numbers.
- $(2, 3, 4, 1)$  and  $(2, 4, 3, 1)$  are transpositions of one another.  
 $(2, 3, 4, 1)$  and  $(1, 3, 4, 2)$  are **not** transpositions of one another.
- Write the numbers  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  in a sequence.
- Using transpositions, get from  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  to the properly ordered sequence  $1, 2, \dots, n$ .
- Number of transpositions used varies depending how we do it, but parity (even-ness) of # of transpositions is always the same.
- If  $\text{sign}(\sigma) = 1$ ,  $\sigma$  is called even, if  $\text{sign}(\sigma) = -1$ ,  $\sigma$  is called odd.


## Definition

If we can get from  $(\sigma(1), \sigma(2), \dots, \sigma(n))$  to  $(1, 2, \dots, n)$  with even # of transpositions, define  $\text{sign}(\sigma)$  to be 1, else define  $\text{sign}(\sigma)$  to be  $-1$ .



- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots (n, \sigma(n))$ .

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board.



- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1))$ ,  $(2, \sigma(2)), \dots (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.

	1	2	3	4
1				
2				
3				
4				

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

	1	2	3	4
1				
2				
3				
4				

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

- Example:  $\sigma(1) = 2$ ,  $\sigma(2) = 3$ ,  $\sigma(3) = 4$ ,  $\sigma(4) = 1$ , pairs:
- $$\begin{array}{ll} (1, \sigma(1)) = (1, 2) \\ (2, \sigma(2)) = (2, 3) \\ (3, \sigma(3)) = (3, 4) \\ (4, \sigma(4)) = (4, 1) \end{array}$$


Corresponding peaceful rook placement:

	1	2	3	4
1	?	?	?	?
2				
3				
4				

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

Example:  $\sigma(1) = 2$ ,  $\sigma(2) = 3$ ,  $\sigma(3) = 4$ ,  $\sigma(4) = 1$ , pairs:  $(1, \sigma(1)) = (1, 2)$ ,  $(2, \sigma(2)) = (2, 3)$ ,  $(3, \sigma(3)) = (3, 4)$ ,  $(4, \sigma(4)) = (4, 1)$ .

Corresponding peaceful rook placement:



- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

$$\sigma(1) = 2 \qquad (1, \sigma(1)) = (1, 2)$$

- Example:  $\sigma(2) = 3$ , pairs:  $(2, \sigma(2)) = (2, 3)$

$$\sigma(3) = 4 \qquad (3, \sigma(3)) = (3, 4)$$

$$\sigma(4) = 1 \qquad (4, \sigma(4)) = (4, 1)$$

Corresponding peaceful rook placement:

	1	2	3	4
1		●		
2	?	?	?	?
3				
4				

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.


$$\sigma(1) = 2 \qquad (1, \sigma(1)) = (1, 2)$$

- Example:  $\sigma(2) = 3$ , pairs:  $(2, \sigma(2)) = (2, 3)$

$$\sigma(3) = 4 \qquad (3, \sigma(3)) = (3, 4)$$

$$\sigma(4) = 1 \qquad (4, \sigma(4)) = (4, 1)$$

Corresponding peaceful rook placement:





- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

- Example:  $\sigma(1) = 2$        $(1, \sigma(1)) = (1, 2)$   
 $\sigma(2) = 3$        $(2, \sigma(2)) = (2, 3)$   
 $\sigma(3) = 4$  , pairs:  $(3, \sigma(3)) = (3, 4)$  .  
 $\sigma(4) = 1$        $(4, \sigma(4)) = (4, 1)$


Corresponding peaceful rook placement:

	1	2	3	4
1		●		
2			●	
3	?	?	?	?
4				

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

- Example:  $\sigma(1) = 2$ ,  $\sigma(2) = 3$ ,  $\sigma(3) = 4$ ,  $\sigma(4) = 1$ , pairs:  $(1, \sigma(1)) = (1, 2)$ ,  $(2, \sigma(2)) = (2, 3)$ ,  $(3, \sigma(3)) = (3, 4)$ ,  $(4, \sigma(4)) = (4, 1)$ .

Corresponding peaceful rook placement:



- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

Example:  $\sigma(1) = 2$ ,  $\sigma(2) = 3$ ,  $\sigma(3) = 4$ ,  $\sigma(4) = 1$ , pairs:  $(1, \sigma(1)) = (1, 2)$ ,  $(2, \sigma(2)) = (2, 3)$ ,  $(3, \sigma(3)) = (3, 4)$ ,  $(4, \sigma(4)) = (4, 1)$ .


Corresponding peaceful rook placement:

	1	2	3	4
1		●		
2			●	
3				●
4	?	?	?	?

- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

Example:  $\sigma(1) = 2$ ,  $\sigma(2) = 3$ ,  $\sigma(3) = 4$ ,  $\sigma(4) = 1$ , pairs:  $(1, \sigma(1)) = (1, 2)$ ,  $(2, \sigma(2)) = (2, 3)$ ,  $(3, \sigma(3)) = (3, 4)$ ,  $(4, \sigma(4)) = (4, 1)$ .


Corresponding peaceful rook placement:



- To each permutation  $\sigma$ , assign  $n$  pairs of numbers  $(1, \sigma(1)), (2, \sigma(2)), \dots, (n, \sigma(n))$ .
- Consider a  $n \times n$  chess board. Interpret pair  $(k, \sigma(k))$  as *(row, column)*-coordinates in the board.
- For each pair  $(k, \sigma(k))$ , place a rook on the board.

- Example:  $\sigma(1) = 2, \sigma(2) = 3, \sigma(3) = 4, \sigma(4) = 1$  , pairs:  $(1, \sigma(1)) = (1, 2), (2, \sigma(2)) = (2, 3), (3, \sigma(3)) = (3, 4), (4, \sigma(4)) = (4, 1)$  .

Corresponding peaceful rook placement:



- $\sigma(k)$  are different  $\Rightarrow$  rook placements are peaceful: rooks never hit one another. i.e., no two points lie on same column or row.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$



# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

## First row

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

## Second row

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

$n^{th}$  row

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

## First column

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

## Second column

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.



# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

$n^{\text{th}}$  column

- Most common convention for matrix notation:
  - $(i, j)^{\text{th}}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.

# Square matrices

- Let  $A$  be  $n \times n$  (square) table of numbers.
- Technical term:  $A$  is a (square) *matrix*.
- Matrices are often denoted by surrounding with  $()$ -parenthesis:

$$A = \begin{pmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{pmatrix}.$$

- Most common convention for matrix notation:
  - $(i, j)^{th}$  entry of a matrix = denoted by letter with indices  $i, j$ , such as  $a_{ij}$
  - no comma between indices  $i, j$  in  $a_{ij}$
  - first index stands for row, second - for column.
- Non-square matrices: used & important but we discuss them elsewhere.

- The determinant  $\det A$  of a square matrix  $A$  is a number written as:

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{vmatrix}$$

- The determinant  $\det A$  of a square matrix  $A$  is a number written as:

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{vmatrix}$$

- The formula for the determinant is:

$$\det A = \sum_{\text{all permutations } \sigma} a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{n\sigma(n)} \text{sign}(\sigma) \quad .$$

- The determinant  $\det A$  of a square matrix  $A$  is a number written as:

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{vmatrix}$$

- The formula for the determinant is:

$$\det A = \sum_{\text{all permutations } \sigma} a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{n\sigma(n)} \text{sign}(\sigma) \quad .$$

- For every permutation  $\sigma$  we have one summand.

- The determinant  $\det A$  of a square matrix  $A$  is a number written as:

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{vmatrix}$$

- The formula for the determinant is:

$$\det A = \sum_{\text{all permutations } \sigma} a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{n\sigma(n)} \text{sign}(\sigma) \quad .$$

- For every permutation  $\sigma$  we have one summand.
- Every pair  $(k, \sigma(k))$  can be identified with a peaceful of a rook placement (as described in previous slides/lectures).

- The determinant  $\det A$  of a square matrix  $A$  is a number written as:

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{vmatrix}$$

- The formula for the determinant is:

$$\det A = \sum_{\text{all permutations } \sigma} a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{n\sigma(n)} \text{sign}(\sigma) \quad .$$

- For every permutation  $\sigma$  we have one summand.
- Every pair  $(k, \sigma(k))$  can be identified with a peaceful of a rook placement (as described in previous slides/lectures).
- For each rook placement we have a summand obtained by multiplying the numbers on which the rooks are standing.

- The determinant  $\det A$  of a square matrix  $A$  is a number written as:

$$\det A = \begin{vmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & & & & \\ a_{n1} & a_{n2} & a_{n3} & \dots & a_{nn} \end{vmatrix}$$

- The formula for the determinant is:

$$\det A = \sum_{\text{all permutations } \sigma} a_{1\sigma(1)} a_{2\sigma(2)} \dots a_{n\sigma(n)} \text{sign}(\sigma) \quad .$$

- For every permutation  $\sigma$  we have one summand.
- Every pair  $(k, \sigma(k))$  can be identified with a peaceful of a rook placement (as described in previous slides/lectures).
- For each rook placement we have a summand obtained by multiplying the numbers on which the rooks are standing.
- The sign of each summand is determined by the sign of the permutation.



## $2 \times 2$ determinants

$$\det \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} =$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 2$ .

## $2 \times 2$ determinants

$$\det \begin{vmatrix} \bullet & a_{12} \\ a_{21} & \bullet \end{vmatrix} =$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 2$ .
- There are two peaceful rook placements for a  $2 \times 2$  chessboard.

## $2 \times 2$ determinants

$$\det \begin{vmatrix} a_{11} & \bullet \\ \bullet & a_{22} \end{vmatrix} =$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 2$ .
- There are two peaceful rook placements for a  $2 \times 2$  chessboard.

## $2 \times 2$ determinants

$$\det \begin{vmatrix} \bullet & a_{12} \\ a_{21} & \bullet \end{vmatrix} = a_{11} a_{22}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 2$ .
- There are two peaceful rook placements for a  $2 \times 2$  chessboard.
- For each peaceful rook placement we got one summand.

## $2 \times 2$ determinants

$$\det \begin{vmatrix} a_{11} & \bullet \\ \bullet & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{12}a_{21}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 2$ .
- There are two peaceful rook placements for a  $2 \times 2$  chessboard.
- For each peaceful rook placement we got one summand.

## $2 \times 2$ determinants

$$\det \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{12}a_{21}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 2$ .
- There are two peaceful rook placements for a  $2 \times 2$  chessboard.
- For each peaceful rook placement we got one summand.
- The permutation  $(\sigma(1), \sigma(2)) = (2, 1)$  is odd, so one of the summands comes with negative sign.

## $3 \times 3$ determinants

$$\det \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} =$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .

## $3 \times 3$ determinants

$$\det \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} =$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.



## $3 \times 3$ determinants

$$\det \begin{vmatrix} \bullet & a_{12} & a_{13} \\ a_{21} & \bullet & a_{23} \\ a_{31} & a_{32} & \bullet \end{vmatrix} = a_{11} a_{22} a_{33}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.

# 3 × 3 determinants

$$\det \begin{vmatrix} a_{11} & \bullet & a_{13} \\ a_{21} & a_{22} & \bullet \\ \bullet & a_{32} & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.

# 3 × 3 determinants

$$\det \begin{vmatrix} a_{11} & a_{12} & \bullet \\ \bullet & a_{22} & a_{23} \\ a_{31} & \bullet & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.

# 3 × 3 determinants

$$\det \begin{vmatrix} a_{11} & a_{12} & \bullet \\ a_{21} & \bullet & a_{23} \\ \bullet & a_{32} & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.

# 3 × 3 determinants

$$\det \begin{vmatrix} \bullet & a_{12} & a_{13} \\ a_{21} & a_{22} & \bullet \\ a_{31} & \bullet & a_{33} \end{vmatrix} = a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32} - a_{13}a_{22}a_{31} - \textcolor{red}{a_{11}a_{23}a_{32}}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.

# 3 × 3 determinants

$$\det \begin{vmatrix} a_{11} & \bullet & a_{13} \\ \bullet & a_{22} & a_{23} \\ a_{31} & a_{32} & \bullet \end{vmatrix} = \begin{aligned} & a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} \\ & + a_{13}a_{21}a_{32} \\ & - a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} \\ & - \textcolor{red}{a_{12}a_{21}a_{33}} \end{aligned}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.

## $3 \times 3$ determinants

$$\det \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = \begin{aligned} & a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} \\ & + a_{13}a_{21}a_{32} \\ & - a_{13}a_{22}a_{31} - a_{11}a_{23}a_{32} \\ & - a_{12}a_{21}a_{33} \end{aligned}$$

- We specialize the  $n \times n$  determinant formula to the case  $n = 3$ .
- There are  $6 = 3!$  peaceful rook placements for a  $3 \times 3$  chessboard.
- For each peaceful rook placement we got one summand.
- The rook placements along the down-right “broken” diagonals correspond to even permutations, and the rook placements along the right-up “broken” diagonals correspond to negative permutations.

# Cross Product in Coordinates

- Let  $\mathbf{i}$ ,  $\mathbf{j}$ ,  $\mathbf{k}$ : unit vectors along coordinate axes.
- We have that

$$\begin{aligned}\mathbf{i} \times \mathbf{i} &= \mathbf{0}, & \mathbf{j} \times \mathbf{j} &= \mathbf{0}, & \mathbf{k} \times \mathbf{k} &= \mathbf{0} \\ \mathbf{i} \times \mathbf{j} &= \mathbf{k}, & \mathbf{j} \times \mathbf{k} &= \mathbf{i}, & \mathbf{k} \times \mathbf{i} &= \mathbf{j} \\ \mathbf{j} \times \mathbf{i} &= -\mathbf{k}, & \mathbf{k} \times \mathbf{j} &= -\mathbf{i}, & \mathbf{i} \times \mathbf{k} &= -\mathbf{j}\end{aligned}$$

- Let 
$$\begin{aligned}\mathbf{u} &= u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k} = (u_1, u_2, u_3) \\ \mathbf{v} &= v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k} = (v_1, v_2, v_3)\end{aligned}$$

- 

$$\begin{aligned}\mathbf{u} \times \mathbf{v} &= (u_1\mathbf{i} + u_2\mathbf{j} + u_3\mathbf{k}) \times (v_1\mathbf{i} + v_2\mathbf{j} + v_3\mathbf{k}) \\ &= (u_2v_3 - u_3v_2)\mathbf{i} + (u_3v_1 - u_1v_3)\mathbf{j} + (u_1v_2 - u_2v_1)\mathbf{k}\end{aligned}$$

$$\mathbf{u} \times \mathbf{v} = (u_1, u_2, u_3) \times (v_1, v_2, v_3) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$



$$\mathbf{u} \times \mathbf{v} = (u_1, u_2, u_3) \times (v_1, v_2, v_3) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

## Example

Find  $\mathbf{u} \times \mathbf{v}$ , where  $\mathbf{u} = (1, 2, 3)$  and  $\mathbf{v} = (6, 5, 4)$ .

$$\mathbf{u} \times \mathbf{v} = (u_1, u_2, u_3) \times (v_1, v_2, v_3) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

## Example

Find  $\mathbf{u} \times \mathbf{v}$ , where  $\mathbf{u} = (1, 2, 3)$  and  $\mathbf{v} = (6, 5, 4)$ .

$$\mathbf{u} \times \mathbf{v} = (1, 2, 3) \times (6, 5, 4) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 3 \\ 6 & 5 & 4 \end{vmatrix}$$

$$\mathbf{u} \times \mathbf{v} = (u_1, u_2, u_3) \times (v_1, v_2, v_3) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

## Example

Find  $\mathbf{u} \times \mathbf{v}$ , where  $\mathbf{u} = (1, 2, 3)$  and  $\mathbf{v} = (6, 5, 4)$ .

$$\begin{aligned} \mathbf{u} \times \mathbf{v} &= (1, 2, 3) \times (6, 5, 4) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 3 \\ 6 & 5 & 4 \end{vmatrix} \\ &= \begin{vmatrix} 2 & 3 \\ 5 & 4 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 3 \\ 6 & 4 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & 2 \\ 6 & 5 \end{vmatrix} \mathbf{k} \end{aligned}$$

$$\mathbf{u} \times \mathbf{v} = (u_1, u_2, u_3) \times (v_1, v_2, v_3) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

## Example

Find  $\mathbf{u} \times \mathbf{v}$ , where  $\mathbf{u} = (1, 2, 3)$  and  $\mathbf{v} = (6, 5, 4)$ .

$$\begin{aligned} \mathbf{u} \times \mathbf{v} &= (1, 2, 3) \times (6, 5, 4) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 3 \\ 6 & 5 & 4 \end{vmatrix} \\ &= \begin{vmatrix} 2 & 3 \\ 5 & 4 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 3 \\ 6 & 4 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & 2 \\ 6 & 5 \end{vmatrix} \mathbf{k} \\ &= (2 \cdot 4 - 3 \cdot 5) \mathbf{i} - (1 \cdot 4 - 3 \cdot 6) \mathbf{j} + (1 \cdot 5 - 2 \cdot 6) \mathbf{k} \end{aligned}$$

$$\mathbf{u} \times \mathbf{v} = (u_1, u_2, u_3) \times (v_1, v_2, v_3) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

## Example

Find  $\mathbf{u} \times \mathbf{v}$ , where  $\mathbf{u} = (1, 2, 3)$  and  $\mathbf{v} = (6, 5, 4)$ .

$$\begin{aligned} \mathbf{u} \times \mathbf{v} &= (1, 2, 3) \times (6, 5, 4) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 2 & 3 \\ 6 & 5 & 4 \end{vmatrix} \\ &= \begin{vmatrix} 2 & 3 \\ 5 & 4 \end{vmatrix} \mathbf{i} - \begin{vmatrix} 1 & 3 \\ 6 & 4 \end{vmatrix} \mathbf{j} + \begin{vmatrix} 1 & 2 \\ 6 & 5 \end{vmatrix} \mathbf{k} \\ &= (2 \cdot 4 - 3 \cdot 5) \mathbf{i} - (1 \cdot 4 - 3 \cdot 6) \mathbf{j} + (1 \cdot 5 - 2 \cdot 6) \mathbf{k} \\ &= -7\mathbf{i} + 14\mathbf{j} - 7\mathbf{k} = (-7, 14, -7). \end{aligned}$$

# Use $\times$ to find vector perpendicular to two given


Recall  $\mathbf{u} \times \mathbf{v}$  is perpendicular to  $\mathbf{u}$  and  $\mathbf{v}$ .

## Example

Find a vector  $\mathbf{w}$  perpendicular to  $\mathbf{u} = (1, 1, 0) = \mathbf{i} + \mathbf{j}$  and  $\mathbf{v} = \mathbf{j} + \mathbf{k} = (0, 1, 1)$ .


$$\begin{aligned}\mathbf{w} &= (\mathbf{i} + \mathbf{j}) \times (\mathbf{j} + \mathbf{k}) = \mathbf{i} \times \mathbf{j} + \mathbf{i} \times \mathbf{k} + \mathbf{j} \times \mathbf{j} + \mathbf{j} \times \mathbf{k} = \\ &= \mathbf{k} - \mathbf{j} + \mathbf{0} + \mathbf{i} = \mathbf{i} - \mathbf{j} + \mathbf{k} = (1, -1, 1) .\end{aligned}$$

# Use $\times$ to find area of triangle in space



- $A, B, C$  points in space,  $\mathbf{u} = \mathbf{AB}$ ,  $\mathbf{v} = \mathbf{AC}$ .


# Use $\times$ to find area of triangle in space



- $A, B, C$  points in space,  $\mathbf{u} = \mathbf{AB}$ ,  $\mathbf{v} = \mathbf{AC}$ .
- Then  
 $|\mathbf{w}| = |\mathbf{orth}_{\mathbf{u}}\mathbf{v}| = \text{distance from } C \text{ to } AB.$




# Use $\times$ to find area of triangle in space




- $A, B, C$  points in space,  $\mathbf{u} = \mathbf{AB}$ ,  $\mathbf{v} = \mathbf{AC}$ .
- Then
$$|\mathbf{w}| = |\mathbf{orth}_u \mathbf{v}| = \text{distance from } C \text{ to } AB.$$
- $|\mathbf{u} \times \mathbf{v}| = |\mathbf{orth}_u \mathbf{v}| |\mathbf{u}| = 2\text{area}(ABC) = \text{area}(ABDC)$

# Use $\times$ to find area of triangle in space



- $A, B, C$  points in space,  $\mathbf{u} = \mathbf{AB}$ ,  $\mathbf{v} = \mathbf{AC}$ .
- Then
  - $|\mathbf{w}| = |\mathbf{orth}_{\mathbf{u}}\mathbf{v}| = \text{distance from } C \text{ to } AB.$
- $|\mathbf{u} \times \mathbf{v}| = |\mathbf{orth}_{\mathbf{u}}\mathbf{v}| |\mathbf{u}| = 2\text{area}(ABC) = \text{area}(ABDC)$
- $|\mathbf{u} \times \mathbf{v}| = \text{Area of parallelogram on sides } \mathbf{u} \text{ and } \mathbf{v}.$

## Example




Find the area of the triangle  $A(1, 2, 3)$ ,  $B(2, 3, 1)$ ,  $C(3, 1, 2)$ .


$$\begin{aligned}\text{Area}(ABC) &= \frac{1}{2}|\mathbf{AB} \times \mathbf{AC}| = \frac{1}{2}|(1, 1, -2) \times (2, -1, -1)| \\ &= \frac{1}{2}|(-3, -3, -3)| \\ &= \frac{3\sqrt{3}}{2}.\end{aligned}$$

# Scalar Triple Product

- $A, B, C, D$  points in space;




# Scalar Triple Product




- $A, B, C, D$  points in space;
- $\mathbf{u} = \mathbf{AB}, \mathbf{v} = \mathbf{AC}, \mathbf{w} = \mathbf{AD}$ ;

# Scalar Triple Product




- $A, B, C, D$  points in space;
- $\mathbf{u} = \mathbf{AB}, \mathbf{v} = \mathbf{AC}, \mathbf{w} = \mathbf{AD}$ ;
- $R = R(\mathbf{u}, \mathbf{v}, \mathbf{w})$ : box on sides  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ .

# Scalar Triple Product



- $A, B, C, D$  points in space;
- $\mathbf{u} = \mathbf{AB}, \mathbf{v} = \mathbf{AC}, \mathbf{w} = \mathbf{AD}$ ;
- $R = R(\mathbf{u}, \mathbf{v}, \mathbf{w})$ : box on sides  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ .
- $\text{Vol}(R) = |\mathbf{u} \times \mathbf{v}| |\mathbf{r}| = |\mathbf{u} \times \mathbf{v}| |\text{proj}_{\mathbf{u} \times \mathbf{v}} \mathbf{w}| = |\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})|$ .

# Scalar Triple Product




- $A, B, C, D$  points in space;
- $\mathbf{u} = \mathbf{AB}, \mathbf{v} = \mathbf{AC}, \mathbf{w} = \mathbf{AD}$ ;
- $R = R(\mathbf{u}, \mathbf{v}, \mathbf{w})$ : box on sides  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ .
- $\text{Vol}(R) = |\mathbf{u} \times \mathbf{v}| |\mathbf{r}| = |\mathbf{u} \times \mathbf{v}| |\text{proj}_{\mathbf{u} \times \mathbf{v}} \mathbf{w}| = |\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})|$ .

## Definition

The quantity  $\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})$  is called the scalar triple product of  $\mathbf{w}, \mathbf{u}, \mathbf{v}$ .



# Scalar Triple Product



- $A, B, C, D$  points in space;
- $\mathbf{u} = \mathbf{AB}$ ,  $\mathbf{v} = \mathbf{AC}$ ,  $\mathbf{w} = \mathbf{AD}$ ;
- $R = R(\mathbf{u}, \mathbf{v}, \mathbf{w})$ : box on sides  $\mathbf{u}$ ,  $\mathbf{v}$ ,  $\mathbf{w}$ .
- $\text{Vol}(R) = |\mathbf{u} \times \mathbf{v}| |\mathbf{r}| = |\mathbf{u} \times \mathbf{v}| |\text{proj}_{\mathbf{u} \times \mathbf{v}} \mathbf{w}| = |\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})|$ .


## Definition

The quantity  $\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v})$  is called the scalar triple product of  $\mathbf{w}$ ,  $\mathbf{u}$ ,  $\mathbf{v}$ .

- If  $\mathbf{u} = (u_1, u_2, u_3)$ ,  $\mathbf{v} = (v_1, v_2, v_3)$ , and  $\mathbf{w} = (w_1, w_2, w_3)$ , then


$$\mathbf{w} \cdot (\mathbf{u} \times \mathbf{v}) = \begin{vmatrix} w_1 & w_2 & w_3 \\ u_1 & u_2 & u_3 \\ v_1 & v_2 & v_3 \end{vmatrix}$$

## Example




Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

## Example




Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

## Example




Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

## Example




Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

## Example




Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

## Example



Find the volume of the **parallelepiped (slanted box)** with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

## Example




Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = \textcolor{red}{?} + ? + ? - ? - ? - ?$$




## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + ? + ? - ? - ? - ?$$


## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + 1 + ? - ? - ? - ?$$


## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + 1 + 1 - ? - ? - ?$$


## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + 1 + 1 - (-1) - ? - ?$$


## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + 1 + 1 - (-1) - (-1) - ?$$


## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + 1 + 1 - (-1) - (-1) - (-1)$$


## Example



Find the volume of the parallelepiped (slanted box) with vertex at the origin spanned by the vectors  $(-1, 1, 1)$ ,  $(1, -1, 1)$ ,  $(1, 1, -1)$ .

$$\begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = -1 + 1 + 1 - (-1) - (-1) - (-1) \\ = 4$$

## Example




Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$\text{Vol}(\text{tetrahedron}) =$




## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\text{Vol}(\text{tetrahedron}) =$$


## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$\text{Vol}(\text{tetrahedron}) =$


## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\text{Vol}(\text{tetrahedron}) =$$


## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$\text{Vol}(\text{tetrahedron}) =$


## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\text{Vol}(\text{tetrahedron}) =$$


## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$\text{Vol}(\text{tetrahedron}) =$


## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\text{Vol}(\text{tetrahedron}) =$$

## Example




Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\text{Vol}(\text{tetrahedron}) = \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges})$$



## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .


$$u_1 = ?$$

$$u_2 = ?$$

$$u_3 = ?$$

$$\text{Vol}(\text{tetrahedron}) = \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges})$$

## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .


$$\mathbf{u}_1 = (1, -1, -1) - (1, 1, 1) = (0, -2, -2)$$

$$\mathbf{u}_2 = (-1, 1, -1) - (1, 1, 1) = (-2, 0, -2)$$

$$\mathbf{u}_3 = (-1, -1, 1) - (1, 1, 1) = (-2, -2, 0)$$

$$\text{Vol}(\text{tetrahedron}) = \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges})$$

## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .


$$\mathbf{u}_1 = (1, -1, -1) - (1, 1, 1) = (0, -2, -2)$$

$$\mathbf{u}_2 = (-1, 1, -1) - (1, 1, 1) = (-2, 0, -2)$$

$$\mathbf{u}_3 = (-1, -1, 1) - (1, 1, 1) = (-2, -2, 0)$$

$$\begin{aligned} \text{Vol}(\text{tetrahedron}) &= \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges}) \\ &= \frac{1}{6} \left| \det \begin{pmatrix} 0 & -2 & -2 \\ -2 & 0 & -2 \\ -2 & -2 & 0 \end{pmatrix} \right| \end{aligned}$$

## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\mathbf{u}_1 = (1, -1, -1) - (1, 1, 1) = (0, -2, -2)$$


$$\mathbf{u}_2 = (-1, 1, -1) - (1, 1, 1) = (-2, 0, -2)$$

$$\mathbf{u}_3 = (-1, -1, 1) - (1, 1, 1) = (-2, -2, 0)$$

$$\text{Vol}(\text{tetrahedron}) = \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges})$$

$$= \frac{1}{6} \left| \det \begin{pmatrix} 0 & -2 & -2 \\ -2 & 0 & -2 \\ -2 & -2 & 0 \end{pmatrix} \right| = \frac{1}{6} | \text{?} |$$

## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .


$$\mathbf{u}_1 = (1, -1, -1) - (1, 1, 1) = (0, -2, -2)$$

$$\mathbf{u}_2 = (-1, 1, -1) - (1, 1, 1) = (-2, 0, -2)$$

$$\mathbf{u}_3 = (-1, -1, 1) - (1, 1, 1) = (-2, -2, 0)$$

$$\begin{aligned} \text{Vol}(\text{tetrahedron}) &= \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges}) \\ &= \frac{1}{6} \left| \det \begin{pmatrix} 0 & -2 & -2 \\ -2 & 0 & -2 \\ -2 & -2 & 0 \end{pmatrix} \right| = \frac{1}{6} | -16 | \end{aligned}$$

## Example



Find the volume of the tetrahedron with vertices  $(1, 1, 1)$ ,  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ .

$$\mathbf{u}_1 = (1, -1, -1) - (1, 1, 1) = (0, -2, -2)$$

$$\mathbf{u}_2 = (-1, 1, -1) - (1, 1, 1) = (-2, 0, -2)$$

$$\mathbf{u}_3 = (-1, -1, 1) - (1, 1, 1) = (-2, -2, 0)$$

$$\begin{aligned} \text{Vol}(\text{tetrahedron}) &= \frac{1}{6} \text{Vol}(\text{Box generated by any 3 edges}) \\ &= \frac{1}{6} \left| \det \begin{pmatrix} 0 & -2 & -2 \\ -2 & 0 & -2 \\ -2 & -2 & 0 \end{pmatrix} \right| = \frac{1}{6} | -16 | = \frac{8}{3}. \end{aligned}$$

### Example

Do the points  $(1, 2, 3)$ ,  $(2, 3, 5)$ ,  $(3, 5, 7)$ ,  $(5, 7, 11)$  lie in one plane?

### Example

Do the points  $(1, -1, -1)$ ,  $(-1, 1, -1)$ ,  $(-1, -1, 1)$ ,  $(1, 2, 3)$  lie in one plane?

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}$ ,  $\mathbf{v}$ ,  $\mathbf{w}$ ;



# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}$ ,  $\mathbf{v}$ ,  $\mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;
  - $\text{Vol}(R(\mathbf{u}, \mathbf{v}, \mathbf{w})) \neq 0$ ;

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;
  - $\text{Vol}(R(\mathbf{u}, \mathbf{v}, \mathbf{w})) \neq 0$ ;
  - $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \neq 0$ .

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;
  - $\text{Vol}(R(\mathbf{u}, \mathbf{v}, \mathbf{w})) \neq 0$ ;
  - $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \neq 0$ .
- If any of the above is valid:  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is a frame.

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;
  - $\text{Vol}(R(\mathbf{u}, \mathbf{v}, \mathbf{w})) \neq 0$ ;
  - $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \neq 0$ .
- If any of the above is valid:  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is a frame.
- Rectangular coordinate system  $\rightarrow$  fundamental frame  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;
  - $\text{Vol}(R(\mathbf{u}, \mathbf{v}, \mathbf{w})) \neq 0$ ;
  - $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \neq 0$ .
- If any of the above is valid:  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is a frame.
- Rectangular coordinate system  $\rightarrow$  fundamental frame  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$
- The hand rules for determining directions of cross products ( $\mathbf{w} = \mathbf{u} \times \mathbf{v}$ ) are consistent with this coordinate system if and only if

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) > 0$$

# Orientations of Space

- The following are equivalent:
  - every vector in space can be decomposed along  $\mathbf{u}, \mathbf{v}, \mathbf{w}$ ;
  - the box  $R(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is non-degenerate;
  - $\text{Vol}(R(\mathbf{u}, \mathbf{v}, \mathbf{w})) \neq 0$ ;
  - $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) \neq 0$ .
- If any of the above is valid:  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is a frame.
- Rectangular coordinate system  $\rightarrow$  fundamental frame  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$
- The hand rules for determining directions of cross products ( $\mathbf{w} = \mathbf{u} \times \mathbf{v}$ ) are consistent with this coordinate system if and only if

$$\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) > 0$$

## Definition

The frame  $(\mathbf{u}, \mathbf{v}, \mathbf{w})$  is positively oriented if  $\mathbf{u} \cdot (\mathbf{v} \times \mathbf{w}) > 0$ .