

# Calculus III

## Lecture 8

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Limits of Functions of Several Variables

# Outline

- 1 Limits of Functions of Several Variables
- 2 Continuity of Functions of Several Variables

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
`https://creativecommons.org/licenses/by/3.0/us/`  
and the links therein.

# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.

# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.
- Let  $P_0$  be point in plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;

# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.
- Let  $P_0$  be point in plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .

# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.
- Let  $P_0$  be point in plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .
- For example,

$$\begin{aligned} f: \mathbb{R}^2 \setminus \{P_0(0,0)\} &\rightarrow \mathbb{R} \\ f(x,y) &= \frac{x^2 y}{x^2 + y^2} \end{aligned}$$

- is defined arbitrarily close to  $(0,0)$ ;
- and is not defined at  $P_0(0,0)$ .



# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.
- Let  $P_0$  be point in plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .
- For example,

$$\begin{aligned} f: \mathbb{R}^2 \setminus \{P_0(0,0)\} &\rightarrow \mathbb{R} \\ f(x,y) &= \frac{x^2 y}{x^2 + y^2} \end{aligned}$$

- is defined arbitrarily close to  $(0,0)$ ;
- and is not defined at  $P_0(0,0)$ .

# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.
- Let  $P_0$  be point in plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .
- For example,

$$\begin{aligned} f: \mathbb{R}^2 \setminus \{P_0(0,0)\} &\rightarrow \mathbb{R} \\ f(x,y) &= \frac{x^2 y}{x^2 + y^2} \end{aligned}$$

- is defined arbitrarily close to  $(0,0)$ ;
- and is not defined at  $P_0(0,0)$ .

# Multivariable Limits

- Let  $f: D \rightarrow \mathbb{R}$ , where  $D$  a subset of the plane.
- Let  $P_0$  be point in plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .
- For example,

$$\begin{aligned} f: \mathbb{R}^2 \setminus \{P_0(0,0)\} &\rightarrow \mathbb{R} \\ f(x,y) &= \frac{x^2 y}{x^2 + y^2} \end{aligned}$$

- is defined arbitrarily close to  $(0,0)$ ;
  - and is not defined at  $P_0(0,0)$ .
- Question: What happens to  $f(Q)$  as  $Q$  gets closer to  $P_0$ ?

# Numerical Exploration of Limits - Example

## Example

$$f(x, y) = \frac{x^2 y}{x^2 + y^2}$$

$$f(Q) \rightarrow ? \text{ as } Q \rightarrow P_0(0, 0)$$

# Numerical Exploration of Limits - Example

## Example

$$f(x, y) = \frac{x^2 y}{x^2 + y^2}$$

$$f(Q) \rightarrow ? \text{ as } Q \rightarrow P_0(0, 0)$$

Numerical approach:

$$Q_1(0.1, 0.1) \mid f(Q_1) = f(0.1, 0.1) \simeq 0.05$$

# Numerical Exploration of Limits - Example

## Example

$$f(x, y) = \frac{x^2 y}{x^2 + y^2}$$

$$f(Q) \rightarrow ? \text{ as } Q \rightarrow P_0(0, 0)$$

Numerical approach:

$$\begin{array}{l|l} Q_1(0.1, 0.1) & f(Q_1) = f(0.1, 0.1) \simeq 0.05 \\ Q_2(0.01, -0.02) & f(Q_2) = f(0.01, -0.02) \simeq -0.004 \end{array}$$

# Numerical Exploration of Limits - Example

## Example

$$f(x, y) = \frac{x^2 y}{x^2 + y^2}$$

$$f(Q) \rightarrow ? \text{ as } Q \rightarrow P_0(0, 0)$$

Numerical approach:

$Q_1(0.1, 0.1)$	$f(Q_1)$	$=$	$f(0.1, 0.1) \simeq 0.05$
$Q_2(0.01, -0.02)$	$f(Q_2)$	$=$	$f(0.01, -0.02) \simeq -0.004$
$Q_3(-0.003, 0.001)$	$f(Q_3)$	$=$	$f(-0.003, 0.001) \simeq 0.0009$

# Numerical Exploration of Limits - Example

## Example

$$f(x, y) = \frac{x^2 y}{x^2 + y^2}$$

$$f(Q) \rightarrow ? \text{ as } Q \rightarrow P_0(0, 0)$$

Numerical approach:

$$\begin{array}{l|l} Q_1(0.1, 0.1) & f(Q_1) = f(0.1, 0.1) \simeq 0.05 \\ Q_2(0.01, -0.02) & f(Q_2) = f(0.01, -0.02) \simeq -0.004 \\ Q_3(-0.003, 0.001) & f(Q_3) = f(-0.003, 0.001) \simeq 0.0009 \end{array}$$

Numerical data suggests  $f(Q)$  approaches 0 as  $Q \rightarrow P_0(0, 0)$ .



# Definition of Multivariable Limit

- Let  $f: D \rightarrow \mathbb{R}$ , with  $D$  a subset of the plane.
- Let  $P_0$  be a point in the plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .

# Definition of Multivariable Limit

- Let  $f: D \rightarrow \mathbb{R}$ , with  $D$  a subset of the plane.
- Let  $P_0$  be a point in the plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .

## Definition

$L$  is **the limit of  $f$  at  $P_0$**  if we can keep the values of  $f(Q)$  as close to  $L$  as we want by keeping  $Q$  close enough to  $P_0$ , but not equal to  $P_0$ .

# Definition of Multivariable Limit

- Let  $f: D \rightarrow \mathbb{R}$ , with  $D$  a subset of the plane.
- Let  $P_0$  be a point in the plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .

## Definition

$L$  is **the limit of  $f$  at  $P_0$**  if we can keep the values of  $f(Q)$  as close to  $L$  as we want by keeping  $Q$  close enough to  $P_0$ , but not equal to  $P_0$ . We write:

$$L = \lim_{Q \rightarrow P_0} f(Q) \quad \text{or}$$

$$L = \lim_{(x,y) \rightarrow (x_0,y_0)} f(x,y)$$

# Definition of Multivariable Limit

- Let  $f: D \rightarrow \mathbb{R}$ , with  $D$  a subset of the plane.
- Let  $P_0$  be a point in the plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .

## Definition

$L$  is **the limit of  $f$  at  $P_0$**  if we can keep the values of  $f(Q)$  as close to  $L$  as we want by keeping  $Q$  close enough to  $P_0$ , but not equal to  $P_0$ . We write:

$$L = \lim_{Q \rightarrow P_0} f(Q) \quad \text{or}$$

$$L = \lim_{(x,y) \rightarrow (x_0,y_0)} f(x,y)$$

- As usual, we extend the definition to allow  $L = \infty$ . By convention, if  $M > N$ , we say that  $M$  is closer to  $\infty$  than  $N$ .

# Definition of Multivariable Limit

- Let  $f: D \rightarrow \mathbb{R}$ , with  $D$  a subset of the plane.
- Let  $P_0$  be a point in the plane such that:
  - $f$  is defined arbitrarily close to  $P_0$ ;
  - $f$  is not necessarily defined at  $P_0$ .

## Definition

$L$  is **the limit of  $f$  at  $P_0$**  if we can keep the values of  $f(Q)$  as close to  $L$  as we want by keeping  $Q$  close enough to  $P_0$ , but not equal to  $P_0$ . We write:

$$L = \lim_{Q \rightarrow P_0} f(Q) \quad \text{or}$$

$$L = \lim_{(x,y) \rightarrow (x_0,y_0)} f(x,y)$$

- As usual, we extend the definition to allow  $L = \infty$ . By convention, if  $M > N$ , we say that  $M$  is closer to  $\infty$  than  $N$ .
- If the limit  $L$  exists, it is unique.

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.



## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if ?

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$   
in polar coordinates.

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} = \lim_{r \rightarrow 0}$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} = \lim_{r \rightarrow 0}$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$   
in polar coordinates.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} = \lim_{r \rightarrow 0} ?$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$   
in polar coordinates.

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} = \lim_{r \rightarrow 0} \frac{r^2 \cos^2 \theta r \sin \theta}{r^2}$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} &= \lim_{r \rightarrow 0} \frac{\cancel{r^2} \cos^2 \theta r \sin \theta}{\cancel{r^2} \sin^2 \theta} \\ &= \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \end{aligned}$$



## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} &= \lim_{r \rightarrow 0} \frac{\cancel{r^2} \cos^2 \theta r \sin \theta}{\cancel{r^2}} \\ &= \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \end{aligned}$$

$$-r \leq r \cos^2 \theta \sin \theta \leq r$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} &= \lim_{r \rightarrow 0} \frac{\cancel{r^2} \cos^2 \theta r \sin \theta}{\cancel{r^2}} \\ &= \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \end{aligned}$$

$$\lim_{r \rightarrow 0} -r \leq \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \leq \lim_{r \rightarrow 0} r$$

## Example

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.


Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} &= \lim_{r \rightarrow 0} \frac{\cancel{r^2} \cos^2 \theta r \sin \theta}{\cancel{r^2}} \\ &= \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \end{aligned}$$

$$0 = \lim_{r \rightarrow 0} -r \leq \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \leq \lim_{r \rightarrow 0} r = 0.$$

# Example



$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2}$$

- $f$  is not defined at  $P_0(0, 0)$ ;
- Even if it were, the actual value might be different from the limit.

Polar coordinates:  $x = r \cos \theta$ ,  $y = r \sin \theta$ .

We have:  $(x, y) \rightarrow (0, 0)$  if and only if  $r \rightarrow 0$  in polar coordinates.

$$\begin{aligned} \lim_{(x,y) \rightarrow (0,0)} \frac{x^2 y}{x^2 + y^2} &= \lim_{r \rightarrow 0} \frac{\cancel{r^2} \cos^2 \theta r \sin \theta}{\cancel{r^2}} \\ &= \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \\ &= 0 \end{aligned}$$

For the last equality, we use the squeeze theorem:

$$0 = \lim_{r \rightarrow 0} -r \leq \lim_{r \rightarrow 0} r \cos^2 \theta \sin \theta \leq \lim_{r \rightarrow 0} r = 0.$$

If  $\mathbf{r} = (a, b)$  is a vector, by  $f(\mathbf{r})$  we understand  $f(a, b)$  (i.e., define  $f(\mathbf{r})$  via the vector-point identification).

## Definition

If  $\mathbf{r} = (a, b)$  is a vector, by  $f(\mathbf{r})$  we understand  $f(a, b)$  (i.e., define  $f(\mathbf{r})$  via the vector-point identification).

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;

If  $\mathbf{r} = (a, b)$  is a vector, by  $f(\mathbf{r})$  we understand  $f(a, b)$  (i.e., define  $f(\mathbf{r})$  via the vector-point identification).

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near  $P$  with position vector  $\mathbf{r}$ ;  $f$  is not necessarily defined at  $P$ ;

If  $\mathbf{r} = (a, b)$  is a vector, by  $f(\mathbf{r})$  we understand  $f(a, b)$  (i.e., define  $f(\mathbf{r})$  via the vector-point identification).

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near  $P$  with position vector  $\mathbf{r}$ ;  $f$  is not necessarily defined at  $P$ ;
- let  $\mathbf{u}$  be an arbitrary vector.



If  $\mathbf{r} = (a, b)$  is a vector, by  $f(\mathbf{r})$  we understand  $f(a, b)$  (i.e., define  $f(\mathbf{r})$  via the vector-point identification).

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near  $P$  with position vector  $\mathbf{r}$ ;  $f$  is not necessarily defined at  $P$ ;
- let  $\mathbf{u}$  be an arbitrary vector.

We say that the one-variable limit

$$\lim_{t \rightarrow 0} f(\mathbf{r} + t\mathbf{u})$$

is the limit of  $f$  along the direction  $\mathbf{u}$ .

If  $\mathbf{r} = (a, b)$  is a vector, by  $f(\mathbf{r})$  we understand  $f(a, b)$  (i.e., define  $f(\mathbf{r})$  via the vector-point identification).

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near  $P$  with position vector  $\mathbf{r}$ ;  $f$  is not necessarily defined at  $P$ ;
- let  $\mathbf{u}$  be an arbitrary vector.

We say that the one-variable limit

$$\lim_{t \rightarrow 0} f(\mathbf{r} + t\mathbf{u})$$

is the limit of  $f$  along the direction  $\mathbf{u}$ .

## Theorem

*If the limit  $\lim_{Q \rightarrow P} f(Q)$  exists, then every directional limit  $\lim_{t \rightarrow 0} f(\mathbf{r} + t\mathbf{u})$  exists and all directional limits are equal.*

## Example (Limit may fail to exist)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

.

## Example (Limit may fail to exist)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\lim_{t \rightarrow 0} f(t\mathbf{u}) = \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm)$$

.

## Example (Limit may fail to exist)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2 t^2} \end{aligned}$$

.

## Example (Limit may fail to exist)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{m t^2 + m^2 t^2} \\ &= \frac{1}{1 + m^2} \end{aligned}$$

## Example (Limit may fail to exist)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2 t^2} \\ &= \frac{m}{1 + m^2} . \end{aligned}$$

Directional limit depends on  $m \Rightarrow$

## Example (Limit may fail to exist)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$


Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2 t^2} \\ &= \frac{m}{1 + m^2} \end{aligned}$$

Directional limit depends on  $m \Rightarrow$  directional limit is not the same for all values of  $\mathbf{u}$



## Example (Limit may fail to exist)




$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2t^2} \\ &= \frac{m}{1 + m^2} \end{aligned}$$

Directional limit depends on  $m \Rightarrow$  directional limit is not the same for all values of  $\mathbf{u} \Rightarrow$  the multivariable limit does not exist.

## Example (Limit may fail to exist)



$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :


$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2 t^2} \\ &= \frac{m}{1 + m^2} \end{aligned}$$

Directional limit depends on  $m \Rightarrow$  directional limit is not the same for all values of  $\mathbf{u} \Rightarrow$  the multivariable limit does not exist.

If we'd used polar coordinates, we would have obtained:

$$\frac{xy}{x^2 + y^2} =$$

## Example (Limit may fail to exist)



$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :


$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2 t^2} \\ &= \frac{m}{1 + m^2} \end{aligned}$$

Directional limit depends on  $m \Rightarrow$  directional limit is not the same for all values of  $\mathbf{u} \Rightarrow$  the multivariable limit does not exist.

If we'd used polar coordinates, we would have obtained:

$$\frac{xy}{x^2 + y^2} = \cos \theta \sin \theta$$

## Example (Limit may fail to exist)



$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy}{x^2 + y^2}$$

Let  $\mathbf{u} = (1, m)$ . Directional limit along  $\mathbf{u}$ :

$$\begin{aligned} \lim_{t \rightarrow 0} f(t\mathbf{u}) &= \lim_{t \rightarrow 0} f(t(1, m)) = \lim_{t \rightarrow 0} f(t, tm) \\ &= \lim_{t \rightarrow 0} \frac{mt^2}{t^2 + m^2 t^2} \\ &= \frac{m}{1 + m^2} \end{aligned}$$

Directional limit depends on  $m \Rightarrow$  directional limit is not the same for all values of  $\mathbf{u} \Rightarrow$  the multivariable limit does not exist.

If we'd used polar coordinates, we would have obtained:

$$\frac{xy}{x^2 + y^2} = \cos \theta \sin \theta$$

This expression depends only on  $\theta$ ; as  $r \rightarrow 0$  permits arbitrary behavior of  $\theta$ , we'd have guessed correctly that the limit doesn't exist.

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.
- Similarities b-n side and directional limits.
  - Single variable functions: limit exists  $\implies$  side limits exist, have the same value.

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.
- Similarities b-n side and directional limits.
  - Single variable functions: limit exists  $\implies$  side limits exist, have the same value.
  - Multivariable functions: limit exists  $\implies$  directional limits exist, have the same value.



# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.
- Similarities b-n side and directional limits.
  - Single variable functions: limit exists  $\implies$  side limits exist, have the same value.
  - Multivariable functions: limit exists  $\implies$  directional limits exist, have the same value.
  - Single variable functions: side limits are different  $\implies$  limit does not exist.

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.
- Similarities b-n side and directional limits.
  - Single variable functions: limit exists  $\implies$  side limits exist, have the same value.
  - Multivariable functions: limit exists  $\implies$  directional limits exist, have the same value.
  - Single variable functions: side limits are different  $\implies$  limit does not exist.
  - Multivariable functions: directional limits have different values  $\implies$  limit does not exist.

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.
- Similarities b-n side and directional limits.
  - Single variable functions: limit exists  $\implies$  side limits exist, have the same value.
  - Multivariable functions: limit exists  $\implies$  directional limits exist, have the same value.
  - Single variable functions: side limits are different  $\implies$  limit does not exist.
  - Multivariable functions: directional limits have different values  $\implies$  limit does not exist.
- Differences b-n side and directional limits.
  - Single variable functions: Side limits are equal  $\implies$  limit exists.

# Side Limits and Directional Limits

- Directional limits in dimension 1 are equal to the left or right hand limits (depending on the direction of the vector).
- Directional limits are therefore the natural analog of 1-dim side limits.
- Similarities b-n side and directional limits.
  - Single variable functions: limit exists  $\implies$  side limits exist, have the same value.
  - Multivariable functions: limit exists  $\implies$  directional limits exist, have the same value.
  - Single variable functions: side limits are different  $\implies$  limit does not exist.
  - Multivariable functions: directional limits have different values  $\implies$  limit does not exist.
- Differences b-n side and directional limits.
  - Single variable functions: Side limits are equal  $\implies$  limit exists.
  - Multivariable functions: even if all directional limits have the same value the limit does not necessarily exist.

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} =$$

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4}$$

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4}$$



## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

# Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

For direction  $x = 0$ ,  $y = m$ : directional limit is again 0.

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

For direction  $x = 0$ ,  $y = m$ : directional limit is again 0.  $\Rightarrow$  all directional limits exist and equal 0.

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

For direction  $x = 0, y = m$ : directional limit is again 0.  $\Rightarrow$  all directional limits exist and equal 0. However, along  $x = y^2$ :

$$\frac{xy^2}{x^2 + y^4} =$$

# Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

For direction  $x = 0$ ,  $y = m$ : directional limit is again 0.  $\Rightarrow$  all directional limits exist and equal 0. However, along  $x = y^2$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{y^4}{y^4 + y^4} =$$

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

For direction  $x = 0, y = m$ : directional limit is again 0.  $\Rightarrow$  all directional limits exist and equal 0. However, along  $x = y^2$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{y^4}{y^4 + y^4} = \frac{1}{2}$$

## Example ( )

$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :


$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0$$

as  $x \rightarrow 0$ .

For direction  $x = 0, y = m$ : directional limit is again 0.  $\Rightarrow$  all directional limits exist and equal 0. However, along  $x = y^2$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{y^4}{y^4 + y^4} = \frac{1}{2} \rightarrow \frac{1}{2} \text{ as } x \rightarrow 0$$

## Example (All directional limits exist, limit doesn't)



$$\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$$

Along  $y = mx$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{m^2 x^3}{x^2 + m^4 x^4} = \frac{m^2 x}{1 + m^2 x^4} \rightarrow 0 \text{ as } x \rightarrow 0.$$

For direction  $x = 0, y = m$ : directional limit is again 0.  $\Rightarrow$  all directional limits exist and equal 0. However, along  $x = y^2$ :

$$\frac{xy^2}{x^2 + y^4} = \frac{y^4}{y^4 + y^4} = \frac{1}{2} \rightarrow \frac{1}{2} \text{ as } x \rightarrow 0$$

Therefore  $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2 + y^4}$  does not exist.



# Limits along paths

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;

# Limits along paths

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near point  $P$  with position vector  $\mathbf{p}$ .

# Limits along paths

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near point  $P$  with position vector  $\mathbf{p}$ .
- Let  $\mathbf{r}(t) = (x(t), y(t))$ ,  $t \in I$  be a continuous path such that:
  - 0 is in  $I$ ,  $\mathbf{r}(0) = \mathbf{p}$ ;

# Limits along paths

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near point  $P$  with position vector  $\mathbf{p}$ .
- Let  $\mathbf{r}(t) = (x(t), y(t))$ ,  $t \in I$  be a continuous path such that:
  - 0 is in  $I$ ,  $\mathbf{r}(0) = \mathbf{p}$ ;
  - $\mathbf{r}$  is continuous at 0;

# Limits along paths

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near point  $P$  with position vector  $\mathbf{p}$ .
- Let  $\mathbf{r}(t) = (x(t), y(t))$ ,  $t \in I$  be a continuous path such that:
  - 0 is in  $I$ ,  $\mathbf{r}(0) = \mathbf{p}$ ;
  - $\mathbf{r}$  is continuous at 0;
  - $\mathbf{r}(t)$  lies in  $D$  for  $t \neq 0$

# Limits along paths

## Definition

- Let  $f : D \rightarrow \mathbb{R}$ , where  $D$  is a region in the plane;
- let  $f$  be defined near point  $P$  with position vector  $\mathbf{p}$ .
- Let  $\mathbf{r}(t) = (x(t), y(t))$ ,  $t \in I$  be a continuous path such that:
  - 0 is in  $I$ ,  $\mathbf{r}(0) = \mathbf{p}$ ;
  - $\mathbf{r}$  is continuous at 0;
  - $\mathbf{r}(t)$  lies in  $D$  for  $t \neq 0$

We say that the one-variable limit

$$\lim_{t \rightarrow 0} f(\mathbf{r}(t))$$

is the limit of  $f$  along the path  $\mathbf{r}(t)$ .

## Theorem

*If the limit  $\lim_{Q \rightarrow P} f(Q)$  exists, then every path limit exists and all path limits are equal.*

## Theorem

*If the limit  $\lim_{Q \rightarrow P} f(Q)$  exists, then every path limit exists and all path limits are equal.*

- If we pick our path to be of the form  $\mathbf{r}(t) = \mathbf{r}_0 + t\mathbf{u}$ , we see that the directional limit is a special case of the path limit.



# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.
- If defined, quotients of continuous are continuous.

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.
- If defined, quotients of continuous are continuous.
- Powers, exponentials of continuous functions are continuous.

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.
- If defined, quotients of continuous are continuous.
- Powers, exponentials of continuous functions are continuous.
- Compositions of continuous functions are continuous.

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.
- If defined, quotients of continuous are continuous.
- Powers, exponentials of continuous functions are continuous.
- Compositions of continuous functions are continuous.
- A function fails to be continuous if:



# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.
- If defined, quotients of continuous are continuous.
- Powers, exponentials of continuous functions are continuous.
- Compositions of continuous functions are continuous.
- A function fails to be continuous if:
  - (Removable discontinuity) limit exists but is different from  $f(x_0, y_0)$  value;

# Continuity

## Definition (Continuous at a point)

We say that  $f$  is continuous at  $(x_0, y_0)$  if  $\lim_{(x,y) \rightarrow (x_0,y_0)} f(x, y) = f(x_0, y_0)$ .

## Definition (Continuity)

We say that  $f$  is a continuous function if it is continuous at all points where it is defined.

- Polynomial functions are continuous.
- Sum, difference, product of continuous functions are continuous.
- If defined, quotients of continuous are continuous.
- Powers, exponentials of continuous functions are continuous.
- Compositions of continuous functions are continuous.
- A function fails to be continuous if:
  - (Removable discontinuity) limit exists but is different from  $f(x_0, y_0)$  value;
  - (Essential discontinuity) the limit does not exist.

# Continuity of vector fields

- Recall that a vector field is a function

$$\begin{aligned}\mathbf{F}: D &\rightarrow \mathbb{R}^2 \\ \mathbf{F}(x, y) &= F_1(x, y)\mathbf{i} + F_2(x, y)\mathbf{j}\end{aligned}$$

# Continuity of vector fields

- Recall that a vector field is a function

$$\begin{aligned}\mathbf{F}: D &\rightarrow \mathbb{R}^2 \\ \mathbf{F}(x, y) &= F_1(x, y)\mathbf{i} + F_2(x, y)\mathbf{j}\end{aligned}$$

- $F_1, F_2$  are two-variable functions with scalar output.

# Continuity of vector fields

- Recall that a vector field is a function

$$\begin{aligned}\mathbf{F}: D &\rightarrow \mathbb{R}^2 \\ \mathbf{F}(x, y) &= F_1(x, y)\mathbf{i} + F_2(x, y)\mathbf{j}\end{aligned}$$

- $F_1, F_2$  are two-variable functions with scalar output.
- We have already defined the notion of continuity of  $F_1$  and  $F_2$ .


# Continuity of vector fields

- Recall that a vector field is a function

$$\begin{aligned}\mathbf{F}: D &\rightarrow \mathbb{R}^2 \\ \mathbf{F}(x, y) &= F_1(x, y)\mathbf{i} + F_2(x, y)\mathbf{j}\end{aligned}$$

- $F_1, F_2$  are two-variable functions with scalar output.
- We have already defined the notion of continuity of  $F_1$  and  $F_2$ .
- We define  $\mathbf{F}$  to be continuous if  $F_1, F_2$  are continuous.


## Example (continuous vector field)



Discuss the continuity of the following vector field.

$$\mathbf{F}(x, y) = \frac{y}{3}\mathbf{i} - \frac{x}{3}\mathbf{j} \quad .$$

## Example (continuous vector field)



Discuss the continuity of the following vector field.


$$\mathbf{F}(x, y) = \frac{y}{3}\mathbf{i} - \frac{x}{3}\mathbf{j} .$$

Then

$$F_1(x, y) = \frac{y}{3} , \quad F_2(x, y) = -\frac{x}{3}$$



## Example (continuous vector field)



Discuss the continuity of the following vector field.


$$\mathbf{F}(x, y) = \frac{y}{3}\mathbf{i} - \frac{x}{3}\mathbf{j} .$$

Then

$$F_1(x, y) = \frac{y}{3} , \quad F_2(x, y) = -\frac{x}{3}$$

We have that  $\frac{y}{3}, -\frac{x}{3}$  are two-variable polynomials

## Example (continuous vector field)



Discuss the continuity of the following vector field.


$$\mathbf{F}(x, y) = \frac{y}{3}\mathbf{i} - \frac{x}{3}\mathbf{j} .$$

Then

$$F_1(x, y) = \frac{y}{3} , \quad F_2(x, y) = -\frac{x}{3}$$

We have that  $\frac{y}{3}, -\frac{x}{3}$  are two-variable polynomials  $\implies F_1$  and  $F_2$  are continuous

## Example (continuous vector field)



Discuss the continuity of the following vector field.

$$\mathbf{F}(x, y) = \frac{y}{3}\mathbf{i} - \frac{x}{3}\mathbf{j} .$$

Then

$$F_1(x, y) = \frac{y}{3} , \quad F_2(x, y) = -\frac{x}{3}$$

We have that  $\frac{y}{3}, -\frac{x}{3}$  are two-variable polynomials  $\implies F_1$  and  $F_2$  are continuous  $\implies$  the vector field is continuous.