

Calculus II

Lecture (not covered in class)

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

- 1 Exponential Functions and logarithms, Review
- 2 Derivatives of Exponential Functions
 - Natural Exponent
- 3 A More Advanced Approach to Exponents
 - Derivative using series definition
- 4 Logarithmic Functions, Review
 - Natural Logarithms
- 5 Derivatives of Logarithms, Review
 - The Natural Logarithm
 - The Number e as a Limit
 - Derivatives of Exponents with Arbitrary Base
 - Derivatives of Arbitrary Exponents with Arbitrary Base

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:


<https://github.com/tmilev/freecalc>

- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:

<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.

Exponential Functions

The function $f(x) = 2^x$ is called an exponential function because the variable x is the exponent.




x	y
2	4
1	2
0	1
-1	$\frac{1}{2}$
-2	$\frac{1}{4}$

(Exponential Function Terminology)

An exponential function is a function of the form $f(x) = a^x$, where a is a positive constant.

Graphs of various exponential functions.



Derivatives of Exponential Functions

Compute the derivative of $f(x) = a^x$ using the definition:

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \lim_{h \rightarrow 0} \frac{a^{x+h} - a^x}{h} \\ &= \lim_{h \rightarrow 0} \frac{a^x a^h - a^x}{h} \\ &= \lim_{h \rightarrow 0} \frac{a^x (a^h - 1)}{h} \\ &= a^x \lim_{h \rightarrow 0} \frac{a^h - 1}{h} \\ &= a^x \lim_{h \rightarrow 0} \frac{a^h - a^0}{h} \\ &= a^x f'(0). \end{aligned}$$

We have shown that, if $f(x) = a^x$ is differentiable at 0, then it is differentiable everywhere, and

$$f'(x) = f'(0)a^x.$$

We leave the following theorem without proof.

Theorem

Let a be a positive number and let $f(x) = a^x$. Then the limit

$$f'(0) = \lim_{h \rightarrow 0} \frac{a^h - 1}{h}$$

exists.




We will later show that

$$f'(0) = \lim_{h \rightarrow 0} \frac{a^h - 1}{h} = \ln(a).$$

Here, \ln is the natural logarithm function.

The Natural Exponential Function

- One base for an exponential function is especially useful.
- It has a special property: its tangent line at $x = 0$ has slope $m = 1$.
- We call this number e , known as Euler's number or Napier's constant.
- e is a number between 2 and 3.
- In fact, $e = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \cdots \approx 2.71828$.



Definition (Natural Exponential Function)

e^x is called the natural exponential function. Its derivative is

$$\frac{d}{dx}(e^x) = e^x.$$

Exponents overview

- For integer x , we know how to compute a^x as a function of a .
- How do we compute $f(x) = a^x$ when x is not an integer?
- We need to go back to the definition of a^x (for x non-integer).
- In what follows we give/recall an elementary way to define exponent.
- Then we give an alternative second definition.
- The second definition will be studied in sufficient depth only much later.
- The two definitions are equivalent: if we choose one definition the other becomes a theorem and the other way round.
- Choosing one definition makes some statements easier to prove and others more difficult.
- We shall discuss pros and cons of the two. In a nutshell:
 - the first elementary definition is easier to motivate;
 - the second alternative definition is easier to compute with.

Exponent definition using limits (approach I)

- For integer p we know to compute a^p .
- Therefore for integer q we know to compute $a^{\frac{1}{q}} = \sqrt[q]{a} = \max\{x \mid x^q \leq a\}$.
- Therefore we know to compute $a^{\frac{p}{q}}$ for all rational $\frac{p}{q}$.
- We can then define

$$a^x = \lim_{\substack{y \rightarrow x \\ y\text{-rational}}} a^y$$

For example, a^π would be defined as the limit of the sequence $a^{3.14}, a^{3.141}, a^{3.1415}, \dots$

- Cons: not computationally effective; not how computers compute.
- Pros: for non-integer x and y , it is very easy to prove that $a^{x+y} = a^x a^y$ - this follows from the definition of limit above.
- This is the definition assumed in many elementary courses.

Exponent definition using series (approach II)

- The following formula (studied much later) can be used as alternative definition.

$$e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!} = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \cdots + \frac{x^n}{n!} + \cdots$$

Here $n! = 1 \cdot 2 \cdot 3 \cdots (n-1) \cdot n$ and is read “ n factorial”.

- For $|x| < 1$ define

$$\ln(1+x) = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{x^n}{n} = x - \frac{x^2}{2} + \frac{x^3}{3} - \cdots + \frac{(-1)^{n+1} x^n}{n} + \cdots$$

Infinite sum studied much later.

- For arbitrary $a > 0$ define a^x as $a^x = e^{x \ln a}$.
- Cons: more difficult to prove $e^{x+y} = e^x e^y$ and $e^{\ln(1+x)} = 1+x$, proof done later.
- Pros: this is how e^x and a^x are actually computed (by modern computers and by humans in the past).

Example

Derive the exponent rule $(e^x)' = e^x$ using the Calc II formula below, the infinite (both sides uniformly convergent) sum rule

$(f_1 + f_2 + f_3 + \dots)' = f_1' + f_2' + f_3' + \dots$ and the power rule $(x^n)' = nx^{n-1}$.

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots,$$


where $n! = 1 \cdot 2 \cdot 3 \cdot \dots \cdot n$. We have that

$$\frac{n}{n!} = \frac{n}{1 \cdot 2 \cdot \dots \cdot (n-1) \cdot n} = \frac{1}{1 \cdot 2 \cdot \dots \cdot (n-1)} = \frac{1}{(n-1)!}.$$

$$\begin{aligned} (e^x)' &= \left(1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots\right)' \\ &= (1)' + (x)' + \frac{(x^2)'}{2!} + \frac{(x^3)'}{3!} + \dots + \frac{(x^n)'}{n!} + \dots \\ &= 0 + 1 + \frac{2x}{2!} + \frac{3x^2}{3!} + \dots + \frac{nx^{n-1}}{n!} + \dots \\ &= 1 + \frac{x}{1!} + \frac{x^2}{2!} + \dots + \frac{x^{n-1}}{(n-1)!} + \dots = e^x \end{aligned}$$

as desired.

Logarithmic Functions



- Suppose $a > 0$, $a \neq 1$.
- Let $f(x) = a^x$.
- Then f is either increasing or decreasing.
- Therefore f is one-to-one.
- Therefore f has an inverse function, f^{-1} .
- The graph shows $y = a^x$ for $a > 1$.
- The graph of $y = \log_a x$ is the reflection of this in the line $y = x$.

Definition ($\log_a x$)

The inverse function of $f(x) = a^x$ is called the logarithmic function with base a , and is written $\log_a x$. It is defined by the formula


$$\log_a x = y \quad \Leftrightarrow \quad a^y = x.$$

If $x > 0$, then $\log_a x$ is the exponent to which the base a must be raised to give x .


Example

Evaluate:

- 1 $\log_3 81 = 4$ because $3^4 = 81$.
- 2 $\log_{25} 5 = \frac{1}{2}$ because $25^{\frac{1}{2}} = \sqrt{25} = 5$.
- 3 $\log_{10} 0.001 = -3$ because $10^{-3} = 0.001$.



- Suppose $a > 1$.
- Domain of a^x : \mathbb{R} .
- Range of a^x : $(0, \infty)$.
- Domain of $\log_a x$: $(0, \infty)$.
- Range of $\log_a x$: \mathbb{R} .
- $\log_a(a^x) = x$ for $x \in \mathbb{R}$.
- $a^{\log_a x} = x$ for $x > 0$.


Graphs of various logarithmic functions with $a > 1$ 

Natural Logarithms

Definition ($\ln x$)

The logarithm with base e is called the natural logarithm, and has a special notation:

$$\log_e x = \ln x.$$



- $\ln x = y \quad \Leftrightarrow \quad e^y = x.$
- $\ln(e^x) = x$ for $x \in \mathbb{R}.$
- $e^{\ln x} = x$ for $x > 0.$

Theorem (Properties of Logarithmic Functions)

If $a > 1$, the function $f(x) = \log_a x$ is a one-to-one, continuous, increasing function with domain $(0, \infty)$ and range \mathbb{R} . If $x, y, a, b > 0$ and r is any real number, then

- 1 $\log_a(xy) = \log_a x + \log_a y.$
- 2 $\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y.$
- 3 $\log_a(x^r) = r \log_a x.$
- 4 $\log_a(x) = \log_b x \log_a b = \frac{\log_b x}{\log_b a} = \frac{\ln x}{\ln a}.$

The Derivative of the Natural Logarithm

Theorem (The Derivative of $\ln x$)

$$\frac{d}{dx}(\ln x) = \frac{1}{x}.$$

Proof.

- Let $y = \ln x$.
- Then $e^y = x$.
- Differentiate this implicitly with respect to x :
- $e^y \frac{dy}{dx} = 1$.
- Rearrange:
- $\frac{d}{dx}(\ln x) = \frac{dy}{dx} = \frac{1}{e^y} = \frac{1}{e^{\ln x}} = \frac{1}{x}$.



Example (Chain Rule,)

Differentiate $y = \ln(x^3 + 1)$.

Let $u = x^3 + 1$.

Then $y = \ln u$.

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left(\frac{1}{u} \right) (3x^2) \\ &= \frac{3x^2}{x^3 + 1}.\end{aligned}$$

Theorem (The Number e as a Limit)

$$e = \lim_{x \rightarrow 0} (1 + x)^{\frac{1}{x}} = \lim_{y \rightarrow \infty} \left(1 + \frac{1}{y}\right)^y.$$

Proof.

Let $f(x) = \ln x$. Then $f'(x) = \frac{1}{x}$, so $f'(1) = 1$.

$$\begin{aligned} 1 = f'(1) &= \lim_{h \rightarrow 0} \frac{f(1+h) - f(1)}{h} = \lim_{x \rightarrow 0} \frac{f(1+x) - f(1)}{x} \\ &= \lim_{x \rightarrow 0} \frac{\ln(1+x) - \ln(1)}{x} = \lim_{x \rightarrow 0} \frac{1}{x} \ln(1+x) \\ &= \lim_{x \rightarrow 0} \ln(1+x)^{\frac{1}{x}}. \end{aligned}$$

Then use the fact that the exponential function is continuous:

$$e = e^1 = e^{\lim_{x \rightarrow 0} \ln(1+x)^{\frac{1}{x}}} = \lim_{x \rightarrow 0} e^{\ln(1+x)^{\frac{1}{x}}} = \lim_{x \rightarrow 0} (1+x)^{\frac{1}{x}}. \quad \square$$

Example

Compute

$$\begin{aligned}\lim_{x \rightarrow \infty} \left(\frac{x+3}{x} \right)^x &= \lim_{x \rightarrow \infty} \left(1 + \frac{3}{x} \right)^x \\ &= \lim_{x \rightarrow \infty} \left(1 + \frac{1}{\frac{x}{3}} \right)^{3 \frac{x}{3}} \\ &= \lim_{\substack{x \rightarrow \infty \\ \frac{x}{3} = y \rightarrow \infty}} \left(1 + \frac{1}{y} \right)^{3y} \\ &= \lim_{y \rightarrow \infty} \left(\left(1 + \frac{1}{y} \right)^y \right)^3 = e^3.\end{aligned}$$

Set $\frac{x}{3} = y$

Example

Compute

$$\begin{aligned}
 & \lim_{x \rightarrow \infty} \left(\frac{x}{x-2} \right)^{2x+2} \\
 = & \lim_{x \rightarrow \infty} \left(\frac{x-2+2}{x-2} \right)^{2x+2} = \lim_{x \rightarrow \infty} \left(1 + \frac{2}{x-2} \right)^{2x+2} \\
 = & \lim_{x \rightarrow \infty} \left(1 + \frac{1}{\frac{x-2}{2}} \right)^{2(x-2+2)+2} \\
 = & \lim_{x \rightarrow \infty} \left(1 + \frac{1}{\frac{x-2}{2}} \right)^{4 \cdot \frac{x-2}{2} + 6} = \lim_{\substack{\frac{x-2}{2} = y \\ y \rightarrow \infty}} \left(1 + \frac{1}{y} \right)^{4y+6} \\
 = & \lim_{y \rightarrow \infty} \left(\left(1 + \frac{1}{y} \right)^y \right)^4 \lim_{y \rightarrow \infty} \left(1 + \frac{1}{y} \right)^6 \\
 = & e^4 \cdot (1+0)^6 = e^4.
 \end{aligned}$$

Set $y = \frac{x-2}{2}$

Theorem (The Derivative of a^x)

$$\frac{d}{dx}(a^x) = a^x \ln a.$$

Proof.

Use the fact that $a = e^{\ln a}$.

$$\begin{aligned}\frac{d}{dx}(a^x) &= \frac{d}{dx} \left((e^{\ln a})^x \right) \\ &= \frac{d}{dx} \left(e^{x \ln a} \right) \\ &= e^{x \ln a} \frac{d}{dx} (x \ln a) \\ &= (e^{\ln a})^x \ln a \\ &= a^x (\ln a).\end{aligned}$$



Example (Chain Rule)

Differentiate $y = 10^{x^2}$.

Let $u = x^2$.

Then $y = 10^u$.

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (10^u (\ln 10)) (2x) \\ &= (2 \ln 10) x 10^{x^2}.\end{aligned}$$

Example

Compute $\frac{d}{dx} \left((\tan x)^{\frac{1}{x}} \right)$, where $x \in (0, \frac{\pi}{2})$.

$$\begin{aligned} \frac{d}{dx} \left((\tan x)^{\frac{1}{x}} \right) &= \frac{d}{dx} \left((e^{\ln \tan x})^{\frac{1}{x}} \right) = \frac{d}{dx} \left(e^{\frac{1}{x} \ln \tan x} \right) \\ &= e^{\frac{1}{x} \ln(\tan x)} \frac{d}{dx} \left(\frac{1}{x} \ln(\tan x) \right) \\ &= (\tan x)^{\frac{1}{x}} \left(-\frac{1}{x^2} \ln(\tan x) + \frac{1}{x} \frac{(\tan x)'}{\tan x} \right) \\ &= (\tan x)^{\frac{1}{x}} \left(-\frac{1}{x^2} \ln(\tan x) + \frac{1}{x} \frac{\frac{1}{\cos^2(x)}}{\frac{\sin x}{\cos x}} \right) \\ &= (\tan x)^{\frac{1}{x}} \left(-\frac{1}{x^2} \ln(\tan x) + \frac{1}{x} \frac{1}{\sin x \cos x} \right) \end{aligned}$$

Example

Suppose $g(x)$ and $f(x)$ are differentiable functions and suppose $g(x) > 0$. Prove that

$$\frac{d}{dx} \left(g(x)^{f(x)} \right) = g(x)^{f(x)} \left(f'(x) \ln(g(x)) + f(x) \frac{g'(x)}{g(x)} \right) .$$

Proof.

$$\begin{aligned} \frac{d}{dx} \left(g(x)^{f(x)} \right) &= \frac{d}{dx} \left(\left(e^{\ln g(x)} \right)^{f(x)} \right) = \frac{d}{dx} \left(e^{f(x) \ln g(x)} \right) \\ &= e^{f(x) \ln g(x)} \frac{d}{dx} (f(x) \ln g(x)) \\ &= g(x)^{f(x)} \left(f'(x) \ln(g(x)) + f(x) \frac{g'(x)}{g(x)} \right) , \end{aligned}$$

as desired. □