

# Calculus I

## Lecture 11

### The Chain Rule

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

- 1 The Chain Rule
  - Chain rule proof

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = ?$  .
- $u = h(x) = ?$  .
- Then  $y = f(x) = g(h(x)) =$

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = ?$  .
- $u = h(x) = ?$  .
- Then  $y = f(x) = g(h(x)) =$



# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = ?$  .
- Then  $y = f(x) = g(h(x)) =$

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = ?$  .
- Then  $y = f(x) = g(h(x)) =$

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = ?$  .
- Then  $y = f(x) = g(h(x)) =$

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) =$

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = ?$  .
- $h'(x) = ?$  .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = ?$  .
- $h'(x) = ?$  .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$ .
- $h'(x) = ?$  .



# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$ .
- $h'(x) = ?$ .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$ .
- $h'(x) = 2x$ .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$ .
- $h'(x) = 2x$ .
- It would be nice if we could find the derivative of  $f$  in terms of the derivatives of  $y$  and  $u$ .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$ .
- $h'(x) = 2x$ .
- It would be nice if we could find the derivative of  $f$  in terms of the derivatives of  $y$  and  $u$ .
- It turns out that the derivative of the composition  $g \circ h$  is the product of the derivative of  $g$  and the derivative of  $h$ .

# The Chain Rule

- What is the derivative of  $f(x) = \sqrt{x^2 + 1}$ ?
- The Power Rule doesn't tell us how to find the derivative.
- $f$  is a composite function  $g \circ h$ :
- $y = g(u) = \sqrt{u}$ .
- $u = h(x) = x^2 + 1$ .
- Then  $y = f(x) = g(h(x)) = g(x^2 + 1) = \sqrt{x^2 + 1}$ .
- We know the derivatives of  $g$  and  $h$ :
- $g'(u) = \frac{1}{2}u^{-\frac{1}{2}}$ .
- $h'(x) = 2x$ .
- It would be nice if we could find the derivative of  $f$  in terms of the derivatives of  $y$  and  $u$ .
- It turns out that the derivative of the composition  $g \circ h$  is the product of the derivative of  $g$  and the derivative of  $h$ .
- This important fact is called the Chain Rule.

# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \qquad \text{(notation 1)} \\ \text{equivalently:}$$

# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

*equivalently:*

$$f'(x) = (g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$



# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \quad \text{(notation 1)}$$

equivalently:

$$f'(x) = (g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad \text{(notation 2)}$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad \text{(notation 3)} .$$

# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

*equivalently:*

$$f'(x) = (g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

equivalently:

$$f'(x) = (g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

# The Chain Rule

Let  $g$  and  $h$  be functions. Recall that the composite function  $f = g \circ h$  is defined via  $f(x) = g(h(x))$ .

## Theorem

*Let  $h$  be differentiable at  $x$  and let  $g$  be differentiable at  $h(x)$ . Then the composite function  $f = g \circ h$  is differentiable at  $x$  and  $f'$  is given by the product*

$$f'(x) = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

*equivalently:*

$$f'(x) = (g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

The last equality uses the Leibniz notation (due to G. Leibniz (1646-1716)).

# Chain rule notations

- As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

# Chain rule notations

- As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

- The three notations are all accepted and can be used interchangeably.

# Chain rule notations

- As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.

# Chain rule notations

- As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.
- In order to exercise ourselves we shall use all three notations throughout our course.



# Chain rule notations

- As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.
- In order to exercise ourselves we shall use all three notations throughout our course.
- There are additional notations (not covered here) used in practice.

# Chain rule notations

- As we saw, the chain rule can be written using a number of notations:

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

- The three notations are all accepted and can be used interchangeably.
- Most authors tend to prefer one of these notations over the others.
- In order to exercise ourselves we shall use all three notations throughout our course.
- There are additional notations (not covered here) used in practice.
- Whenever in doubt about derivative notation, if possible, request clarification.

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = ?$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\text{Chain Rule: } f'(x) = g'(h(x))h'(x)$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( ? \right) ( ? ) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (2x) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (2x) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } h(x) = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (2x) \\ &= \frac{x}{\sqrt{x^2 + 1}}. \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = ?$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u).$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(u).$$

$$\text{Chain Rule: } f'(x) = g'(u)u'$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= \left( ? \right) ( ? ) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= \left( \frac{1}{2\sqrt{u}} \right) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= \left( \frac{1}{2\sqrt{u}} \right) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= \left( \frac{1}{2\sqrt{u}} \right) (2x) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\begin{aligned} \text{Differentiate } f(x) &= \sqrt{x^2 + 1}. \\ \text{Let } u &= x^2 + 1. \\ \text{Let } g(u) &= \sqrt{u}. \\ \text{Then } f(x) &= g(u). \\ \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= \left( \frac{1}{2\sqrt{u}} \right) (2x) \\ &= \frac{x}{\sqrt{x^2 + 1}}. \end{aligned}$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

Differentiate  $y = \sqrt{x^2 + 1}.$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = ?$$

$$\text{Then } y = ?$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = ?$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = ?$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = \sqrt{u}.$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = \sqrt{u}.$$

$$\text{Chain Rule: } \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = \sqrt{u}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left( ? \right) \left( ? \right) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = \sqrt{u}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left( \frac{1}{2\sqrt{u}} \right) (?) \end{aligned}$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = \sqrt{u}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left( \frac{1}{2\sqrt{u}} \right) (\text{?}) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\text{Differentiate } y = \sqrt{x^2 + 1}.$$

$$\text{Let } u = x^2 + 1.$$

$$\text{Then } y = \sqrt{u}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left( \frac{1}{2\sqrt{u}} \right) (2x) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3)

$$\begin{aligned} \text{Differentiate } y &= \sqrt{x^2 + 1}. \\ \text{Let } u &= x^2 + 1. \\ \text{Then } y &= \sqrt{u}. \\ \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left( \frac{1}{2\sqrt{u}} \right) (2x) \\ &= \frac{x}{\sqrt{x^2 + 1}}. \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = ?$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\text{Chain Rule: } f'(x) = g'(h(x))h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( ? \right) ( ? ) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (\cos x) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

Example (Chain Rule, Notation 1, square root of a trigonometric function)

$$\text{Differentiate } f(x) = \sqrt{\sin x + 2}.$$

$$\text{Let } h(x) = \sin x + 2.$$

$$\text{Let } g(u) = \sqrt{u}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( \frac{1}{2\sqrt{h(x)}} \right) (\cos x) \\ &= \frac{1}{2\sqrt{\sin x + 2}}. \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = ?$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u) .$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u) .$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u) .$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$\text{Chain Rule: } f'(x) = g'(u)u'$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (\text{?}) (\text{?}) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (-\sin u) \left( ? \right) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (-\sin u) \left( ? \right) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (-\sin u) (3x^2) \end{aligned}$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos(x^3) .$$

$$\text{Let } u = x^3 .$$

$$\text{Let } g(u) = \cos u .$$

$$\text{Then } f(x) = g(u) .$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (-\sin u) (3x^2) \\ &= -3x^2 \sin(x^3) . \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate} \quad f(x) = \cos^3 x.$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = ?$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) \quad .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$\text{Chain Rule: } f'(x) = g'(u)u'$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= \left( ? \right) \left( ? \right) \end{aligned}$$



$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (3u^2) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (3u^2) (?) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (3u^2) (-\sin x) \end{aligned}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 2)

$$\text{Differentiate } f(x) = \cos^3 x.$$

$$\text{Let } u = \cos x.$$

$$\text{Let } g(u) = u^3.$$

$$\text{Then } f(x) = g(u).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(u)u' \\ &= (3u^2) (-\sin x) \\ &= -3 \sin x (\cos x)^2. \end{aligned}$$

- In the example  $y = \cos^3 x$ , the outer function was a power function:  $y = u^3$ .

- In the example  $y = \cos^3 x$ , the outer function was a power function:  $y = u^3$ .
- The derivative was  $\frac{dy}{dx} = 3u^2 \frac{du}{dx} = (3 \cos^2 x)(-\sin x)$ .

- In the example  $y = \cos^3 x$ , the outer function was a power function:  $y = u^3$ .
- The derivative was  $\frac{dy}{dx} = 3u^2 \frac{du}{dx} = (3 \cos^2 x)(-\sin x)$ .
- We can generalize this:

### Observation (The Power Rule Combined with the Chain Rule)

*If  $n$  is any real number and  $u = h(x)$  is differentiable, then*

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

*Alternatively,*

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate} \quad y = (x^3 - 1)^{100}.$$



$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = ?$$

$$\text{Then } y = ?$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = ?$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = ?$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\text{Chain Rule: } \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= \left( ? \right) \left( ? \right) \end{aligned}$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (100u^{99}) \left( ? \right) \end{aligned}$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (100u^{99}) \quad ( ? ) \end{aligned}$$



$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (100u^{99}) (3x^2) \end{aligned}$$

$$\frac{d}{dx}(u^n) = nu^{n-1} \frac{du}{dx}$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 3, Power Rule)

$$\text{Differentiate } y = (x^3 - 1)^{100}.$$

$$\text{Let } u = x^3 - 1.$$

$$\text{Then } y = u^{100}.$$

$$\begin{aligned} \text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (100u^{99}) (3x^2) \\ &= 300x^2(x^3 - 1)^{99}. \end{aligned}$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = ?$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = ?$$

$$\text{Then } f(x) = g(h(x)).$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3}) .$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\text{Chain Rule: } f'(x) = g'(h(x))h'(x)$$



$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( ? \right) ( ? ) \end{aligned}$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( -\frac{1}{3}(h(x))^{-\frac{4}{3}} \right) (?) \end{aligned}$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( -\frac{1}{3}(h(x))^{-\frac{4}{3}} \right) (?) \end{aligned}$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( -\frac{1}{3}(h(x))^{-\frac{4}{3}} \right) (2x + 1) \end{aligned}$$

$$\frac{d}{dx}[h(x)]^n = n[h(x)]^{n-1} \cdot h'(x)$$

$$(g(h(x)))' = g'(h(x)) \cdot h'(x) \quad (\text{notation 1})$$

$$(g(u))' = g'(u)u' \quad \text{where } u = h(x) \quad (\text{notation 2})$$

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} \quad \text{where } y = g(u) \quad (\text{notation 3})$$

### Example (Chain Rule, Notation 1, Power Rule)

$$\text{Differentiate } f(x) = \frac{1}{\sqrt[3]{x^2+x+1}}.$$

$$\text{Let } h(x) = x^2 + x + 1.$$

$$\text{Let } g(u) = u^{-\frac{1}{3}}.$$

$$\text{Then } f(x) = g(h(x)).$$

$$\begin{aligned} \text{Chain Rule: } f'(x) &= g'(h(x))h'(x) \\ &= \left( -\frac{1}{3}(h(x))^{-\frac{4}{3}} \right) (2x+1) \\ &= -\frac{2x+1}{3}(x^2+x+1)^{-\frac{4}{3}}. \end{aligned}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2}$$



## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2}$$

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{? \cdot (2t+1) - (t-2) \cdot ?}{(2t+1)^2}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2}$$

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{1 \cdot (2t+1) - (t-2) \cdot ?}{(2t+1)^2}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2}$$

$$= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{1 \cdot (2t+1) - (t-2) \cdot ?}{(2t+1)^2}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$\begin{aligned} &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2} \\ &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{1 \cdot (2t+1) - (t-2) \cdot 2}{(2t+1)^2} \end{aligned}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$\begin{aligned} &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2} \\ &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{1 \cdot (2t+1) - (t-2) \cdot 2}{(2t+1)^2} \\ &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{2t+1-2t+4}{(2t+1)^2} \end{aligned}$$

## Example (Chain Rule and Quotient Rule)

Find the derivative of

$$g(t) = \left( \frac{t-2}{2t+1} \right)^9.$$

Power Rule and Chain Rule:

$$g'(t) = 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{d}{dt} \left( \frac{t-2}{2t+1} \right)$$

Quotient Rule:

$$\begin{aligned} &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\frac{d}{dt}(t-2) \cdot (2t+1) - (t-2) \frac{d}{dt}(2t+1)}{(2t+1)^2} \\ &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{1 \cdot (2t+1) - (t-2) \cdot 2}{(2t+1)^2} \\ &= 9 \left( \frac{t-2}{2t+1} \right)^8 \frac{\cancel{2t} + 1 - \cancel{2t} + 4}{(2t+1)^2} = \frac{45(t-2)^8}{(2t+1)^{10}}. \end{aligned}$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .



## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$= \left( ? \right) (x^3 - x + 1)^4 + (2x + 1)^5 \left( ? \right)$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 + (2x + 1)^5 \left( ? \right)$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 + (2x + 1)^5 \left( ? \right)$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \end{aligned}$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(2x + 1)^4 (?) (x^3 - x + 1)^4 + 4(2x + 1)^5 (x^3 - x + 1)^3 (?) \end{aligned}$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(2x + 1)^4 (?) (x^3 - x + 1)^4 + 4(2x + 1)^5 (x^3 - x + 1)^3 (3x^2 - 1) \end{aligned}$$



## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(2x + 1)^4 (?) (x^3 - x + 1)^4 + 4(2x + 1)^5 (x^3 - x + 1)^3 (3x^2 - 1) \end{aligned}$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(2x + 1)^4 (2) (x^3 - x + 1)^4 + 4(2x + 1)^5 (x^3 - x + 1)^3 (3x^2 - 1) \end{aligned}$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(\mathbf{2x + 1})^4 (2) (\mathbf{x^3 - x + 1})^4 + 4(\mathbf{2x + 1})^5 (\mathbf{x^3 - x + 1})^3 (3x^2 - 1) \end{aligned}$$

Common factor  $2(\mathbf{2x + 1})^4 (\mathbf{x^3 - x + 1})^3$  :

$$= 2(2x + 1)^4 (x^3 - x + 1)^3 (2(2x + 1)(3x^2 - 1) + 5(x^3 - x + 1))$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(2x + 1)^4 (2) (x^3 - x + 1)^4 + 4(2x + 1)^5 (x^3 - x + 1)^3 (3x^2 - 1) \end{aligned}$$

Common factor  $2(2x + 1)^4(x^3 - x + 1)^3$  :

$$= 2(2x + 1)^4(x^3 - x + 1)^3(2(2x + 1)(3x^2 - 1) + 5(x^3 - x + 1))$$

## Example

Find the derivative of  $y = (2x + 1)^5(x^3 - x + 1)^4$ .

Product Rule:

$$y' = \frac{d}{dx} ((2x + 1)^5) (x^3 - x + 1)^4 + (2x + 1)^5 \frac{d}{dx} ((x^3 - x + 1)^4)$$

Chain Rule:

$$\begin{aligned} &= \left( 5(2x + 1)^4 \frac{d}{dx} (2x + 1) \right) (x^3 - x + 1)^4 \\ &\quad + (2x + 1)^5 \left( 4(x^3 - x + 1)^3 \frac{d}{dx} (x^3 - x + 1) \right) \\ &= 5(2x + 1)^4 (2) (x^3 - x + 1)^4 + 4(2x + 1)^5 (x^3 - x + 1)^3 (3x^2 - 1) \end{aligned}$$

Common factor  $2(2x + 1)^4(x^3 - x + 1)^3$  :

$$\begin{aligned} &= 2(2x + 1)^4(x^3 - x + 1)^3 (2(2x + 1)(3x^2 - 1) + 5(x^3 - x + 1)) \\ &= 2(2x + 1)^4(x^3 - x + 1)^3 (17x^3 + 6x^2 - 9x + 3) \end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left( e^? \right)^x$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$



## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left( e^{\ln 2} \right)^x$$

$$y = e^{x \ln 2}.$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = ?$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\text{Chain Rule: } \frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (\text{?})(\text{?})\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(?)\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(?)\end{aligned}$$



## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln 2)\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln 2) \\ &= \left(e^{(?) }\right)(\ln 2)\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln 2) \\ &= \left(e^{(x \ln 2)}\right)(\ln 2)\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = \left(e^{\ln 2}\right)^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln 2) \\ &= \left(e^{(x \ln 2)}\right)(\ln 2) \\ &= \left(e^{\ln 2}\right)^x (\ln 2)\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = (e^{\ln 2})^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln 2) \\ &= (e^{(x \ln 2)}) (\ln 2) \\ &= (e^{\ln 2})^x (\ln 2) \\ &= 2^x \ln 2.\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = 2^x$ .

$$y = (e^{\ln 2})^x$$

$$y = e^{x \ln 2}.$$

Let  $u = x \ln 2$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln 2) \\ &= (e^{(x \ln 2)}) (\ln 2) \\ &= (e^{\ln 2})^x (\ln 2) \\ &= 2^x \ln 2.\end{aligned}$$

## Example (Chain Rule, general exponential function)

Differentiate  $y = a^x$ .

$$y = \left(e^{\ln a}\right)^x$$

$$y = e^{x \ln a}.$$

Let  $u = x \ln a$ .

Then  $y = e^u$ .

$$\begin{aligned}\text{Chain Rule: } \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dx} \\ &= (e^u)(\ln a) \\ &= \left(e^{(x \ln a)}\right)(\ln a) \\ &= \left(e^{\ln a}\right)^x (\ln a) \\ &= a^x \ln a.\end{aligned}$$

## Theorem (The Derivative of $a^x$ )

$$\frac{d}{dx}(a^x) = a^x \ln a.$$



- We can add more “links” when we use the Chain Rule.

- We can add more “links” when we use the Chain Rule.
- $y = f(u)$
- $u = g(x)$
- $x = h(t)$

- We can add more “links” when we use the Chain Rule.
- $y = f(u)$
- $u = g(x)$
- $x = h(t)$
- Use the Chain Rule twice:

$$\frac{dy}{dt} = \frac{dy}{du} \frac{du}{dt}$$

- We can add more “links” when we use the Chain Rule.
- $y = f(u)$
- $u = g(x)$
- $x = h(t)$
- Use the Chain Rule twice:

$$\frac{dy}{dt} = \frac{dy}{du} \frac{du}{dt} = \frac{dy}{du} \frac{du}{dx} \frac{dx}{dt}$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( ? \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$



## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \left( ? \right) \frac{d}{dx} (10^x + 1)$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) \frac{d}{dx} (10^x + 1)$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) \frac{d}{dx} (10^x + 1)$

$$= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) ( ? )$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) \frac{d}{dx} (10^x + 1)$

$$= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) (10^x \ln 10)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = \sin \sqrt{10^x + 1}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( \sin \sqrt{10^x + 1} \right)$$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \frac{d}{dx} \left( \sqrt{10^x + 1} \right)$

Chain Rule:  $= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) \frac{d}{dx} (10^x + 1)$

$$= \left( \cos \sqrt{10^x + 1} \right) \left( \frac{1}{2\sqrt{10^x + 1}} \right) (10^x \ln 10)$$

$$= \frac{(\ln 10) 10^x \cos \sqrt{10^x + 1}}{2\sqrt{10^x + 1}}.$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .



## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

$$\text{Chain Rule:} \quad = \left( ? \right) \frac{d}{dx} \left( \tan(\pi x) \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

$$\text{Chain Rule:} \quad = \left( e^{\tan(\pi x)} \right) \frac{d}{dx} \left( \tan(\pi x) \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

$$\text{Chain Rule:} \quad = \left( e^{\tan(\pi x)} \right) \frac{d}{dx} \left( \tan(\pi x) \right)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \frac{d}{dx} (\tan(\pi x))$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \left( ? \right) \frac{d}{dx} (\pi x)$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

$$\text{Chain Rule:} \quad = \left( e^{\tan(\pi x)} \right) \frac{d}{dx} \left( \tan(\pi x) \right)$$

$$\text{Chain Rule:} \quad = \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) \frac{d}{dx} (\pi x)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \frac{d}{dx} \left( \tan(\pi x) \right)$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) \frac{d}{dx} (\pi x)$

$$= \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) (?)$$



## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \frac{d}{dx} \left( \tan(\pi x) \right)$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) \frac{d}{dx} (\pi x)$

$$= \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) (\pi)$$

## Example (Using the Chain Rule twice)

Differentiate:  $y = e^{\tan(\pi x)}$ .

$$\frac{dy}{dx} = \frac{d}{dx} \left( e^{\tan(\pi x)} \right)$$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \frac{d}{dx} \left( \tan(\pi x) \right)$

Chain Rule:  $= \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) \frac{d}{dx} (\pi x)$

$$= \left( e^{\tan(\pi x)} \right) \left( \sec^2(\pi x) \right) (\pi)$$
$$= \pi e^{\tan(\pi x)} \sec^2(\pi x).$$

## Theorem (Chain rule)

*Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .*

$$(f(g(x)))'_{|x=a} = f'(g(a))g'(a)$$

Proof with additional assumptions -motivation for actual proof.



## Theorem (Chain rule)

*Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .*

$$(f(g(x)))'_{|x=a} = f'(g(a))g'(a)$$

Proof with **additional assumptions** -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ .



## Theorem (Chain rule)

*Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .*

$$(f(g(x)))'_{|x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ .

$$(f \circ g)'(a) =$$



## Theorem (Chain rule)

*Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .*

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ .

$$(f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ .

$$\begin{aligned}(f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right)\end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ .

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \end{aligned}$$





## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ .

$$\begin{aligned}(f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\&= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\&= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right)\end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y)-f(g(a))}{y-g(a)}$ .

$$\begin{aligned}(f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\&= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\&= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right)\end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y)-f(g(a))}{y-g(a)}$ .

$G(y)$  is continuous at  $g(a) \Rightarrow$

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y)-f(g(a))}{y-g(a)}$ .  
 $G(y)$  is continuous at  $g(a) \Rightarrow G(g(x))$  is continuous at  $a$ .

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y) - f(g(a))}{y - g(a)}$ .  
 $G(y)$  is continuous at  $g(a) \Rightarrow G(g(x))$  is continuous at  $a$ .

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \left( \lim_{y=g(x), y \rightarrow g(a)} \frac{f(y) - f(g(a))}{y - g(a)} \right) g'(a) = \end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y)-f(g(a))}{y-g(a)}$ .  
 $G(y)$  is continuous at  $g(a) \Rightarrow G(g(x))$  is continuous at  $a$ . **Furthermore  $g(x)$  is continuous at  $a$ .**

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(\textcolor{red}{g(x)}) - f(g(a))}{\textcolor{red}{g(x)} - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \left( \lim_{y=g(x), y \rightarrow g(a)} \frac{f(\textcolor{red}{y}) - f(g(a))}{\textcolor{red}{y} - g(a)} \right) g'(a) = \end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y) - f(g(a))}{y - g(a)}$ .  $G(y)$  is continuous at  $g(a) \Rightarrow G(g(x))$  is continuous at  $a$ . Furthermore  $g(x)$  is continuous at  $a$ .

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \left( \lim_{y=g(x), y \rightarrow g(a)} \frac{f(y) - f(g(a))}{y - g(a)} \right) g'(a) = f'(g(a))g'(a) . \end{aligned}$$



## Theorem (Chain rule)

Let  $g$ -differentiable at neighborhood of  $a$ ,  $f$ -diff. at neighb. of  $g(a)$ .

$$(f(g(x)))'|_{x=a} = f'(g(a))g'(a)$$

## Proof with additional assumptions -motivation for actual proof.

Suppose that  $g(x) \neq g(a)$  so long as  $x \neq a$ . Set  $G(y) = \frac{f(y) - f(g(a))}{y - g(a)}$ .  $G(y)$  is continuous at  $g(a) \Rightarrow G(g(x))$  is continuous at  $a$ . Furthermore  $g(x)$  is continuous at  $a$ .

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \lim_{x \rightarrow a} \left( \frac{f(g(x)) - f(g(a))}{g(x) - g(a)} \right) \lim_{x \rightarrow a} \left( \frac{g(x) - g(a)}{x - a} \right) \\ &= \left( \lim_{y=g(x), y \rightarrow g(a)} \frac{f(y) - f(g(a))}{y - g(a)} \right) g'(a) = f'(g(a))g'(a) \end{aligned}$$





## Theorem (Chain rule)

*$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a) \Rightarrow (f(g(a)))' = f'(g(a))g'(a)$  .*

Proof.



## Theorem (Chain rule)

*$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a) \Rightarrow (f(g(a)))' = f'(g(a))g'(a)$  .*

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$  .



## Theorem (Chain rule)

*$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a)$   $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$  .*

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$  .  $Q(g(x))$  - defined for all  $x$  near  $a$ .



## Theorem (Chain rule)

$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a) \Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .



## Theorem (Chain rule)

*$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a)$   $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .*

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .



## Theorem (Chain rule)

*$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a)$   $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .*

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .



## Theorem (Chain rule)

*$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a)$   $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .*

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$Q(g(x)) \frac{g(x)-g(a)}{x-a} =$$



## Theorem (Chain rule)

$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a) \Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$Q(g(x)) \frac{g(x)-g(a)}{x-a} = \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a}, & g(x) = g(a) \end{cases}$$





## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g(a)*  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$Q(g(x)) \frac{g(x)-g(a)}{x-a} = \begin{cases} \frac{(f(g(x))-f(g(a)))(g(x)-g(a))}{(g(x)-g(a))(x-a)}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases}$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g(a)*  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a))) \cancel{(g(x)-g(a))}}{\cancel{(g(x)-g(a))} x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g(a)*  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g(a)*  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$(f \circ g)'(a) =$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g*(*a*)  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$(f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} =$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g*(*a*)  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$(f \circ g)'(a) = \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} Q(g(x)) \frac{g(x) - g(a)}{x - a}$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g*(*a*)  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(a)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} Q(g(x)) \frac{g(x) - g(a)}{x - a} \\ &= \lim_{x \rightarrow a} Q(g(x)) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} \end{aligned}$$



## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g(a)*  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(x)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} Q(g(x)) \frac{g(x) - g(a)}{x - a} \\ &= \lim_{x \rightarrow a} Q(g(x)) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} = f'(g(a))g'(a). \end{aligned}$$





## Theorem (Chain rule)

*g*-diff. near *a*, *f*-diff. near *g(a)*  $\Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(x)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} Q(g(x)) \frac{g(x) - g(a)}{x - a} \\ &= \lim_{x \rightarrow a} Q(g(x)) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} = f'(g(a))g'(a). \end{aligned}$$



## Theorem (Chain rule)

$g$ -diff. near  $a$ ,  $f$ -diff. near  $g(a) \Rightarrow (f(g(a)))' = f'(g(a))g'(a)$ .

### Proof.

Define  $Q(y) = \begin{cases} \frac{f(y)-f(g(a))}{y-g(a)}, & y \neq g(a) \\ f'(g(a)), & y = g(a) \end{cases}$ .  $Q(g(x))$  - defined for all  $x$  near  $a$ . Therefore  $f'(g(a)) = \lim_{y \rightarrow a} Q(y) = \lim_{x \rightarrow a} Q(g(x))$ .

$$\begin{aligned} Q(g(x)) \frac{g(x)-g(a)}{x-a} &= \begin{cases} \frac{(f(g(x))-f(g(a)))}{(g(x)-g(a))} \frac{(g(x)-g(a))}{x-a}, & g(x) \neq g(a) \\ f'(g(a)) \frac{g(x)-g(a)}{x-a} = 0, & g(x) = g(a) \end{cases} \\ &= \frac{f(g(x))-f(a)}{x-a}. \end{aligned}$$

$$\begin{aligned} (f \circ g)'(a) &= \lim_{x \rightarrow a} \frac{f(g(x)) - f(g(a))}{x - a} = \lim_{x \rightarrow a} Q(g(x)) \frac{g(x) - g(a)}{x - a} \\ &= \lim_{x \rightarrow a} Q(g(x)) \lim_{x \rightarrow a} \frac{g(x) - g(a)}{x - a} = f'(g(a))g'(a). \end{aligned}$$

