

Calculus I

Lecture 0

Representing Functions

Todor Milev

<https://github.com/tmilev/freecalc>

2020

Outline

- 1 Ways to Represent a Function
 - The Definition of a Function
 - The Vertical Line Test
 - Piecewise Defined Functions
 - Symmetry
 - Increasing and Decreasing Functions
 - A Note on Domains of Functions

License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
<https://creativecommons.org/licenses/by/3.0/us/>
and the links therein.


License to use and redistribute

These lecture slides and their \LaTeX source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:
`https://github.com/tmilev/freecalc`
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:
`https://creativecommons.org/licenses/by/3.0/us/`
and the links therein.




Definition (Function)

A function f is a rule that assigns to each element x in a set D exactly one element, called $f(x)$, in a set E .

- Functions are also synonymously called “maps”.

Definition (Domain)

The set D in the definition of f is called the domain of f .




Definition (Function)

A function f is a rule that assigns to each element x in a set D exactly one element, called $f(x)$, in a set E .

Definition (Co-domain)

The set E in the definition of f is called the co-domain of f .




Definition (Function)

A function f is a rule that assigns to each element x in a set D exactly one element, called $f(x)$, in a set E .

Definition (Value of f at x)

The number $f(x)$ is called *the value of f at x* and is read " f of x ".

- The value of f at x is also called the image of x under the map f .



Definition (Function)

A function f is a rule that assigns to each element x in a set D exactly one element, called $f(x)$, in a set E .

Definition (Range)

The set of all possible values taken by $f(x)$ as the element x runs over elements of D is called the range of f .

The Vertical Line Test


Question

Given a curve in the plane, is it the graph of a function or not?

The answer is as follows.

Proposition (The Vertical Line Test)

A curve in the plane is the graph of a function if and only if no vertical line intersects it more than once.




Piecewise Defined Functions

Definition (Piecewise Defined Function)

A piecewise defined function is a function that is defined by different algebraic formulas on different subsets of its domain.

Example



$$f(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ -1 & \text{if } x < 0 \end{cases}$$

The filled red circle means $(0, 1)$ is on the curve.


The open circle means $(0, -1)$ is not on the curve.

Example

The absolute value $|x|$ of a number a is defined to be


$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$

Sketch a graph of the function $f(x) = |x|$.



Example

Find a formula for the function f whose graph is given below.




Different formulas on $[0, 1)$, $[1, 2)$, and $[2, 5)$.

$$f(x) = \begin{cases} x & \text{if } 0 \leq x < 1 \\ 2 - x & \text{if } 1 \leq x < 2 \\ 0 & \text{if } 2 \leq x < 5 \end{cases}$$

Example

Sketch the function $f(x) = |2x - 3|$.



$$|x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0. \end{cases}$$


$$|2x - 3| = \begin{cases} 2x - 3 & \text{if } 2x - 3 \geq 0 \\ -(2x - 3) & \text{if } 2x - 3 < 0 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } 2x \geq 3 \\ -2x + 3 & \text{if } 2x < 3 \end{cases}$$

$$= \begin{cases} 2x - 3 & \text{if } x \geq 3/2 \\ -2x + 3 & \text{if } x < 3/2. \end{cases}$$

Example

Sketch the function $f(x) = \frac{|4x + 2|}{2x + 1}$.



$$|u| = \begin{cases} u & \text{if } u \geq 0 \\ -u & \text{if } u < 0. \end{cases}$$

$$\frac{|4x + 2|}{2x + 1} = \begin{cases} \frac{4x+2}{2x+1} & \text{if } 4x + 2 > 0 \\ \frac{-(4x+2)}{2x+1} & \text{if } 4x + 2 < 0 \end{cases}$$

$$= \begin{cases} \frac{2(2x+1)}{2x+1} & \text{if } 4x > -2 \\ \frac{-2(2x+1)}{2x+1} & \text{if } 4x < -2 \end{cases}$$

$$= \begin{cases} 2 & \text{if } x > -\frac{1}{2} \\ -2 & \text{if } x < -\frac{1}{2}. \end{cases}$$

Symmetry

Definition (Even and Odd Functions)

A function f is called even if $f(-x) = f(x)$ for all x in its domain. A function f is called odd if $f(-x) = -f(x)$ for all x in its domain.

Example (x^2 is Even, x^3 is Odd)

The function $f(x) = x^2$ is even:

$$f(-x) = (-x)^2 = x^2 = f(x).$$

The function $g(x) = x^3$ is odd:

$$g(-x) = (-x)^3 = -x^3 = -g(x).$$

Definition (Even and Odd Functions)

A function f is called even if $f(-x) = f(x)$ for all x in its domain. A function f is called odd if $f(-x) = -f(x)$ for all x in its domain.

Example

Determine whether each of the following functions is even, odd, or neither even nor odd.

$$f(x) = x^5 + x$$

$$\begin{aligned} f(-x) &= (-x)^5 + (-x) \\ &= -x^5 - x \\ &= -(x^5 + x) \\ &= -f(x) \end{aligned}$$

Therefore f is odd.

$$g(x) = 1 - x^4$$

$$\begin{aligned} g(-x) &= 1 - (-x)^4 \\ &= 1 - x^4 \\ &= g(x) \end{aligned}$$

Therefore g is even.

$$h(x) = 2x - 1$$

$$\begin{aligned} h(-x) &= 2(-x) - 1 \\ &= -2x - 1 \\ &\neq h(x), -h(x) \end{aligned}$$

Therefore h is neither even nor odd.


Increasing and Decreasing Functions

Definition (Increasing and Decreasing Functions)

A function f is called increasing on an interval I if $f(x_1) < f(x_2)$ whenever $x_1 < x_2$ in I .

It is called decreasing on the interval I if $f(x_1) > f(x_2)$ whenever $x_1 < x_2$ in I .

Example (Increasing and Decreasing)



- f is increasing on $[-1, -\frac{1}{2}]$.
- f is decreasing on $[-\frac{1}{2}, \frac{1}{2}]$.
- f is increasing on $[\frac{1}{2}, 1]$.

A Note on Domains of Functions

If the domain of a function isn't specified, it is implied to be all numbers x for which the formula $f(x)$ is defined. There are some restrictions to consider:

- Can't divide by 0.
- Even roots of a negative number are not defined in this course ($\sqrt{-1}$, $\sqrt[4]{-2053}$, $\sqrt[6]{-15} \dots$ not allowed).
- Taking $\log x$ if $x \leq 0$ is not allowed in this course; taking $\log 0$ is not allowed in any course.