

# Precalculus

## Lecture 16

### Factoring Polynomials

Todor Milev

<https://github.com/tmilev/freecalc>

2020

# Outline

## 1 Factorization overview

# Outline

- 1 Factorization overview
- 2 Polynomial division

# Outline

- 1 Factorization overview
- 2 Polynomial division
- 3 Factoring cubics with rational root

# Outline

- 1 Factorization overview
- 2 Polynomial division
- 3 Factoring cubics with rational root
- 4 Polynomial inequalities

# License to use and redistribute

These lecture slides and their  $\text{\LaTeX}$  source code are licensed to you under the Creative Commons license CC BY 3.0. You are free

- to Share - to copy, distribute and transmit the work,
- to Remix - to adapt, change, etc., the work,
- to make commercial use of the work,

as long as you reasonably acknowledge the original project.

- Latest version of the .tex sources of the slides:  
<https://github.com/tmilev/freecalc>
- Should the link be outdated/moved, search for “freecalc project”.
- Creative Commons license CC BY 3.0:  
<https://creativecommons.org/licenses/by/3.0/us/>  
and the links therein.

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 =$$

$$x^2 + 1 =$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (\quad)(\quad)$$

$$x^2 + 1 =$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$



Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1)$$

$$x^2 + 1 =$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 =$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1)$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left( x - \left( -\frac{5}{2} \right) \right) (x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x+i)(x-i)$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^{\textcolor{red}{2}} - i^{\textcolor{red}{2}} = (x + i)(x - i)$$

$$x^4 - 1 =$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1)$$

$$x^4 + 1 =$$



Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x+i)(x-i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x+i)(x-i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left( x - \left( -\frac{5}{2} \right) \right) (x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 = (\textcolor{red}{x}^2 - 1)(x^2 + 1) = (\textcolor{red}{x} - 1)(\textcolor{red}{x} + 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$x^4 + 1 =$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x+i)(x-i)$$

$$\begin{aligned} x^4 - 1 &= (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1) \\ &= (x - 1)(x + 1)(x - i)(x + i) \end{aligned}$$

$$x^4 + 1 =$$



Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$\begin{aligned} x^4 - 1 &= (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1) \\ &= (x - 1)(x + 1)(x - i)(x + i) \end{aligned}$$

$$x^4 + 1 = \left(x^2 - \sqrt{2}x + 1\right) \left(x^2 + \sqrt{2}x + 1\right)$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$\begin{aligned} x^4 - 1 &= (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1) \\ &= (x - 1)(x + 1)(x - i)(x + i) \end{aligned}$$

$$\begin{aligned} x^4 + 1 &= \left(x^2 - \sqrt{2}x + 1\right) \left(x^2 + \sqrt{2}x + 1\right) \\ &= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right) \\ &\quad \left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right) \end{aligned}$$

Recall that  $i^2 = -1$ ,  $\sqrt{-1} = i$ .

## Example (Polynomial factorizations)

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2 \left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x + i)(x - i)$$

$$\begin{aligned} x^4 - 1 &= (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1) \\ &= (x - 1)(x + 1)(x - i)(x + i) \end{aligned}$$

$$\begin{aligned} x^4 + 1 &= \left(x^2 - \sqrt{2}x + 1\right) \left(x^2 + \sqrt{2}x + 1\right) \\ &= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right) \\ &\quad \left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right) \end{aligned}$$

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- Every pol. of deg.  $n$  can be factored as **product of  $n$  linear factors**.

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- Every pol. of **deg.  $n$**  can be factored as product of  $n$  linear factors.

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- Every pol. of deg.  $n$  can be factored as product of  $n$  linear factors.

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- Every pol. of deg.  $n$  can be factored as product of  $n$  linear factors.
- $x_1, \dots, x_n$  may be complex numbers. Reminder: complex numbers are of the form  $p + qi$ , where  $i^2 = -1$  and  $\sqrt{-1} = i$ .



## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- Every pol. of deg.  $n$  can be factored as product of  $n$  linear factors.
- $x_1, \dots, x_n$  may be complex numbers. Reminder: complex numbers are of the form  $p + qi$ , where  $i^2 = -1$  and  $\sqrt{-1} = i$ .
- While we can find  $x_1, \dots, x_n$  with arbitrary precision, **there may not exist a formula involving radicals for computing each  $x_1, \dots, x_n$ .**

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- Every pol. of deg.  $n$  can be factored as product of  $n$  linear factors.
- $x_1, \dots, x_n$  may be complex numbers. **Reminder: complex numbers are of the form  $p + qi$ , where  $i^2 = -1$  and  $\sqrt{-1} = i$ .**
- While we can find  $x_1, \dots, x_n$  with arbitrary precision, there may not exist a formula involving radicals for computing each  $x_1, \dots, x_n$ .

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- $x_1, \dots, x_n$  may be complex numbers. Reminder: complex numbers are of the form  $p + qi$ , where  $i^2 = -1$  and  $\sqrt{-1} = i$ .

## Theorem (The Fundamental Theorem of Algebra)

*Every polynomial can be factored into product of linear terms*

$$p(x) = a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n),$$

*where  $x_1, \dots, x_n$  are the (not necessarily different) roots of  $p(x)$ .*

- $x_1, \dots, x_n$  may be complex numbers. Reminder: complex numbers are of the form  $p + qi$ , where  $i^2 = -1$  and  $\sqrt{-1} = i$ .

## Corollary

*Every real polynomial can be factored into a product of real linear terms and real quadratic terms with no real roots, i.e., factors of form*

- $(x - r)$ , where  $r$  is real and
- $ax^2 + bx + c$  with  $b^2 - 4ac < 0$  where  $a, b, c$  are real.

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1) \quad \left| \text{real roots} \right.$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1) \quad \left| \text{real roots}\right.$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1) \quad \left| \begin{array}{l} \text{real roots} \end{array} \right.$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i) \quad \left| \begin{array}{l} \text{complex roots} \end{array} \right.$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$



$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1) \quad \left| \begin{array}{l} \text{real roots} \end{array} \right.$$

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i) \quad \left| \begin{array}{l} \text{complex roots} \end{array} \right.$$

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

real roots

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

complex roots

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

mixed roots

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

real roots

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

complex roots

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

mixed roots

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2\left(x - \left(-\frac{5}{2}\right)\right)(x - 1)$$

real roots

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

complex roots

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

mixed roots

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

complex roots

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

=prod. real quadratics no roots & lin. terms.

## Example

$$2x^2 + 3x - 5 = (2x + 5)(x - 1) = 2(x - (-\frac{5}{2}))(x - 1)$$

real roots

$$x^2 + 1 = x^2 - (-1) = x^2 - i^2 = (x - (-i))(x - i)$$

complex roots

$$x^4 - 1 = (x^2 - 1)(x^2 + 1) = (x - 1)(x + 1)(x^2 + 1)$$

$$= (x - 1)(x - (-1))(x - i)(x - (-i))$$

mixed roots

$$x^4 + 1 = (x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$$

$$= \left(x - \left(\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

$$\left(x - \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right)\right) \left(x - \left(-\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)\right)$$

complex roots

# Factoring polynomials in practice

- In theory every polynomial can be factored.

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

# Factoring polynomials in practice

- In theory every polynomial can be factored.

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- Theory guarantees numerical approximations for roots  $x_1, \dots, x_n$ .

# Factoring polynomials in practice

- In theory every polynomial can be factored.

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- Theory guarantees numerical approximations for roots  $x_1, \dots, x_n$ .
- **Can we find algebraic formulas** for  $x_1, \dots, x_n$ ?



# Factoring polynomials in practice

- In theory every polynomial can be factored.

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- Theory guarantees numerical approximations for roots  $x_1, \dots, x_n$ .
- **Can we find algebraic formulas** for  $x_1, \dots, x_n$ ?
- **No**, if using finitely many operations  $+$ ,  $-$ ,  $*$ ,  $/$ ,  $\sqrt[n]{\phantom{x}}$ .

# Factoring polynomials in practice

- In theory every polynomial can be factored.

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- Theory guarantees numerical approximations for roots  $x_1, \dots, x_n$ .
- Can we find algebraic formulas for  $x_1, \dots, x_n$ ?
- No, if using finitely many operations  $+$ ,  $-$ ,  $*$ ,  $/$ ,  $\sqrt[n]{\phantom{x}}$ .
- First (advanced) proof by Norwegian Niels Henrik Abel(1824) based on work of Italian Paolo Ruffini(1799).

# Factoring polynomials in practice

- In theory every polynomial can be factored.

$$a_0x^n + a_1x^{n-1} + \dots + a_n = a_0(x - x_1) \dots (x - x_n)$$

- Theory guarantees numerical approximations for roots  $x_1, \dots, x_n$ .
- **Can we find algebraic formulas** for  $x_1, \dots, x_n$ ?
- No, if using finitely many operations  $+, -, *, /$ ,  $\sqrt[n]{\phantom{x}}$ .
- First (advanced) proof by Norwegian Niels Henrik Abel(1824) based on work of Italian Paolo Ruffini(1799).
- **Yes**, with extra operations. Difficult: google Galois Theory to get started.

# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:

These poly's are equal	Type of factorization
$x^4 + 1$	
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right)$ $\left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	

# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:
  - Factor the polynomial over the rational numbers. Use integers/quotients, but no  $\sqrt{\phantom{x}}$ .

These poly's are equal	Type of factorization
$x^4 + 1$	factored over rationals
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right)$ $\left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	

# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:
  - Factor the polynomial **over the rational numbers**. Use integers/quotients, but no  $\sqrt{\phantom{x}}$ .

These poly's are equal	Type of factorization
$x^4 + 1$	<b>factored over rationals</b>
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	

# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:
  - Factor the polynomial over the rational numbers. Use integers/quotients, but no  $\sqrt{\phantom{x}}$ .
  - Factor the polynomial over the real numbers. Use radicals and/or numerical approximations, no use of  $i = \sqrt{-1}$ .

These poly's are equal	Type of factorization
$x^4 + 1$	factored over rationals
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	factored over the reals
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right)$ $\left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	

# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:
  - Factor the polynomial over the rational numbers. Use integers/quotients, but no  $\sqrt{\phantom{x}}$ .
  - Factor the polynomial **over the real numbers**. Use radicals and/or numerical approximations, **no use of  $i = \sqrt{-1}$** .

These poly's are equal	Type of factorization
$x^4 + 1$	factored over rationals
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	<b>factored over the reals</b>
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right)$ $\left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	



# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:
  - Factor the polynomial over the rational numbers. Use integers/quotients, but no  $\sqrt{\phantom{x}}$ .
  - Factor the polynomial over the real numbers. Use radicals and/or numerical approximations, no use of  $i = \sqrt{-1}$ .
  - Fully factor the polynomial using complex numbers.

These poly's are equal	Type of factorization
$x^4 + 1$	factored over rationals
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	factored over the reals
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right)$ $\left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	full complex factorization

# What does factorization mean?

- Based on context, “to factor a polynomial” means one of:
  - Factor the polynomial over the rational numbers. Use integers/quotients, but no  $\sqrt{\phantom{x}}$ .
  - Factor the polynomial over the real numbers. Use radicals and/or numerical approximations, no use of  $i = \sqrt{-1}$ .
  - Fully factor the polynomial **using complex numbers**.

These poly's are equal	Type of factorization
$x^4 + 1$	factored over rationals
$(x^2 - \sqrt{2}x + 1)(x^2 + \sqrt{2}x + 1)$	factored over the reals
$\left( x - \left( \frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i \right) \right)$ $\left( x - \left( \frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right) \left( x - \left( -\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i \right) \right)$	full complex factorization

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.
  - Kronecker algorithm (German Leopold Kronecker (1823-1891)).

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.
  - Kronecker algorithm (German Leopold Kronecker (1823-1891)).
  - Methods based on finite fields.



# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.
  - Kronecker algorithm (German Leopold Kronecker (1823-1891)).
  - Methods based on finite fields.
  - Lenstra-Lenstra-Lovász algorithm (Dutch, Dutch, Hungarian mathematicians, all contemporary).

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.
  - Kronecker algorithm (German Leopold Kronecker (1823-1891)).
  - Methods based on finite fields.
  - Lenstra-Lenstra-Lovász algorithm (Dutch, Dutch, Hungarian mathematicians, all contemporary).
- **Above methods require computer**; no rational roots assumption.

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.
  - Kronecker algorithm (German Leopold Kronecker (1823-1891)).
  - Methods based on finite fields.
  - Lenstra-Lenstra-Lovász algorithm (Dutch, Dutch, Hungarian mathematicians, all contemporary).
- Above methods require computer; **no rational roots assumption**.
- **If we assume rational roots** there are practical algorithms **by hand**.

# Factorization over the rationals

- Suppose we want to factor a polynomial using only rational numbers (no  $\sqrt[n]{\phantom{x}}$  or numerical approximations).
- No guarantee to get:

$$a_0x^n + a_1x^{n-1} + \cdots + a_n = a_0(x - x_1) \cdots (x - x_n)$$

- A factorization using rationals may have arbitrarily large factors.
- Efficient algorithms for factoring using rationals exist.
  - Kronecker algorithm (German Leopold Kronecker (1823-1891)).
  - Methods based on finite fields.
  - Lenstra-Lenstra-Lovász algorithm (Dutch, Dutch, Hungarian mathematicians, all contemporary).
- Above methods require computer; no rational roots assumption.
- If we assume rational roots there are practical algorithms by hand.
- We study those for cubics with the aid of scientific calculator.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1}$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x + 1} + 1}$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \text{?} \\
 x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1}
 \end{array}$$

Divide  $x^3$  by  $x$ .



## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r} x^2 \\ x - 1 \overline{) x^3 + 2x^2 + 1} \end{array}$$

Divide  $x^3$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1} \\
 \underline{\phantom{x^3} ? \phantom{+ 0x} ?} \\
 \phantom{x^3 + 2x^2 + 0x}
 \end{array}$$

Multiply  $x^2$  by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 \phantom{+ 0x} + 1} \\
 \underline{x^3 - x^2} \phantom{+ 0x} \\
 \phantom{x^3} 3x^2 \phantom{+ 0x} + 1
 \end{array}$$

Multiply  $x^2$  by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 \phantom{x^3} 3x^2 \phantom{+ 1} \\
 \phantom{x^3} \phantom{3x^2} \phantom{+ 1} \phantom{+ 1}
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \phantom{x^3} x^2 \quad ? \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1}
 \end{array}$$

Divide  $3x^2$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1}
 \end{array}$$

Divide  $3x^2$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{\phantom{3}x^2 \phantom{+ 1}} \\
 \phantom{3}x^2 \phantom{+ 1}
 \end{array}$$

Multiply  $3x$  by divisor.



## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \underline{3x^2 - 3x} \\
 \phantom{3x^2} 4x + 1
 \end{array}$$

Multiply  $3x$  by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 \phantom{3x^2} 3x \phantom{+ 1} \\
 \phantom{3x^2} \underline{3x - 3} \phantom{+ 1} \\
 \phantom{3x^2} \phantom{3x} 4 \phantom{+ 1}
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \\
 3x + 1
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x \quad ? \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1
 \end{array}$$

Divide  $3x$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \underline{3x^2 - 3x} \\
 3x + 1
 \end{array}$$

Divide  $3x$  by  $x$ .

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{\phantom{3x} ? \phantom{+ 1}}
 \end{array}$$

Multiply 3 by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 + 1 \\
 \underline{3x^2 - 3x} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

Multiply 3 by divisor.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 ?
 \end{array}$$

Subtract last two polynomials.



## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

Subtract last two polynomials.

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \text{Quotient: } x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(x^3 + 2x^2 + 1) = (x^2 + 3x + 3) \cdot (x - 1) + 4$$

## Example (Polynomial long division)

Divide with quotient and remainder  $x^3 + 2x^2 + 1$  by  $x - 1$ .

$$\begin{array}{r}
 \text{Quotient: } x^2 + 3x + 3 \\
 x - 1 \overline{) x^3 + 2x^2 + 1} \\
 \underline{x^3 - x^2} \phantom{+ 1} \\
 3x^2 \phantom{+ 1} \\
 \underline{3x^2 - 3x} \phantom{+ 1} \\
 3x + 1 \\
 \underline{3x - 3} \\
 4
 \end{array}$$

**Remainder:** 4

$$\begin{aligned}
 (\text{Dividend}) &= (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder}) \\
 (x^3 + 2x^2 + 1) &= (x^2 + 3x + 3) \cdot (x - 1) + 4
 \end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r} \text{?} \\ 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \end{array}$$

Divide  $6x^3$  by  $2x$ .



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r} 3x^2 \\ 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \end{array}$$

Divide  $6x^3$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r} 3x^2 \\ 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\ \underline{\phantom{2x - 3} ? \phantom{00} ? \phantom{00}} \end{array}$$

Multiply  $3x^2$  by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r} 3x^2 \\ 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\ \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \end{array}$$

Multiply  $3x^2$  by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 10x^2 + 17x - 3
 \end{array}$$

?      ?      ?

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 \quad ? \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3
 \end{array}$$

Divide  $-10x^2$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3
 \end{array}$$

Divide  $-10x^2$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \phantom{-} \underline{\phantom{-10x^2 + } ? \phantom{+ 17x - 3}}
 \end{array}$$

Multiply  $-5x$  by divisor.



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Multiply  $-5x$  by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 \phantom{-10x^2 + } 2x - 3 \\
 \phantom{-10x^2 + } \underline{2x - 3} \\
 \phantom{-10x^2 + } 0
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x \quad ? \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \\
 2x - 3
 \end{array}$$

Divide  $2x$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3
 \end{array}$$

Divide  $2x$  by  $2x$ .

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{\phantom{2x} ? \phantom{0} ?}
 \end{array}$$

Multiply **1** by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

Multiply **1** by divisor.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 ?
 \end{array}$$

Subtract last two polynomials.



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

Subtract last two polynomials.

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 \text{Quotient: } 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 \text{Quotient:} \quad 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

**Remainder:** 0

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. **Use your work to factor the cubic.** Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{array}{r}
 \text{Quotient:} \quad 3x^2 - 5x + 1 \\
 2x - 3 \overline{) 6x^3 - 19x^2 + 17x - 3} \\
 \underline{6x^3 - 9x^2} \phantom{+ 17x - 3} \\
 -10x^2 + 17x - 3 \\
 \underline{-10x^2 + 15x} \phantom{- 3} \\
 2x - 3 \\
 \underline{2x - 3} \\
 0
 \end{array}$$

**Remainder:** 0

$$(\text{Dividend}) = (\text{Quotient}) \cdot (\text{Divisor}) + (\text{Remainder})$$

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$(6x^3 - 19x^2 + 17x - 3) = (3x^2 - 5x + 1) \cdot (2x - 3)$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = ?$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - ? \right) \left( x - ? \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5 + \sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5 - \sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. **Use your work to factor the cubic.** Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5 + \sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5 - \sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. **Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .**

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5 + \sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5 - \sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$6x^3 - 19x^2 + 17x - 3 = 0$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right)\end{aligned}$$



## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right)\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right)\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 &= 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right) \\ x &= \frac{3}{2}\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.

$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ \textcolor{red}{2}x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right) \\ x &= \textcolor{red}{\frac{3}{2}}\end{aligned}$$

## Example

Demonstrate that  $6x^3 - 19x^2 + 17x - 3$  is divisible by  $2x - 3$  using polynomial long division. Use your work to factor the cubic. Solve the equation  $6x^3 - 19x^2 + 17x - 3 = 0$ .

$$\begin{aligned}(6x^3 - 19x^2 + 17x - 3) &= (3x^2 - 5x + 1) \cdot (2x - 3) \\ &= 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3)\end{aligned}$$

No easy factorization of quadratic, so use formula:

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-(-5) \pm \sqrt{(-5)^2 - 4 \cdot 3 \cdot 1}}{2 \cdot 3} = \frac{5 \pm \sqrt{13}}{6}$$

We are ready to solve the equation.


$$\begin{aligned}6x^3 - 19x^2 + 17x - 3 &= 0 \\ 3 \left( x - \left( \frac{5+\sqrt{13}}{6} \right) \right) \left( x - \left( \frac{5-\sqrt{13}}{6} \right) \right) (2x - 3) &= 0 \\ 2x - 3 = 0 \quad \text{or} \quad x = \left( \frac{5+\sqrt{13}}{6} \right) \quad \text{or} \quad x = \left( \frac{5-\sqrt{13}}{6} \right) \\ x &= \frac{3}{2}\end{aligned}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s).

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the x axis at: ?



## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the x axis at:  $-1.5, -1, 2$ .

## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$?(x - ?)(x - ?)(x - ?)$$

## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$?(x - (-1.5))(x - (-1))(x - 2)$$

## Example




Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$2(x - (-1.5))(x - (-1))(x - 2)$$

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$2(x - (-1.5))(x - (-1))(x - 2) = (2x + 3)(x + 1)(x - 2)$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$2(x - (-1.5))(x - (-1))(x - 2) = (2x + 3)(x + 1)(x - 2)$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$


Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.



## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\ &= 2x^3 + x^2 - 7x - 6 \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$2x^3 + x^2 - 7x - 6 = 0$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\ &= 2x^3 + x^2 - 7x - 6 \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$\begin{aligned} 2x^3 + x^2 - 7x - 6 &= 0 \\ (2x + 3)(x + 1)(x - 2) &= 0 \end{aligned}$$

Make sure to practice with the graphing calculator you will on your exam(s). The graph appears to intersect the  $x$  axis at:  $-1.5, -1, 2$ . The left hand side should factor as:

$$\begin{aligned} 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\ &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\ &= 2x^3 + x^2 - 7x - 6 \end{aligned}$$

Check work to make sure we guessed the roots correctly.

## Example



Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$\begin{aligned}
 2x^3 + x^2 - 7x - 6 &= 0 \\
 (2x + 3)(x + 1)(x - 2) &= 0 \\
 x = -\frac{3}{2} \quad \text{or} \quad x = -1 \quad \text{or} \quad x = 2
 \end{aligned}$$

Make sure to practice with the graphing calculator you will on your exam(s). **The graph appears to intersect the x axis at:  $-1.5, -1, 2$ .** The left hand side should factor as:

$$\begin{aligned}
 2(x - (-1.5))(x - (-1))(x - 2) &= (2x + 3)(x + 1)(x - 2) \\
 &= (2x^2 + 5x + 3)(x - 2) = (2x^3 + 5x^2 + 3x) - (4x^2 + 10x + 6) \\
 &= 2x^3 + x^2 - 7x - 6
 \end{aligned}$$

**Check work to make sure we guessed the roots correctly.**



## Example


Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
?




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
? , ? , 3.




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?




$$x - 3 \overline{) x^3 - x^2 - 8x + 6}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?




$$x - 3 \overline{) x^3 - x^2 - 8x + 6}$$

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$x - 3 \quad \overline{x^3 - x^2 - 8x + 6}$$


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$x - 3 \overline{) x^3 - x^2 - 8x + 6}$$

Divide  $x^3$  by  $x$ .




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$x - 3 \overline{) \overset{x^2}{x^3 - x^2 - 8x + 6}}$$


Divide  $x^3$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{\phantom{x^3} ? \phantom{x^2} ?}
 \end{array}$$

Multiply  $x^2$  by divisor.


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6
 \end{array}$$


Multiply  $x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 \phantom{x^3} - 2x^2 - 8x + 6 \\
 \phantom{x^3} \phantom{- 2x^2} - 2x^2 - 8x + 6 \\
 \phantom{x^3} \phantom{- 2x^2} \phantom{- 2x^2} - 8x + 6 \\
 \phantom{x^3} \phantom{- 2x^2} \phantom{- 2x^2} \phantom{- 8x} 6 \\
 \phantom{x^3} \phantom{- 2x^2} \phantom{- 2x^2} \phantom{- 8x} \phantom{6} 6
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 \quad ? \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6
 \end{array}$$

Divide  $2x^2$  by  $x$ .


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \hline
 x^3 - x^2 - 8x + 6 \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \hline
 x^3 - x^2 - 8x + 6 \\
 \underline{x^3 - 3x^2} \\
 2x^2 - 8x + 6 \\
 \underline{\phantom{2x^2} \phantom{-8x} \phantom{+6} \phantom{0}} \\
 \phantom{2x^2} \phantom{-8x} \phantom{+6} \phantom{0}
 \end{array}$$

Multiply  $2x$  by divisor.




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \hline
 x^3 - x^2 - 8x + 6 \\
 \underline{x^3 - 3x^2} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \\
 6x + 6
 \end{array}$$


Multiply  $2x$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \overline{x^3 - x^2 - 8x + 6} \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6 \\
 2x^2 - 6x \\
 \hline
 \phantom{2x^2} 2x + 6 \\
 \phantom{2x^2} 2x + 4 \\
 \hline
 \phantom{2x^2} 2
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \\
 \overline{x^3 - x^2 - 8x + 6} \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6 \\
 2x^2 - 6x \\
 \hline
 -2x + 6
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x \quad ? \\
 \hline
 x^3 - x^2 - 8x + 6 \\
 x^3 - 3x^2 \\
 \hline
 2x^2 - 8x + 6 \\
 2x^2 - 6x \\
 \hline
 -2x + 6
 \end{array}$$

Divide  $-2x$  by  $x$ .


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6
 \end{array}$$


Divide  $-2x$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{\phantom{-2x} ? \phantom{+ 6} ?}
 \end{array}$$


Multiply  $-2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$


Multiply  $-2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 ?
 \end{array}$$

Subtract last two polynomials.




## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



$$\begin{array}{r}
 x - 3 \quad \overline{) \begin{array}{r} x^3 + 2x - 2 \\ x^3 - x^2 - 8x + 6 \\ \hline x^3 - 3x^2 \\ \hline 2x^2 - 8x + 6 \\ \hline 2x^2 - 6x \\ \hline -2x + 6 \\ \hline -2x + 6 \\ \hline 0 \end{array}} \\
 \hline
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



$$\begin{array}{r}
 x^2 + 2x - 2 \\
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) + 0 = 0$$

The graph appears to intersect the x axis at:   
 ? , ? , 3. What are the two roots besides 3?



**Quotient:**  $x^2 + 2x - 2$

$$\begin{array}{r}
 x - 3 \overline{) x^3 - x^2 - 8x + 6} \\
 \underline{x^3 - 3x^2} \phantom{+ 6} \\
 2x^2 - 8x + 6 \\
 \underline{2x^2 - 6x} \phantom{+ 6} \\
 -2x + 6 \\
 \underline{-2x + 6} \\
 0
 \end{array}$$


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) + 0 = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



<b>Quotient:</b>	$x^2 + 2x - 2$
$x - 3$	$\begin{array}{r} x^3 - x^2 - 8x + 6 \\ x^3 - 3x^2 \\ \hline 2x^2 - 8x + 6 \\ 2x^2 - 6x \\ \hline -2x + 6 \\ -2x + 6 \\ \hline 0 \end{array}$

**Remainder:**

0


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



<b>Quotient:</b>	$x^2 + 2x - 2$
$x - 3$	$\begin{array}{r} x^3 - x^2 - 8x + 6 \\ x^3 - 3x^2 \\ \hline 2x^2 - 8x + 6 \\ 2x^2 - 6x \\ \hline -2x + 6 \\ -2x + 6 \\ \hline 0 \end{array}$
<b>Remainder:</b>	$0$


## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x =$$



The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x =$$

$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?





## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.


$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.

$$x^3 - x^2 - 8x + 6 = 0$$


$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3$$


The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?



## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$


$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3 \quad x = \frac{-2 \pm \sqrt{12}}{2}$$

The graph appears to intersect the x axis at:  
 ? , ? , 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3 \quad x = \frac{-2 \pm \sqrt{12}}{2}$$


$$x = \frac{-2 \pm 2\sqrt{3}}{2}$$

The graph appears to intersect the x axis at:

?, ?, 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3 \quad x = \frac{-2 \pm \sqrt{12}}{2}$$


$$x = \frac{-2 \pm 2\sqrt{3}}{2} = -1 \pm \sqrt{3}.$$

The graph appears to intersect the x axis at:

**?**, **?**, 3. What are the two roots besides 3?

## Example

Plot the left hand side of the equation with a graphing calculator. Solve the equation.



$$x^3 - x^2 - 8x + 6 = 0$$

$$(x - 3)(x^2 + 2x - 2) = 0$$

$$x - 3 = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{(2)^2 - 4 \cdot 1 \cdot (-2)}}{2 \cdot 1}$$

$$x = 3$$

$$x = \frac{-2 \pm \sqrt{12}}{2}$$

$$x = \frac{-2 \pm 2\sqrt{3}}{2} = -1 \pm \sqrt{3}.$$

The graph appears to intersect the x axis at:

$-1 - \sqrt{3}$ ,  $-1 + \sqrt{3}$ , 3. What are the two roots besides 3?

Final answer:

$$x = 3 \quad \text{or} \quad x = -1 - \sqrt{3} \quad \text{or} \quad x = -1 + \sqrt{3}.$$

## Example


Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$






## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = ?$  .




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ .




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?  
Is there another root (far away from 0)?




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?  
**Is there another root (far away from 0)?**




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$

Divide  $2x^3$  by  $x$ .




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3}$$


Divide  $2x^3$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} ? \phantom{x} ?}
 \end{array}$$


Multiply  $2x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3}
 \end{array}$$


Multiply  $2x^2$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 \phantom{2x^3 - } 2x^2 + 5x - 3
 \end{array}$$

? ? ?


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \begin{array}{l} 2x^2 \quad ? \end{array} \\
 \hline
 2x^3 + x^2 + 5x - 3 \\
 \underline{2x^3 - x^2} \\
 2x^2 + 5x - 3
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 2x^2 + 2x \\
 \hline
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \quad \quad \quad \underline{2x^3 - x^2} \\
 \quad \quad \quad \quad \quad 2x^2 + 5x - 3
 \end{array}$$


Divide  $2x^2$  by  $x$ .

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \phantom{2x^2 + } \underline{? \quad ?}
 \end{array}$$

Multiply  $2x$  by divisor.




## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \\
 6x - 3
 \end{array}$$


Multiply  $2x$  by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 \phantom{2x^2} 6x - 3 \\
 \phantom{2x^2} \underline{6x - 3} \\
 \phantom{2x^2} 0
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} -} \phantom{2x^3 +} x^2 \phantom{+ 5x - 3} \\
 \phantom{x - \frac{1}{2}} \phantom{2x^3 +} 2x^2 + 5x - 3 \\
 \phantom{x - \frac{1}{2}} \phantom{2x^3 +} \underline{\phantom{2x^2 +} - x} \phantom{- 3} \\
 \phantom{x - \frac{1}{2}} \phantom{2x^3 +} \phantom{2x^2 +} 6x - 3
 \end{array}$$


Subtract last two polynomials.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$


We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3
 \end{array}$$

Divide  $6x$  by  $x$ .

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3
 \end{array}$$

Divide  $6x$  by  $x$ .

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} \phantom{6x} - 3} \\
 \phantom{x - \frac{1}{2}} \phantom{6x} 0
 \end{array}$$


Multiply 6 by divisor.

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$


We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:



$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 6x - 3} \\
 0
 \end{array}$$

Multiply 6 by divisor.

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$


We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{+ 5x - 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 ?
 \end{array}$$

Subtract last two polynomials.



## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 2x^2 + 2x + 6 \\
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 6x - 3} \\
 0
 \end{array}$$

Subtract last two polynomials.

## Example




Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)? Factor:

$$\begin{array}{r}
 x - \frac{1}{2} \quad \overline{2x^3 + x^2 + 5x - 3} \\
 \underline{\phantom{x - \frac{1}{2}} 2x^3 - x^2} \\
 2x^2 + 5x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 2x^2 - x} \\
 6x - 3 \\
 \underline{\phantom{x - \frac{1}{2}} 6x - 3} \\
 0
 \end{array}$$

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) + 0 = 0$$


We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?

Is there another root (far away from 0)? Factor:

**Quotient:**  $2x^2 + 2x + 6$

$$\begin{array}{r}
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) + 0 = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?

Is there another root (far away from 0)? Factor:


**Quotient:**  $2x^2 + 2x + 6$

$$\begin{array}{r}
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

**Remainder:**

0

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct?

Is there another root (far away from 0)? Factor:

**Quotient:**  $2x^2 + 2x + 6$

$$\begin{array}{r}
 x - \frac{1}{2} \overline{) 2x^3 + x^2 + 5x - 3} \\
 \underline{2x^3 - x^2} \phantom{- 3} \\
 2x^2 + 5x - 3 \\
 \underline{2x^2 - x} \phantom{- 3} \\
 6x - 3 \\
 \underline{6x - 3} \\
 0
 \end{array}$$

**Remainder:**

0


## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x =$$



We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example


Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x =$$

$$x = \frac{1}{2}$$



We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?





## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?



## Example

Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.


$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$


$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?



## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$


$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2} \quad x = \frac{-2 \pm \sqrt{-44}}{2 \cdot 2}$$

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example



Plot the left hand side of the equation with a graphing calculator. Find all real solutions of the equation.

$$2x^3 + x^2 + 5x - 3 = 0$$

$$(x - \frac{1}{2})(2x^2 + 2x + 6) = 0$$

$$x - \frac{1}{2} = 0 \quad \text{or} \quad x = \frac{-2 \pm \sqrt{2^2 - 4 \cdot 2 \cdot 6}}{2 \cdot 2}$$

$$x = \frac{1}{2} \quad x = \frac{-2 \pm \sqrt{-44}}{2 \cdot 2}$$

no real solution

We see only one root,  $x = 0.5 = \frac{1}{2}$ . Is our guess correct? Is there another root (far away from 0)?

## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

## Example

Solve the inequality.

$$\begin{array}{rcl} 2x^2 + 3x - 5 & \geq & 0 \\ (? \quad \quad)(? \quad \quad) & \geq & 0 \end{array}$$

## Example

Solve the inequality.


$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

## Example

Solve the inequality.

$$\begin{aligned}2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0\end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .





## Example

Solve the inequality.

$$\begin{aligned}2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0\end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .




## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$			


## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(?) ( ? )$		


## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(?)$		

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(?)$		

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(+)$		




## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(+)$	<b>?</b>	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$			
$(1, \infty)$	$(+)(+)$	$+$	


## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(-)(-)$		
$(1, \infty)$	$(+)(+)$	+	


## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(?)$		
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(?)$		
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(-)$		
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(-)$	<b>?</b>	
$(1, \infty)$	$(+)(+)$	$+$	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$			
$(-\frac{5}{2}, 1)$	$(+)(-)$	$-$	
$(1, \infty)$	$(+)(+)$	$+$	




## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(\textcolor{red}{?})(\textcolor{red}{?})$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	


## Example

Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(?)$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(?)$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(-)$		
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(-)$	<b>?</b>	
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	
$(1, \infty)$	$(+)(+)$	+	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	
$(-\infty, -\frac{5}{2})$	$(-)(-)$	$+$	
$(-\frac{5}{2}, 1)$	$(+)(-)$	$-$	
$(1, \infty)$	$(+)(+)$	$+$	

## Example

Solve the inequality.

$$\begin{aligned} 2x^2 + 3x - 5 &\geq 0 \\ (2x + 5)(x - 1) &\geq 0 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$

## Example


Solve the inequality.

$$2x^2 + 3x - 5 \geq 0$$

$$(2x + 5)(x - 1) \geq 0$$

$x \in ?$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$




## Example

Solve the inequality.

$$\begin{aligned}
 2x^2 + 3x - 5 &\geq 0 \\
 (2x + 5)(x - 1) &\geq 0 \\
 x &\in \left(-\infty, -\frac{5}{2}\right] \cup [1, \infty)
 \end{aligned}$$


Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
 The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .



Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$


# Example

Solve the inequality.



$$\begin{aligned}
 2x^2 + 3x - 5 &\geq 0 \\
 (2x + 5)(x - 1) &\geq 0 \\
 x &\in (-\infty, -\frac{5}{2}] \cup [1, \infty)
 \end{aligned}$$

Left hand side vanishes when  $x = -\frac{5}{2}$  and when  $x = 1$ .  
 The two roots split the real line into three intervals:  
 $(-\infty, -\frac{5}{2})$ ,  $(-\frac{5}{2}, 1)$ ,  $(1, \infty)$ .




Interval	Factor signs	Final sign	Sample pt	Value at sample pt
$(-\infty, -\frac{5}{2})$	$(-)(-)$	+	-100	$f(-100) > 0$
$(-\frac{5}{2}, 1)$	$(+)(-)$	-	0	$f(0) = -5 < 0$
$(1, \infty)$	$(+)(+)$	+	100	$f(100) > 0$

## Example

Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.  
$$2x^3 - 5x^2 + x + 2 > 0$$


## Example



Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - \quad) (x - \quad) (x - \quad) > 0$$

# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - ?) (x - ?)(x - ?) > 0$$

## Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - (-\frac{1}{2})) (x - 1)(x - 2) > 0$$

## Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$? (x - (-\frac{1}{2})) (x - 1)(x - 2) > 0$$



## Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

## Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.


$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ .



## Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.


$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ .



## Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.


$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ .



# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		

# Example



Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		
$(-\frac{1}{2}, 1)$		

# Example



Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		
$(-\frac{1}{2}, 1)$		
$(1, 2)$		
$(2, \infty)$		

# Example



Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$


Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$		
$(-\frac{1}{2}, 1)$		
$(1, 2)$		
$(2, \infty)$		



# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	?	?
$(-\frac{1}{2}, 1)$	?	?
$(1, 2)$	?	?
$(2, \infty)$	?	?

# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$


$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	$-$
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	$+$
$(1, 2)$	$(+)(+)(-)$	$-$
$(2, \infty)$	$(+)(+)(+)$	$+$

# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2\left(x - \left(-\frac{1}{2}\right)\right)(x - 1)(x - 2) > 0$$


$$x \in ?$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	-
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	+
$(1, 2)$	$(+)(+)(-)$	-
$(2, \infty)$	$(+)(+)(+)$	+

# Example



Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$


$$x \in (-\frac{1}{2}, 1) \cup (2, \infty)$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	-
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	+
$(1, 2)$	$(+)(+)(-)$	-
$(2, \infty)$	$(+)(+)(+)$	+

# Example




Plot the function  $2x^3 - 5x^2 + x + 2$ . Solve the inequality.

$$2x^3 - 5x^2 + x + 2 > 0$$

$$2(x - (-\frac{1}{2}))(x - 1)(x - 2) > 0$$

$$x \in (-\frac{1}{2}, 1) \cup (2, \infty)$$

Left hand side vanishes when  $x = -\frac{1}{2}$ , when  $x = 1$  and when  $x = 2$ . The two roots split the real line into four intervals:  $(-\infty, -\frac{1}{2})$ ,  $(-\frac{1}{2}, 1)$ ,  $(1, 2)$ ,  $(2, \infty)$ .



Interval	Factor signs	Final sign from plot
$(-\infty, -\frac{1}{2})$	$(-)(-)(-)$	-
$(-\frac{1}{2}, 1)$	$(+)(-)(-)$	+
$(1, 2)$	$(+)(+)(-)$	-
$(2, \infty)$	$(+)(+)(+)$	+