

Directional Partial-Wave Sweeps and the Krylov Solve

The propagator matvec for elastic multiple scattering on a depth-stratified voxel grid

Tod Nestor

16 June 2026

Abstract

The multiple scattering of elastic waves by voxels superimposed on a depth-stratified background is a Foldy–Lax / volume integral equation, $T = T_0(I - G_0T_0)^{-1}$, solved by a Krylov method whose only nontrivial operation is the application of the background propagator G_0 . This note describes that application as an alternating-direction partial-wave summation: the lateral coupling is summed by constant- z left-right (k_x) and constant- (z, x) in-out (k_y) cumulative-phase sweeps, and the coupling between depth planes by the up-down Kennett reflectivity recursion (k_z). Choosing the lateral directions for the in-plane coupling sums the propagator along axes in which the voxels are separated, so the k_z partial-wave sum that diverges at zero depth separation is never evaluated. The sweeps are a pure forward matvec; all multiple scattering, to every order, is resolved by the Krylov iterations.

Companion series. *One of a companion series on elastic multiple scattering from cubic heterogeneities on a depth-stratified background, the T -matrix programme $T = T_0(I - G_0T_0)^{-1}$: (I) the stratified inter-site propagator, the same-depth convergence theory, the SH and P/SV traction kernels, interplane side scattering and the first-shell near stencil (`EwaldIntraPlanePropagator`); (II) the contrast-source volume integral equation and the CG-FFT / seismic grid T -matrix lineage (`ContrastSourceVIE`); (III) the directional partial-wave sweep matvec and the Krylov multiple-scattering solve (`DirectionalSweepSolver`); with the empirical companion `RQ1_Planar-Decomposition-Proposal`. This is note (III).*

Contents

1	The system, and where the cost lives	2
2	The matvec is a partial-wave summation	2
2.1	Lateral sweeps: constant- z left-right and constant- (z, x) in-out	2
2.2	Vertical sweep: up-down Kennett between depth planes	3
2.3	Why these directions	3
2.4	The same-depth term in a stratified background	3
3	Alternating-direction composition	4
4	The Krylov solve	4
5	Validation status	5
6	What is distinctive	6

1 The system, and where the cost lives

Voxels (cubic cells) are superimposed on a background that carries the lateral average of the medium at each depth: laterally homogeneous within a constant- z plane, layered in z . Each voxel has an on-site T-matrix T_0 , the single-cell scatterer (Eshelby with the $r = 0$ delta-function correction up to the 27-component Galerkin form, the self-term already closed), and the voxels are coupled by the background propagator G_0 . The multiple scattering is the Foldy–Lax system [1, 2, 3]

$$(I - G_0 T_0) \psi = \psi^{\text{inc}}, \quad T = T_0 (I - G_0 T_0)^{-1}, \quad (1)$$

where ψ collects the exciting (incoming) fields at the voxels. The operator T_0 is block-diagonal and local. The whole cost is the off-diagonal G_0 : the inter-voxel communication. The strategy is to solve (1) by a Krylov method whose matrix-vector product is one application of G_0 , and to make that application fast.

The essential point is the division of labour. The application of G_0 is a *forward* operation, a summation of partial waves that carries the radiation of every voxel to every other. It contains no inversion and no embedding. All of the multiple scattering, the reverberations to every order, is built and resummed by the Krylov iterations, not by the propagator.

2 The matvec is a partial-wave summation

Given source amplitudes $b = T_0 \psi$ at the voxels, the matvec returns the field they radiate, $\psi' = G_0 b$. For the homogeneous part of the background the propagator is a sum of partial waves, and a partial wave’s phase accumulates linearly in space: e^{ikx} at the next voxel is the current one times e^{ikp} , with p the pitch. The sum is therefore built by carrying that running phase along a line, one term added per voxel. This is the sweep. The only freedom is which axis to sum along, and the choice is dictated by where the voxels are separated.




Figure 1: The G_0 matvec as three directional partial-wave sweeps over the voxel grid (cross-section; the third axis y runs into the page). Within each depth plane the lateral coupling is summed by constant- z left-right (k_x) and constant- (z, x) in-out (k_y) cumulative-phase sweeps; the coupling between depth planes is the up-down Kennett reflectivity recursion (k_z). The lateral directions carry the in-plane coupling along axes of nonzero separation, so the k_z sum is never evaluated at $\Delta z = 0$.

2.1 Lateral sweeps: constant- z left-right and constant- (z, x) in-out

Within a depth plane the background is homogeneous, so the lateral propagation is pure phase, with no background reflections to sum. Split the in-plane propagator on the k_x pole into a right-going and a left-going partial wave and sum each by a sweep in its own direction. With a^+ the right-going accumulated field, s_n the source radiated at voxel n , and $\Phi_x = e^{ik_x p}$ the

one-pitch lateral propagator,

$$a_{n+1}^+ = (a_n^+ + s_n) \Phi_x \quad (\text{left to right}), \quad a_{n-1}^- = (a_n^- + s_n) \Phi_x \quad (\text{right to left}). \quad (2)$$

Each partial wave is carried along the direction in which it decays: the right-going evanescent branch $e^{-\kappa_x x}$ is swept to the right, the left-going branch to the left, so the growing exponential is never formed and no reflection/transmission bookkeeping is needed for stability. The in-out sweep in y is identical, run along each constant- (z, x) line on the k_y pole. The two together sum the full in-plane partial-wave series as nested one-dimensional phase accumulations.

2.2 Vertical sweep: up-down Kennett between depth planes

In z the background is not homogeneous: the lateral average changes from plane to plane, so the depth-average layering has its own reflections and transmissions. The coupling between depth planes is therefore the up-down Kennett reflectivity recursion [4] on the k_z pole, propagating the source amplitudes through the layered background and summing its internal up/down multiples. This is the one sweep that carries reflection and transmission coefficients, and it does so for the *background* layering, not for the heterogeneity; it is the standard invariant-embedding recursion, stable by construction, and its coefficients depend only on the background and are formed once. The heterogeneity scattering remains entirely in T_0 and the Krylov loop.

2.3 Why these directions

The conventional construction sums the in-plane coupling on the k_z pole, the up/down decomposition, whose surviving (k_x, k_y) integral loses its convergence factor $e^{-\kappa_\perp |\Delta z|}$ at $\Delta z = 0$ and diverges at equal depth. The lateral split avoids it outright: the in-plane coupling is summed on the k_x and k_y poles, along which distinct in-plane voxels are separated, so the divergent same-depth k_z sum is never the one evaluated. The k_z sweep is used only between planes, where $\Delta z \neq 0$ and the Kennett recursion is exact and stable. Every sweep marches along an axis of nonzero separation.

2.4 The same-depth term in a stratified background

The lateral sweep of §2.1 sums the same-depth coupling of the *whole space*. That is not the whole of it. When the background is stratified, two voxels at the same depth are also coupled by paths that leave one voxel, reflect from a layer boundary above or below, and return to the plane. Those paths are absent from the lateral sweep by construction, because the k_x residue carries only the direct term.

They cannot simply be added by evaluating the stratified propagator at $\Delta z = 0$ and integrating: that is the divergence of §2.3, and it is present in the stratified kernel exactly as in the whole-space one, its strain-strain block growing like $|k_x|$. The resolution is to subtract it. Writing G_0^{lay} for the stratified propagator between a plane and itself and G_0^∞ for the whole-space kernel at equal depth,

$$G_0^{\text{lay}}(\Delta z = 0) = \underbrace{G_0^\infty(\Delta z = 0)}_{\text{lateral sweep, by } k_x \text{ residue}} + \underbrace{\Delta G_0}_{\text{reverberation only}}, \quad (3)$$

and the two terms on the right are individually non-integrable while their difference is not. Every path contributing to ΔG_0 travels at least twice the distance H to the nearest interface, so it carries $e^{ik_z 2H}$ and ΔG_0 decays exponentially in k_x . Its k_x integral therefore converges, and the divergence cancels between the two terms rather than being regularised in either.

Two consequences are worth stating plainly. First, ΔG_0 vanishes identically when the background is uniform, so the construction reduces to the whole-space sweep with nothing left

over. Second, ΔG_0 is retained at $\Delta x = 0$: a voxel *does* interact with itself through the layering, and since T_0 is the single-site T -matrix of the whole space, its self-term does not contain that path. Dropping the $\Delta x = 0$ entry — the natural thing to do, since the whole-space self-term is excluded everywhere else — would silently discard every layer-return-to-self.

3 Alternating-direction composition

The three sweep pairs compose by alternating direction, in the spirit of the alternating-direction methods of Peaceman and Rachford [5]: the two-dimensional in-plane sum factors into an x pass and a y pass, and the full three-dimensional propagator into those plus the z recursion. Each pass is a one-dimensional, $O(N)$ accumulation, and the matvec is their ordered application:

1. **up-down** (k_z , **Kennett**): propagate the source amplitudes between depth planes through the layered background;
2. **left-right** (k_x): along each constant- z row, sweep left to right for the right-going field and right to left for the left-going field;
3. **in-out** (k_y): along each constant- (z, x) line, sweep in and out the same way.

The result is $\psi' = G_0 b$, the field every voxel radiates onto every other, summed along directions of nonzero separation throughout. Six direction-pure sweeps in all, none of which ever forms the $\Delta z = 0$ sum.

4 The Krylov solve

The matvec of Section 3 is a pure forward application of G_0 . The multiple scattering is solved around it by a Krylov method (GMRES [6]) applied to (1). One iteration is

$$v \mapsto v - G_0 T_0 v, \quad (4)$$

that is, one block-diagonal T_0 (local, the cube T -matrix) followed by one G_0 (the three sweeps). The Krylov subspace builds the scattering series and resums it, reaching the solution in far fewer applications than the bare Neumann (Born) series and remaining stable at strong contrast where that series would diverge. The reverberations and back-and-forth that the forward sweep deliberately does *not* contain are produced here, across the iterations.

For strong contrast the iteration is preconditioned, the role played by the renormalised scattering series of Jakobsen and Wu [7] and the convergent Born series of Osnabrugge, Leedum-rongwatthanakun and Vellekoop [8]: a preconditioner on the Krylov solve, transparent to the sweep, which never needs to know the contrast level.

The stack, end to end.

1. **Setup.** Form T_0 per voxel (self-term closed) and the background Kennett reflectivity coefficients (once).
2. **Matvec** $G_0 b$. Up-down Kennett between planes, then left-right (k_x) and in-out (k_y) cumulative-phase sweeps within planes.
3. **Solve.** GMRES on $(I - G_0 T_0)\psi = \psi^{\text{inc}}$, preconditioned at strong contrast, each step one T_0 and one matvec.
4. **Output.** The exciting fields ψ , hence the global T and any observable (reflection matrix, shot gather) by one more radiation of $b = T_0 \psi$.

5 Validation status

The 2½-D case (heterogeneity in (z, x) , y invariant, k_y a parameter) is implemented and gated. Each claim below is the same physics through two independent code paths, so the residual measures defect, not discretisation, except where noted.

Claim	Arbiter	
amplitude $\times$ phase ^{<i>n</i>} reproduces the bundled kernel	post- k_x -residue kernel	
and the residual is <i>flat</i> in separation $n = 1 \dots 32$	—	
left-going amplitude is the x -reflection of the right	parity identity	
lateral sweep = pairwise double sum, distinct sources	$O(N^2)$ sum	
and the residual is <i>flat</i> in lattice size	—	
k_y -integrated kernel = closed-form propagator	Kupradze closed form	falls
inter-plane sweep = exact real-space pairwise sum	Kupradze closed form	falls
lateral coupling does not wrap (no periodicity)	kernel at true separation	
stratified operator reduces to whole space	independent k_z -residue kernel	
same-depth reverberation vanishes for a uniform background	—	
self-energy $\equiv$ dressed T-matrix, $T_d = T_0(I - DT_0)^{-1}$	direct vs dressed solve	
reverberation obeys symplectic reciprocity	J_6 law, $\pm k$	
self-energy recedes with the reflector	—	monotone, 10
partition: every pair once, self-term per background	support count	
GMRES = dense LU and Neumann series, $\rho(G_0 T_0) = 0.92$	two direct methods	

Three points of method are worth recording because each cost time to learn. Symmetry and reciprocity invariants are homogeneous of degree one and therefore blind to an overall scale error; every such gate here is paired with one absolute comparison. Where a tolerance had to be loosened it was loosened only after measuring why, and the accompanying test asserts the property that actually discriminates — that the residual is flat in separation, or flat in lattice size — since the dangerous failure is a factor that accumulates along the sweep. Finally, a residual that does *not* fall under refinement is a convention error, not a quadrature error: two of the discrepancies met here were exactly that, one an index ordering and one an attenuation mismatch between an operator and the Green’s function it multiplied.

On the self-energy, and which invariant applies to it. The same-depth reverberation ΔG_0 does *not* satisfy the whole-space symmetry $WM = (WM)^T$: measured 0.33 to 1.06 across k_x , at every separation. That is not a defect but a misapplied invariant. At coincident planes the *full* layered kernel fails the symplectic reciprocity law at 9.5×10^{-3} while ΔG_0 satisfies it at 8×10^{-14} : the non-reciprocal part is the one-sided jump carried by the direct term, and removing that term is precisely what (3) does. The reverberation is the clean reciprocal object; the whole-space symmetry is the wrong thing to demand of it. This is recorded because the same pattern — a plausible invariant that the exact reference also fails — has already cost this programme one false gate.

Not yet done. Two things. The in-out k_y pair, and with it the full three-dimensional Cartesian case, is the remaining piece; until it lands, the solver is 2½-D and a comparison against a three-dimensional solver measures the geometry rather than the sweeps. And the *absolute magnitude* of ΔG_0 is not yet checked against an independent physical reference: what is established is that it is reciprocal, that it vanishes without layering, that it recedes with the reflector, and that it behaves as a self-energy under the dressed-T identity. The route to closing it is the single-bounce limit — as the contrast weakens, ΔG_0 must approach the interface reflection coefficient times the whole-space propagator over twice the standoff — which is checkable in isolation.

6 What is distinctive

Two choices set this apart from a standard FFT-accelerated volume integral equation. First, the forward operator is a directional partial-wave summation rather than a circular-convolution FFT, so the in-plane coupling is summed along the separated axes and the $\Delta z = 0$ divergence is structurally absent rather than regularised — and where the stratified background does contribute at equal depth, §2.4, it enters as a difference in which the divergence cancels rather than as a term requiring regularisation. Second, the depth structure is carried exactly by the stratified propagator on the true layered background, not approximated on a homogeneous whole-space kernel. The multiple scattering is left where it belongs, in a Krylov solve that calls the sweep as its matvec, so the propagator stays a clean, stable, forward summation and the solver carries the contrast.

References

- [1] L. L. Foldy, *The multiple scattering of waves*, Phys. Rev. **67**, 107 (1945).
- [2] M. Lax, *Multiple scattering of waves*, Rev. Mod. Phys. **23**, 287 (1951).
- [3] J. E. Gubernatis, E. Domany and J. A. Krumhansl, *Formal aspects of the theory of the scattering of ultrasound by flaws in elastic materials*, J. Appl. Phys. **48**(7), 2804 (1977).
- [4] B. L. N. Kennett, *Seismic Wave Propagation in Stratified Media*, Cambridge University Press (1983).
- [5] D. W. Peaceman and H. H. Rachford, *The numerical solution of parabolic and elliptic differential equations*, J. Soc. Indust. Appl. Math. **3**, 28 (1955).
- [6] Y. Saad and M. H. Schultz, *GMRES: a generalized minimal residual algorithm for solving nonsymmetric linear systems*, SIAM J. Sci. Stat. Comput. **7**, 856 (1986).
- [7] M. Jakobsen and R.-S. Wu, *Renormalized scattering series for frequency-domain waveform modelling of strong velocity contrasts*, Geophys. J. Int. **206**(2), 880 (2016).
- [8] G. Osnabrugge, S. Leedumrongwatthanakun and I. M. Vellekoop, *A convergent Born series for solving the inhomogeneous Helmholtz equation in arbitrarily large media*, J. Comput. Phys. **322**, 113 (2016).