

Coupling Integrals of Polynomial Cubes at a Welded Elastic Interface

The Static Image in Closed Form, and the Plane Waves for the Rest

T. M. Nestor

Independent researcher, Australia

Draft, 11 October 2026

Abstract

In a stratified elastic reference medium the Green's tensor is that of a homogeneous medium plus the waves reflected and transmitted at the interfaces, and it is most naturally available not as a tabulated function but through its plane-wave, up- and down-going, factorisation. A volume-integral method that divides the heterogeneity into cubic cells carrying polynomial fields needs the Galerkin coupling of every pair of cells through this reference. Between cells in planes that do not touch, the plane-wave representation with the cells' moments in closed form is exact, and its integral over the lateral wavenumber converges exponentially. Next to a welded interface it does not: a cell is coupled to itself and to its neighbours through its own mirror image, which touches them, and the reflected coupling converges only algebraically. We show that the singular part is the static field of a point force at the interface, to which the reflected and transmitted kernels tend as $(\omega/\beta q)^2$ at large lateral wavenumber q , and obtain it and its coupling blocks in closed form. Solved in the lateral-wavenumber domain, Rongved's problem gives every entry as $e^{-q\zeta}/q$ times a polynomial; in space the reflected and transmitted static fields are 356 and 275 terms built on Kelvin's potential and two logarithmic potentials, Mindlin's terms, which at a welded interface are present for every pair of distinct media and which we write as a line of images. The moments of these kernels over a box with the singular point at a vertex reduce, by Euler's identity for homogeneous functions, to faces, edges and edges to infinity, with Hadamard finite parts where the degree is negative and two base integrals where it is zero, $B_1 = \frac{5}{4}\text{Cl}_2(\pi/3) - G$ and $B_2 = \frac{1}{3}\log[(1 + \sqrt{2})^2/(2(1 + \sqrt{3}))]$, so that every moment is elementary apart from Catalan's constant and one value of Clausen's function. The corner moments and the blocks agree with independent routes to 10^{-15} and 10^{-13} or better, coincide with the blocks of the homogeneous medium to 2 to 8×10^{-15} when the two media are the same, and with the static part subtracted the plane-wave remainder converges as Q^{-5} in the cutoff Q instead of Q^{-3} .

1 Introduction

A volume-integral method for elastic waves represents a heterogeneous body as a contrast with a reference medium, divides the contrast into cells, and couples every pair of cells through the Green's tensor of the reference. When the reference is homogeneous the coupling of cubic cells

carrying polynomial fields can be obtained in closed form and evaluated to round-off [1]. In seismology the natural reference is stratified, and its Green's tensor is that of a homogeneous medium plus waves reflected and transmitted at the interfaces. The stratified Green's tensor is rarely tabulated: it is computed through its plane-wave representation, in which, at each lateral wavenumber, waves travel up and down through the layers and are reflected and transmitted at each interface [2, 3]. A volume method built on a stratified reference then needs the coupling blocks in a form that uses only this up- and down-going factorisation [4]; tabulating the Green's tensor between all pairs of depths would cost storage quadratic in the number of depth planes, where the plane waves need only linear storage.

For cells in different planes that do not touch, the plane-wave representation with the cells' moments in closed form is exact, and its integral over the lateral wavenumber converges exponentially because the evanescent waves decay between the planes. For cells in touching planes it diverges, and those blocks are the closed forms of the homogeneous medium [1]. At an interface the same failure reaches further. The reflection of a cell's field is the field of its mirror image, and the image of a cell that touches the interface touches the cell itself and its neighbours along the interface. The reflected coupling of every such cell is therefore singular, and its plane-wave integral converges only algebraically.

This paper removes that obstacle. Its contributions are the following.

1. The plane-wave representation of the coupling blocks, stated for the direct, reflected and transmitted couplings with the cells' moments in closed form (Section 2).
2. The identification of the singular part of the reflected and transmitted couplings as the static field of a point force at a welded interface (Rongved's problem), with the dynamic kernels tending to it as $(\omega/\beta q)^2$; subtracting it changes the convergence of the plane-wave remainder from Q^{-3} to Q^{-5} (Sections 2 and 3).
3. The static image and the transmitted static field in space, as 356 and 275 canonical terms on three potentials, with Mindlin's logarithmic terms written as a line of images (Section 3).
4. The coupling blocks of these fields in closed form: the moments over a box with the singular point at a vertex reduce to elementary functions and two constants, B_1 and B_2 , which we evaluate in closed form (Sections 4 and 5).
5. Verification of each step against routes that share no step with it (Section 6).

The dynamic remainder, smooth by construction, is left to the plane waves; Section 7 describes how they carry it through a stack of layers.

2 Cells, blocks and the plane-wave representation

Cells and blocks. A cell is a cube of half-width h centred at $\mathbf{x}_c$, with local coordinates $\xi = (\mathbf{x} - \mathbf{x}_c)/h \in [-1, 1]^3$. It carries a nine-component state, the displacement and an independent engineering strain, as a polynomial in ξ [5]; the field is tested against products of Legendre

polynomials $L_a(\xi) = \prod_i P_{l_{a,i}}(\xi_i)$, and the source density, the contrast times the state, is written in the monomials $m_c(\xi') = \prod_i \xi_i'^{e_{c,i}}$. With $\mathbf{P}(\mathbf{x}, \mathbf{x}')$ the 9×9 point propagator of the reference medium, which maps a point source of force and engineering stress at $\mathbf{x}'$ to the displacement and engineering strain at $\mathbf{x}$, a field cell V_m and a source cell V_n couple through the blocks

$$\mathbf{K}_{ac} = \int_{V_m} \int_{V_n} L_a(\xi) \mathbf{P}(\mathbf{x}, \mathbf{x}') m_c(\xi') d^3x d^3x'. \quad (1)$$

In a homogeneous medium $\mathbf{P}$ depends on $\mathbf{x} - \mathbf{x}'$ only, and with $\mathbf{x} = \mathbf{x}_m + h\xi$, $\mathbf{x}' = \mathbf{x}_n + h\xi'$ the block becomes an integral over the separation $\mathbf{s} = h(\xi - \xi') \in [-2h, 2h]^3$ against the weight $W_{ac}(\mathbf{s}) = \prod_i w_{l_{a,i}e_{c,i}}(s_i)$, with $w_{\alpha\beta}(s) = h \int v^\alpha (v - s/h)^\beta dv$ the cross-correlation of the two cells' one-dimensional factors, a piecewise polynomial on eight boxes [1]. The kernel is singular at $\mathbf{s} = -(\mathbf{x}_m - \mathbf{x}_n)$, which for touching cells is a vertex of some of the boxes.

The plane-wave representation. Take coordinates in the order (z, x, y) , z the depth, and a lateral wavenumber $\mathbf{k} = (k_x, k_y) \in \mathbb{R}^2$, $q = |\mathbf{k}|$. For each wave $v \in \{P, SV, SH\}$ of speed c_v ($c_P = \alpha$, $c_{SV} = c_{SH} = \beta$) let

$$k_{z,v} = (\omega^2/c_v^2 - q^2)^{1/2}, \quad \text{Im } k_{z,v} \geq 0, \quad \mathbf{k}_v^\sigma = (\sigma k_{z,v}, k_x, k_y), \quad \sigma = \pm 1,$$

the wave vector travelling down ($\sigma = +1$) or up ($\sigma = -1$). Its polarisations are

$$\mathbf{e}_P^\sigma = \frac{\alpha}{\omega} \mathbf{k}_P^\sigma, \quad \mathbf{e}_{SH} = \frac{(0, -k_y, k_x)}{q}, \quad \mathbf{e}_{SV}^\sigma = \frac{\beta}{\omega} \mathbf{k}_{SV}^\sigma \times \mathbf{e}_{SH},$$

with $\mathbf{e}_{SH} = (0, 1, 0)$ at $\mathbf{k} = \mathbf{0}$. Each is a unit vector in the bilinear sense, $\mathbf{e} \cdot \mathbf{e} = 1$, which, unlike $\mathbf{e} \cdot \bar{\mathbf{e}} = 1$, continues analytically into the evanescent range $q > \omega/c_v$. The state of the wave is the nine-vector $\mathbf{d}_v^\sigma = (\mathbf{e}, \boldsymbol{\varepsilon})$ of its displacement and engineering strain, $\varepsilon_{ij} = \frac{i}{2}(k_i e_j + k_j e_i)$ with $\mathbf{k} = \mathbf{k}_v^\sigma$ and the three shears doubled, and $\mathbf{M} = \text{diag}(1, 1, 1, 1, 1, 1, \frac{1}{2}, \frac{1}{2}, \frac{1}{2})$ pairs the engineering-stress entries of a source with the engineering-strain entries of a state. In a homogeneous medium of density ρ , for $\sigma(z - z') > 0$,

$$\mathbf{P}(\mathbf{x}, \mathbf{x}') = \frac{1}{4\pi^2} \int_{\mathbb{R}^2} \sum_v w_v \mathbf{d}_v^\sigma \mathbf{d}_v^{\sigma T} \mathbf{M} e^{i\mathbf{k}_v^\sigma \cdot (\mathbf{x} - \mathbf{x}')} d^2k, \quad w_v = \frac{i}{2\rho c_v^2 k_{z,v}}, \quad (2)$$

the source entering through $\mathbf{d}_v^{\sigma T} \mathbf{M}$ and the field through $\mathbf{d}_v^\sigma$, as reciprocity requires. For cells in planes that do not touch, $|z_m - z_n| \geq 4h$, set $\sigma = \text{sgn}(z_m - z_n)$; Eq. (1) then becomes

$$\mathbf{K}_{ac} = \frac{1}{4\pi^2} \int_{\mathbb{R}^2} \sum_v w_v \Lambda_a(\mathbf{k}_v^\sigma) \mu_c(-\mathbf{k}_v^\sigma) \mathbf{d}_v^\sigma \mathbf{d}_v^{\sigma T} \mathbf{M} e^{i\mathbf{k}_v^\sigma \cdot (\mathbf{x}_m - \mathbf{x}_n)} d^2k, \quad (3)$$




Figure 1. Cells at a welded interface, in a vertical section; z is depth. (a) Two cells in medium A that touch the interface and each other. The reflected coupling is the field of the source cell's mirror image (dashed), at the depth $\zeta = -(z + z')$ below the field point, and the image touches the field cell at the singular point. (b) A source cell in A and a field cell in B that touch at a point of the interface. The transmitted coupling depends on $\zeta = z - z'$, and the singular point is where the cells touch.

with the scalar moments of the two cells, both in closed form for complex arguments [1],

$$\Lambda_a(\kappa) = h^3 \int_{[-1,1]^3} L_a(\xi) e^{ih\kappa \cdot \xi} d^3\xi = h^3 \prod_i 2i^{l_{a,i}} j_{l_{a,i}}(h\kappa_i), \quad (4)$$

$$\mu_c(\kappa) = h^3 \int_{[-1,1]^3} m_c(\xi') e^{ih\kappa \cdot \xi'} d^3\xi' = h^3 \prod_i F_{e,i}(h\kappa_i), \quad F_n(t) = \int_{-1}^1 \xi^n e^{it\xi} d\xi, \quad (5)$$

j_l the spherical Bessel functions and F_n a finite combination of them.

Reflection and transmission. Let medium A, with Lamé constants λ_A , μ_A and density ρ_A , fill $z < 0$ and medium B fill $z > 0$, welded at $z = 0$ (Figure 1). With the amplitudes referred to the interface and the polarisations of each medium, a down-going wave ν in A gives the up-going wave μ in A with amplitude $R_{\mu\nu}(\mathbf{k})$ and the down-going wave μ in B with amplitude $T_{\mu\nu}(\mathbf{k})$, obtained at each $\mathbf{k}$ from the continuity of displacement and traction, a 6×6 linear system [3]. For $\mathbf{x}$ and $\mathbf{x}'$ both in A the reflected propagator is

$$\mathbf{P}^R(\mathbf{x}, \mathbf{x}') = \frac{1}{4\pi^2} \int_{\mathbb{R}^2} \sum_{\mu, \nu} \mathbf{d}_\mu^{A-} R_{\mu\nu} w_\nu^A \mathbf{d}_\nu^{A+T} \mathbf{M} e^{i(\mathbf{k}_\mu^{A-} \cdot \mathbf{x} - \mathbf{k}_\nu^{A+} \cdot \mathbf{x}')} d^2k, \quad (6)$$

and for $\mathbf{x}'$ in A and $\mathbf{x}$ in B the whole coupling is the transmitted one,

$$\mathbf{P}^T(\mathbf{x}, \mathbf{x}') = \frac{1}{4\pi^2} \int_{\mathbb{R}^2} \sum_{\mu, \nu} \mathbf{d}_\mu^{B+} T_{\mu\nu} w_\nu^A \mathbf{d}_\nu^{A+T} \mathbf{M} e^{i(\mathbf{k}_\mu^{B+} \cdot \mathbf{x} - \mathbf{k}_\nu^{A+} \cdot \mathbf{x}')} d^2k, \quad (7)$$

the superscripts naming the medium whose wave numbers and polarisations are used. The blocks follow as in Eq. (3), with $\Lambda_a(\mathbf{k}_\mu^\pm) \mu_c(-\mathbf{k}_\nu^\pm)$ in place of $\Lambda_a \mu_c$ and the phase $e^{i(\mathbf{k}_\mu \cdot \mathbf{x}_m - \mathbf{k}_\nu \cdot \mathbf{x}_n)}$. The vertical phase of the reflected term is $e^{i(k_{z,\mu}|z| + k_{z,\nu}|z'|)}$: its decay on the evanescent branch is set by the distance of the receiver from the source's mirror image $\mathbf{x}^* = (-z', x', y')$.

Where the plane waves fail. For cells in touching planes, $|z_m - z_n| = 2h$, the integral (3) does not converge: on the evanescent branch $k_{z,\nu} = i\kappa$, $\kappa \rightarrow \infty$, the moments Λ_a and μ_c grow as

$e^{\kappa h}$ each and cancel the decay $e^{-2\kappa h}$ of the phase. In the reflected coupling the relevant distance is that from the image, $|z_m + z_n|$, which is $2h$ for any two cells that touch the interface, and the same cancellation occurs (Figure 1a). The integrand then decays only algebraically in q , and so does the error of a truncated integral. Table 1 shows it for a cell that touches the interface, coupled to its own image: the change of the integral between successive fourfold cutoffs Q falls by a factor of 68, close to the $4^3 = 64$ of convergence as Q^{-3} .

The singular part of the integrand is its behaviour at large q . There $k_{z,v} = iq[1 - \omega^2/(2c_v^2 q^2) + O(q^{-4})]$, and the reflected and transmitted kernels depend on the frequency only through $(\omega/c_v q)^2$. Their limit is the static kernel of Section 3, and the difference falls as $(\omega/\beta q)^2$. Writing $\mathbf{K}^R = \mathbf{K}_0^R + (\mathbf{K}^R - \mathbf{K}_0^R)$, with $\mathbf{K}_0^R$ the block of the static image, the first term is obtained in closed form (Section 4) and the second by the plane-wave integral with the static spectrum subtracted under the integral sign. The remainder converges as Q^{-5} (Table 1), and the same holds for the transmitted coupling.

	$Qh = 10 \rightarrow 40$	$Qh = 40 \rightarrow 160$
reflected kernel	7.5×10^{-3}	1.1×10^{-4}
reflected kernel less its static part	4.8×10^{-5}	5.1×10^{-8}

Table 1. The plane-wave integral of the reflected block of a cell touching the interface, coupled to its own image: the change between successive fourfold cutoffs of the lateral wavenumber, relative to the block. The ratios of successive changes, 68 and 940, are those of convergence as Q^{-3} and Q^{-5} .

3 The static field of a point force at a welded interface

In the lateral-wavenumber domain. Let a unit point force act in A at depth $z' < 0$. In the frame whose x axis lies along $\mathbf{k}$, the static Navier equations of an isotropic medium have the solutions $(a + bz)e^{\pm qz}$, the double root replacing the separate P and S waves of the dynamic problem. The field in A is Kelvin's field plus a reflected field regular as $z \rightarrow -\infty$; the field in B is transmitted and regular as $z \rightarrow \infty$; displacement and traction on the planes $z = \text{const}$ are continuous at $z = 0$. This is the problem of Rongved [6], solved here in the lateral-wavenumber domain. Its solution has a simple structure: every entry of the reflected tensor is $e^{q(z+z')}/q$ times a polynomial of degree at most one in each of qz and qz' , and every entry of the transmitted tensor is $e^{-q(z-z')}/q$ times a polynomial in qz and qz' , the coefficients rational in the Lamé constants. When the two media coincide the reflected tensor vanishes and the transmitted one is Kelvin's.

In space. With $\zeta = -(z + z') > 0$ for the reflected field and $\zeta = z - z' > 0$ for the transmitted one, ρ the lateral separation and $R = (|\rho|^2 + \zeta^2)^{1/2}$, the transform to space follows from

$$\frac{1}{4\pi^2} \int_{\mathbb{R}^2} e^{i\mathbf{k}\cdot\boldsymbol{\rho}} q^m e^{-q\zeta} d^2k = \begin{cases} (-\partial_\zeta)^{m+1} \Phi_0, & m \geq -1, \\ \Phi_1, & m = -2, \\ \Phi_2, & m = -3, \end{cases} \quad (8)$$

with the three potentials

$$\Phi_0 = \frac{1}{2\pi R}, \quad \Phi_1 = -\frac{\log(R + \zeta)}{2\pi}, \quad \Phi_2 = \frac{\zeta \log(R + \zeta) - R}{2\pi}, \quad (9)$$

each the ζ -antiderivative of the one before, $\partial_\zeta \Phi_1 = -\Phi_0$ and $\partial_\zeta \Phi_2 = -\Phi_1$. Rotating from the frame of $\mathbf{k}$ to fixed axes brings factors k_h/q ; since ik_h becomes ∂_{ρ_h} , an entry $(k_h/q)f(q)$ is ∂_{ρ_h} of the transform of $f/(iq)$, and an entry $(k_h k_l/q^2)f(q)$ is $-\partial_{\rho_h} \partial_{\rho_l}$ of the transform of f/q^2 . These divisions by q are what produce the powers $m = -2$ and -3 , and so the logarithmic potentials. They are the terms that Mindlin [7] found for the half-space with a free surface; at a welded interface they are present for every pair of media, with coefficients that vanish only when the media coincide.

Canonical terms. The point propagator differentiates the Green's tensor: a receiver derivative ∂_z acts on the explicit z and on ζ , as $\partial/\partial z - \partial_\zeta$ for the image and $\partial/\partial z + \partial_\zeta$ for the transmitted field, and similarly for source derivatives and the lateral ones. Collected, the 9×9 static image is a sum of 356 terms

$$c(\lambda_A, \mu_A, \lambda_B, \mu_B) z^A z'^B \partial_{\rho_x}^{\alpha_x} \partial_{\rho_y}^{\alpha_y} \partial_\zeta^{\alpha_z} \Phi_j(\boldsymbol{\rho}, \zeta), \quad (10)$$

in 61 families $(j, \boldsymbol{\alpha})$: Φ_0 with up to four derivatives, and Φ_1 and Φ_2 under lateral derivatives only (one to four, and two to four), since a ζ -derivative of a logarithmic potential is the potential below it. The transmitted field has 275 terms in 46 families of the same kind. Every kernel $\partial^\alpha \Phi_j$ is homogeneous in $(\boldsymbol{\rho}, \zeta)$, of degree $-1 - |\boldsymbol{\alpha}|$, $-|\boldsymbol{\alpha}|$ and $1 - |\boldsymbol{\alpha}|$ for $j = 0, 1, 2$.

A line of images. Every logarithmic term carries a lateral derivative and therefore vanishes as $\zeta \rightarrow \infty$. Integrating the relations between the potentials from ζ to infinity gives

$$\partial^\alpha \Phi_1(\boldsymbol{\rho}, \zeta) = \int_\zeta^\infty \partial^\alpha \Phi_0(\boldsymbol{\rho}, t) dt, \quad \partial^\alpha \Phi_2(\boldsymbol{\rho}, \zeta) = \int_\zeta^\infty (t - \zeta) \partial^\alpha \Phi_0(\boldsymbol{\rho}, t) dt. \quad (11)$$

Mindlin's terms are thus Kelvin's image spread along the vertical half-line beyond the image point, with uniform and with linearly growing density. This is the form in which their moments are computed.

4 Coupling blocks with the static image

The separation form. For two cells in A the image kernel depends on the lateral differences $\rho_h = x_h - x'_h$ but on the vertical sum $\zeta = -(z + z')$. The block is therefore an integral over (ρ_x, ρ_y, ζ) against a weight that is a cross-correlation of the cells' factors in the two lateral coordinates, as in the homogeneous medium, and a convolution in the vertical, $u_{\alpha\beta}(\tau) = \int v^\alpha (\tau - v)^\beta dv$ in $\tau = v + v'$, into which the factors z^A and z'^B of Eq. (10) are expanded. The weights are piecewise polynomials with rational coefficients, computed exactly. The domain divides into boxes on which the weight is a polynomial.

The singular point. The image kernel is singular only at $\boldsymbol{\rho} = 0, \zeta = 0$, which is reached when both cells touch the interface and touch each other laterally (Figure 1a), and then at a vertex of the boxes that hold it. There the vertical weight vanishes as ζ^{1+A+B} , and for every one of the 356 terms the integrand is at worst $O(R^{-2})$: it is absolutely integrable, with no distribution and no derivative to be moved onto the weight. Boxes away from the singular point have smooth integrands and are integrated by Gauss rules in coordinates centred on each box. On a box with the singular point at a vertex, reflected onto the positive octant, every term requires the corner moments

$$M_{j,\alpha}[p, q, r] = \int_0^a \int_0^b \int_0^c \rho_x^p \rho_y^q \zeta^r \partial^\alpha \Phi_j(\boldsymbol{\rho}, \zeta) d\zeta d\rho_y d\rho_x, \quad (12)$$

with $a = b = c = 2h$. They are universal: by homogeneity they scale as a power of h times a number independent of h , the frequency and the media.

Kelvin's potential. For $j = 0$, $\partial^\alpha R^{-1} = P_\alpha(\boldsymbol{\rho}, \zeta) R^{-2|\alpha|-1}$ with P_α a homogeneous polynomial of degree $|\alpha|$, so the moments are combinations of the master integrals

$$I[p, q, r; n] = \int_0^a \int_0^b \int_0^c x^p y^q z^r (x^2 + y^2 + z^2)^{n/2} dz dy dx, \quad (13)$$

here for odd n down to -9 . The integrand is homogeneous of degree $p + q + r + n$ about the singular point, and Euler's identity with the divergence theorem reduces the box to its three far faces, the faces through the singular point carrying no flux; on a face the same step in two dimensions reduces to edges, and on an edge the integral is elementary [1]. One transcendental function enters, through the face integral of R^{-3} : $\int_0^b \int_0^c R^{-3} dz dy = a^{-1} \arctan(bc/(a\sqrt{a^2 + b^2 + c^2}))$ on the face $x = a$, which is continued to lower n by differentiation with respect to a .

The logarithmic potentials. By Eq. (11) the ζ -integral of the box can be done first, which leaves Kelvin's potential against a polynomial weight in t over the box and over the semi-infinite column $[0, a] \times [0, b] \times [c, \infty)$ above it. On $[0, c]$ the weight is $t^{r+1}/(r+1)$ ($j = 1$) or $t^{r+2}/((r+1)(r+2))$ ($j = 2$); on the column it is $c^{r+1}/(r+1)$, or $t c^{r+1}/(r+1) - c^{r+2}/(r+2)$. The column is the semi-infinite box less the finite one. Euler's identity holds on the semi-infinite box, the face at infinity carrying no flux, and its faces reduce to edges that run to infinity,

$$\int_0^\infty z^r (A^2 + z^2)^{n/2} dz = \frac{1}{2} A^{r+n+1} B\left(\frac{r+1}{2}, -\frac{r+n+1}{2}\right), \quad (14)$$

Beta functions at half-integers, elementary. Two cases need care. Where the degree $p + q + r + n + 3$ of a column term is negative, each of the two boxes holds an integral that diverges at the singular corner; Euler's identity then gives Hadamard's finite part of each, and the two finite parts differ by the exact integral over the column, because their singular contributions at the shared corner are equal. Where the degree is zero the identity is degenerate, and the column is reduced instead by integration by parts in t , then in ρ_x and ρ_y , whose boundary terms are faces at height c and

strips at $\rho_x = a$ or $\rho_y = b$, away from the singular point. Two base integrals remain:

$$B_1 = \int_0^a \int_0^b \int_c^\infty R^{-3} dt d\rho_y d\rho_x, \quad B_2 = \int_0^a \int_0^b \int_c^\infty \rho_x \rho_y R^{-5} dt d\rho_y d\rho_x. \quad (15)$$

The base integrals. Both are of degree zero, so they depend only on the ratios of the sides, which here are equal. The t -integrals are elementary, and in polar coordinates (r, θ) in the ρ -plane so are the radial ones: with $c = 1$ and $s = \sqrt{1 + r^2}$,

$$B_1 = \int \log \frac{1+s}{2} d\theta, \quad B_2 = \int \cos \theta \sin \theta \left[\frac{1}{3s} + \frac{2}{3} \log(1+s) - \frac{1}{3} - \frac{2}{3} \log 2 \right] d\theta,$$

with $r = \sec \theta$ on $[0, \pi/4]$ and $r = \csc \theta$ on $[\pi/4, \pi/2]$, the two halves equal by symmetry. For B_2 the substitution $v = \cos^2 \theta$ makes the remaining integral elementary. For B_1 the integral $2 \int_0^{\pi/4} \log(\frac{1}{2}(1 + \sqrt{1 + \sec^2 \theta})) d\theta$ evaluates to Catalan's constant G , $\pi \log 2$, and the imaginary parts of $\text{Li}_2(2^{-1/2} e^{i\pi/12})$ and $\text{Li}_2(2^{-1/2} e^{7i\pi/12})$. Kummer's formula writes $\text{Im Li}_2(re^{i\theta})$ in Clausen functions $\text{Cl}_2(\theta) = \text{Im Li}_2(e^{i\theta})$ through the auxiliary angle $\omega = \arctan(r \sin \theta / (1 - r \cos \theta))$ [8], and at both points $\omega = \pi/6$. The logarithms then cancel, and with $\text{Cl}_2(\pi/2) = G$ and the duplication formula $\text{Cl}_2(2\theta) = 2\text{Cl}_2(\theta) - 2\text{Cl}_2(\pi - \theta)$ at $\theta = \pi/6$,

$$B_1 = \frac{5}{4} \text{Cl}_2\left(\frac{\pi}{3}\right) - G = 0.35271\,14138\dots, \quad B_2 = \frac{1}{3} \log \frac{(1 + \sqrt{2})^2}{2(1 + \sqrt{3})} = 0.02151\,58182\dots \quad (16)$$

Every corner moment is therefore elementary apart from the two constants G and $\text{Cl}_2(\pi/3)$. All of them are universal numbers, computed once and stored.

5 Cells on opposite sides of the interface

For a source cell in A and a field cell in B the static coupling is the transmitted field, the whole of it, with no direct part. Its potentials are those of Eq. (9) with $\zeta = z - z' = |z| + |z'|$, and the singular point is where the two cells touch at the interface (Figure 1b). The 275 canonical terms are integrable at that point, as the image's are. In the separation form the vertical weight is now a cross-correlation in $v - v'$, like the lateral ones, and the corner moments are those of Section 4.

The other arrangements follow by symmetry. A field cell in A with a source in B, and two cells in B, are the problems above reflected in the plane $z = 0$ with the two media exchanged. Under the reflection the displacement u_z and the shear strains γ_{zx}, γ_{zy} change sign, so the block is conjugated by $\text{diag}(-1, 1, 1, 1, 1, 1, 1, -1, -1)$ in the order $(u_z, u_x, u_y, \varepsilon_{zz}, \varepsilon_{xx}, \varepsilon_{yy}, \gamma_{xy}, \gamma_{zx}, \gamma_{zy})$, and each cell polynomial is multiplied by (-1) to its power of ξ_z .

6 Verification

Each step was checked against a route that shares no step with it; Table 2 collects the results. The media are A with $\alpha = 5.0$, $\beta = 3.0 \text{ km s}^{-1}$, $\rho = 2.5 \text{ g cm}^{-3}$ and B with $\alpha = 6.5$, $\beta = 3.6 \text{ km s}^{-1}$,

$\rho = 2.8 \text{ g cm}^{-3}$, and $h = 0.05 \text{ km}$. Errors are relative to the largest entry of the quantity compared, and the corner moments are measured on the scale 2^{p+q+r} of each monomial.

The checks are of four kinds. The spectral solutions were derived independently in two computer-algebra systems, and their spatial forms were compared with direct two-dimensional Fourier integration of the spectral kernels. The corner moments were compared with three routes: Euler's reduction with Gauss rules on the far faces, tanh-sinh quadrature of the singular volume integral itself, and an evaluation at thirty digits that integrates one coordinate exactly and then the other two with Duffy coordinates at the corner [9]. We note that adaptive three-dimensional cubature applied directly with the singular point at a corner, or over the semi-infinite column, can report convergence while in error by as much as 5×10^{-7} , which is why the references integrate one coordinate exactly. The blocks of cells away from the singular point were compared with a tensor Gauss rule over both cells of the canonical kernel, which shares neither the separation form nor the exact weights. Two limits close the chain: for identical media the transmitted blocks must be the touching blocks of the homogeneous medium [1], computed by an unrelated method, and the image must vanish; and the touching block of a cell with its own image must be the limit of the plane-wave integral (6) of the static spectrum as the cutoff grows.

Two remarks qualify the table. The touching reflected blocks agree with the face-quadrature route to 5×10^{-15} , but both routes contract the corner moments with the weights expanded in powers about the singular corner, which in the homogeneous medium costs about one digit at degree two [1]; the only fully independent check of these blocks, the plane-wave integral, is limited to 2.3×10^{-8} by its own convergence. And with identical media the image vanishes identically, to the last bit, on both sides of the interface.

quantity	compared with	agreement
<i>Plane-wave representation</i>		
integrand of Eq. (2)	residue form of the plane-to-plane kernel	3×10^{-15}
blocks (3), planes not touching	closed-form blocks [1]	$2\text{--}6 \times 10^{-13}$
<i>Static image</i>		
reflected spectral tensor	second computer-algebra derivation	5×10^{-16}
dynamic reflected kernel, $\omega/\beta q \rightarrow 0$	static kernel	$\propto (\omega/\beta q)^2$
spatial form	Fourier integral of the spectrum	$7 \times 10^{-13}\text{--}1.3 \times 10^{-12}$
canonical terms (10)	direct differentiation of the spatial kernel	3×10^{-15}
<i>Corner moments</i>		
Euler's reduction, 7 cases, up to 4 derivatives	tanh-sinh quadrature of the volume integral	$6 \times 10^{-16}\text{--}4.8 \times 10^{-14}$
closed forms, all 53 352 moments, monomials to degree (6, 6, 8)	Euler's reduction with Gauss rules on the faces	1.2×10^{-15}
13 moments, all potentials, 3 degenerate columns	thirty-digit evaluation	$5 \times 10^{-23}\text{--}1.6 \times 10^{-16}$
B_1, B_2 , Eq. (16)	forty-digit angular quadrature	10^{-40}
<i>Reflected blocks</i>		
cells away from the singular point, 4 pairs, degree-one and degree-two fields	tensor Gauss rule over both cells	$3 \times 10^{-15}\text{--}8.5 \times 10^{-15}$
touching blocks, closed-form corner moments, 3 pairs	Gauss rules on the far faces	$1.8\text{--}5.3 \times 10^{-15}$
cell with its own image	static plane-wave integral, $Qh = 320$ to $20\,480$	$7.9 \times 10^{-6} \rightarrow 2.3 \times 10^{-8}$
<i>Transmitted field and blocks</i>		
transmitted spectral tensor	second computer-algebra derivation	4.2×10^{-16}
spatial form	Fourier integral of the spectrum	3.9×10^{-16}
canonical terms, different media	Fourier integral of the spectrum	$6.6\text{--}7.8 \times 10^{-13}$
identical media: face, edge, corner contact; mirrored; degree two	homogeneous-medium blocks [1]	$1.9\text{--}7.6 \times 10^{-15}$
different media, cells not touching	tensor Gauss rule over both cells	$1.7\text{--}5.0 \times 10^{-13}$

Table 2. Independent checks. The last entry of each group of blocks is limited by the accuracy of the comparison route: the Gauss rule over both cells at order 10 to about 10^{-13} , and the static plane-wave integral by its algebraic convergence.

7 Discussion

The static image carries the whole singularity of the coupling at a welded interface, and in closed form it costs no more than the touching blocks of a homogeneous medium: a fixed set of universal corner moments, contracted with exact rational weights and with coefficients rational in the Lamé constants. The dynamic remainder is smooth, and the plane waves of Section 2 carry it at a cost that does not depend on how close the cells are to the interface, since the subtraction has removed what made the integral converge slowly.

For a stack of layers the same construction applies interface by interface. The singular part of the reflected and transmitted couplings comes only from the interface that the cells touch, and is the static image of that interface; reflections from other interfaces reach a cell after travelling at least one layer thickness and are smooth. The plane-wave representation extends to the stack through the reflection and transmission matrices of the layered medium, composed by the recursions of the reflectivity method [2, 3]. The coupling of every cell with every other is then available from the up- and down-going factorisation alone, without tabulating the stratified Green’s tensor, which is the storage advantage that motivates the approach; a two-way march in depth that applies the volume operator in this way is the subject of further work.

Two limitations remain. The corner moments are contracted with the weights in the power basis about the singular corner; the Legendre basis that restores full double-precision accuracy in the homogeneous medium [1] has not yet been applied to the image boxes, and the touching reflected blocks are correspondingly expected to lose about one digit at degree two. And the closed form of the base integrals is stated for cubic corner boxes, $a = b = c$, which is all a grid of cubes requires; for unequal sides the angular integrals are evaluated by quadrature.

Data availability

The Mathematica notebooks and Python scripts that derive and check every result, and the package they test, are available at <https://github.com/tmnestor/MultipleScatteringCalculations>: the static image and transmitted field, `Mathematica/StaticInterfaceImage.wl`, `Mathematica/StaticImageSpace.wl` and `Mathematica/StaticInterfaceTransmission.wl`, with their Python twins `scripts/derive_static_interface_image.py`, `scripts/derive_static_image_space.py` and `scripts/derive_static_interface_transmission.py`; the canonical terms, `scripts/derive_interface_image_terms.py`; the corner moments and base integrals, `Mathematica/ImageCornerMoments.wl`, `scripts/derive_image_corner_moments.py` and `cubic_scattering/image_moments.py`; the blocks, `cubic_scattering/interface_image.py`; the plane-wave representation, reflection and transmission, `cubic_scattering/stratified_march.py`; and the checks of Table 2, `scripts/gate_march_modes.py` and `scripts/gate_interface_image.py`.

Acknowledgments

This work began thirty years ago in the author’s doctoral research at the Research School of Earth Sciences of the Australian National University, Canberra [4].

Declaration of generative AI and AI-assisted technologies in the manuscript preparation process

Software development. Software development followed a human-directed, AI-assisted workflow. The author designed the software originally, but only directed and validated recent improvements, which build on the author’s original Fortran implementation (the [PhD_fortran_code](#) folder of the repository given under Data availability) and an extensive library of Mathematica notebooks, developed and extended since 1992, that carry out and verify the symbolic derivations; an agentic coding assistant (Anthropic Claude) operated under human-in-the-loop supervision to implement specified changes and diagnostic tests. The design, the choice of numerical methods, and the resolution of the principal technical difficulties were the author’s. All AI-generated changes were subject to code review by the author and validated by automated tests and independent checks.

Manuscript preparation. During the preparation of this work, the author used Claude (Anthropic) in order to search for relevant open-access research, to write the TikZ code of the diagrams and to help redraft the paper into more coherent sections for readability. After using this tool, the author reviewed and edited the content as needed and takes full responsibility for the content of the published article.

References

- [1] T. M. Nestor. The coupling integrals of polynomial cubes in closed form: Exact for elastic waves, and accurate in double precision. In preparation, 2026.
- [2] K. Fuchs and G. Müller. Computation of synthetic seismograms with the reflectivity method and comparison with observations. *Geophysical Journal of the Royal Astronomical Society*, 23(4):417–433, 1971.
- [3] B. L. N. Kennett. *Seismic Wave Propagation in Stratified Media*. Cambridge University Press, Cambridge, 1983.
- [4] T. M. Nestor. *Seismic Wave Propagation in Complex Media: Summing the Multiple Scattering Series*. PhD thesis, Research School of Earth Sciences, Australian National University, Canberra, ACT, Australia, 1996.
- [5] T. M. Nestor. The single-site T-matrix of a Legendre cell in closed form: Elastic waves, every power of the frequency on six constants. In preparation, 2026.

- [6] L. Rongved. Force interior to one of two joined semi-infinite solids. In *Proceedings of the Second Midwestern Conference on Solid Mechanics*, pages 1–13, 1955.
- [7] R. D. Mindlin. Force at a point in the interior of a semi-infinite solid. *Physics*, 7(5):195–202, 1936.
- [8] Leonard Lewin. *Polylogarithms and Associated Functions*. North-Holland, New York, 1981.
- [9] M. G. Duffy. Quadrature over a pyramid or cube of integrands with a singularity at a vertex. *SIAM Journal on Numerical Analysis*, 19(6):1260–1262, 1982. doi: 10.1137/0719090.