

# In Search of the Perfect Fluid

Thomas Schaefer, North Carolina State University



See T. Schäfer, D. Teaney, “Perfect Fluidity” [arXiv:0904.3107]


## The bottom-line

Remarkably, the best fluids that have been observed are the *coldest* and the *hottest* fluid ever created in the laboratory, cold atomic gases ( $10^{-6}\text{K}$ ) and the quark gluon plasma ( $10^{12}\text{K}$ ) at RHIC.

Both of these fluids come close to a bound on the shear viscosity that was first proposed based on calculations in string theory, involving non-equilibrium evolution of black holes in 5 (and more) dimensions.


# Measures of Perfection

Viscosity determines shear stress (“friction”) in fluid flow



$$F = A \eta \frac{\partial v_x}{\partial y}$$

Dimensionless measure of shear stress: Reynolds number



$$Re = \underbrace{\frac{n}{\eta}}_{\text{fluid property}} \times \underbrace{mvr}_{\text{flow property}}$$

- $[\eta/n] = \hbar$

- Relativistic systems  $Re = \frac{s}{\eta} \times \tau T$


Other sources of dissipation (thermal conductivity, bulk viscosity, ...) vanish for certain fluids, but shear viscosity is always non-zero.

There are reasons to believe that  $\eta$  is bounded from below by a constant times  $\hbar s/k_B$ . In a large class of theories  $\eta/s \geq \hbar/(4\pi k_B)$ .


A fluid that saturates the bound is a “perfect fluid”.

# Fluids: Gases, Liquids, Plasmas, ...

Hydrodynamics: Long-wavelength, low-frequency dynamics of conserved or spontaneously broken symmetry variables.




$$\tau \sim \tau_{micro}$$



$$\tau \sim \lambda^{-1}$$

Historically: Water  
 $(\rho, \epsilon, \vec{\pi})$



Simple fluid: Conservation laws for mass, energy, momentum

$$\frac{\partial \rho}{\partial t} + \vec{\nabla}(\rho \vec{v}) = 0$$

$$\frac{\partial \epsilon}{\partial t} + \vec{\nabla} \vec{j}^\epsilon = 0$$

$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0$$



Constitutive relations: Energy momentum tensor

$$\Pi_{ij} = P\delta_{ij} + \rho v_i v_j + \eta \left( \partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right) + O(\partial^2)$$

reactive

dissipative

2nd order

Measure of perfection:


$$\frac{\delta \Pi_{ij}}{\Pi_{ij}} \sim \frac{\eta(\partial \cdot v)}{P} \sim \frac{\eta}{s} \frac{1}{\tau T}$$

# Kinetic Theory

Kinetic theory: conserved quantities carried by quasi-particles

$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p + \vec{F} \cdot \vec{\nabla}_p f_p = C[f_p]$$

$$\eta \sim \frac{1}{3} n \bar{p} l_{mfp}$$



Normalize to density. Uncertainty relation suggests

$$\frac{\eta}{n} \sim \bar{p} l_{mfp} \geq \hbar$$

Also:  $s \sim k_B n$  and  $\eta/s \geq \hbar/k_B$

Validity of kinetic theory as  $\bar{p} l_{mfp} \sim \hbar$ ?

# Effective Theories for Fluids (Here: Weak Coupling QCD)



$$\mathcal{L} = \bar{q}_f (i \not{D} - m_f) q_f - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a$$



$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p = C[f_p] \quad (\omega < T)$$



$$\frac{\partial}{\partial t} (\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad (\omega < g^4 T)$$



# Holographic Duals: Transport Properties





Thermal (conformal) field theory  $\equiv$   $AdS_5$  black hole

CFT entropy  $\Leftrightarrow$  Hawking-Bekenstein entropy  
 $\sim$  area of event horizon

shear viscosity  $\Leftrightarrow$  Graviton absorption cross section  
 $\sim$  area of event horizon

$$T_{\mu\nu} = \frac{1}{\sqrt{-g}} \frac{\delta \mathcal{L}}{\delta g_{\mu\nu}}$$

$$g_{\mu\nu} = g_{\mu\nu}^0 + \gamma_{\mu\nu}$$



# Holographic Duals: Transport Properties

Thermal (conformal) field theory  $\equiv$   $AdS_5$  black hole


CFT entropy  $\Leftrightarrow$  Hawking-Bekenstein entropy  
 $\sim$  area of event horizon

shear viscosity  $\Leftrightarrow$  Graviton absorption cross section  
 $\sim$  area of event horizon

Strong coupling limit

$$\frac{\eta}{s} = \frac{\hbar}{4\pi k_B}$$

Son and Starinets (2001)



Strong coupling limit universal? Provides lower bound for all theories?

# Effective Theories (Strong coupling)




$$\mathcal{L} = \bar{\lambda}(i\sigma \cdot D)\lambda - \frac{1}{4}G_{\mu\nu}^a G_{\mu\nu}^a + \dots \Leftrightarrow S = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} \mathcal{R} + \dots$$




$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad (\omega < T)$$

## Kinetics vs No-Kinetics




AdS/CFT low viscosity goo




pQCD kinetic plasma

# Kinetics vs No-Kinetics

Spectral function  $\rho(\omega) = \text{Im}G_R(\omega, 0)$  associated with  $T_{xy}$



weak coupling QCD



strong coupling AdS/CFT

transport peak vs no transport peak

## Perfect Fluids: How to be a contender?

Bound is quantum mechanical

need quantum fluids



Bound is incompatible with weak coupling and kinetic theory

strong interactions, no quasi-particles

Model system has conformal invariance (essential?)

(Almost) scale invariant systems

## Perfect Fluids: The contenders





QGP ( $T=180$  MeV)

Trapped Atoms  
( $T=0.1$  neV)



Liquid Helium  
( $T=0.1$  meV)

# Perfect Fluids: The contenders



QGP  $\eta = 5 \cdot 10^{11} Pa \cdot s$

Trapped Atoms

$\eta = 1.7 \cdot 10^{-15} Pa \cdot s$



Liquid Helium

$\eta = 1.7 \cdot 10^{-6} Pa \cdot s$

Consider ratios

$\eta/s$



# Kinetic Theory: Quasiparticles



low temperature

high temperature

unitary gas

phonons



atoms



helium

phonons, rotons



atoms






QCD

pions

quarks, gluons



# Theory Summary



# I. Experiment (Liquid Helium)





FIG. 1. The viscosity of liquid helium II measured by flow through a  $10^{-4}$  cm channel.

Kapitza (1938)

viscosity vanishes below  $T_c$   
capillary flow viscometer



Hollis-Hallett (1955)


roton minimum, phonon rise  
rotation viscometer

$$\eta/s \simeq 0.8 \hbar/k_B$$

## II. Collective Modes (Fermions)


Radial breathing mode

Ideal fluid hydrodynamics ( $P \sim n^{5/3}$ )



$$\frac{\partial n}{\partial t} + \vec{\nabla} \cdot (n \vec{v}) = 0$$

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\frac{\vec{\nabla} P}{mn} - \frac{\vec{\nabla} V}{m}$$



Hydro frequency at unitarity

$$\omega = \sqrt{\frac{10}{3}} \omega_{\perp}$$

Damping small, depends on  $T/T_F$ .

# Viscous Hydrodynamics


Energy dissipation ( $\eta, \zeta, \kappa$ : shear, bulk viscosity, heat conductivity)

$$\begin{aligned}\dot{E} = & -\frac{1}{2} \int d^3x \eta(x) \left( \partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right)^2 \\ & - \int d^3x \zeta(x) (\partial_i v_i)^2 - \frac{1}{T} \int d^3x \kappa(x) (\partial_i T)^2\end{aligned}$$







Shear viscosity to entropy ratio  
(assuming  $\zeta = \kappa = 0$ )

$$\frac{\eta}{s} = (3\lambda N)^{\frac{1}{3}} \frac{\Gamma}{\omega_{\perp}} \frac{E_0}{E_F} \frac{N}{S}$$


Schaefer (2007), see also Bruun, Smith



# Dissipation



# Dissipation





$$\left\{ \begin{array}{l} (\delta t_0)/t_0 \\ (\delta a)/a \end{array} \right\} = \left\{ \begin{array}{l} 0.008 \\ 0.024 \end{array} \right\} \left( \frac{\langle \alpha_s \rangle}{1/(4\pi)} \right) \left( \frac{2 \cdot 10^5}{N} \right)^{1/3} \left( \frac{S/N}{2.3} \right) \left( \frac{0.85}{E_0/E_F} \right)$$

$t_0$ : “Crossing time” ( $b_{\perp} = b_z$ ,  $\theta = 45^{\circ}$ )

$a$ : amplitude

### III. Elliptic Flow (QGP)

Hydrodynamic  
expansion converts  
coordinate space  
anisotropy  
to momentum space  
anisotropy



source: U. Heinz (2005)




# Viscosity and Elliptic Flow

Consistency condition  $T_{\mu\nu} \gg \delta T_{\mu\nu}$   
(applicability of Navier-Stokes)

$$\frac{\eta + \frac{4}{3}\zeta}{s} \ll \frac{3}{4}(\tau T)$$

Danielewicz, Gyulassy (1985)

Very restrictive for  $\tau < 1$  fm



Romatschke (2007), Teaney (2003)

Many questions: Dependence on initial conditions, freeze out, etc.

## Outlook

Too early to declare a winner.

$$\eta/s \simeq 0.8 \text{ (He)}, \quad \eta/s \leq 0.5 \text{ (CA)}, \quad \eta/s \leq 0.5 \text{ (QGP)}$$

Other experimental constraints, more analysis needed.

Kinetic theory: o.k. in He (all  $T$ ), o.k. close to  $T_c$  in CA, QGP?

New theory tools: AdS/Cold Atom correspondence? Field theory approaches in cross over regime (large  $N$ , epsilon expansions, ...)