

In Search of the Perfect Fluid

Thomas Schaefer, North Carolina State University



See T. Schäfer, D. Teaney, “Perfect Fluidity” [arXiv:0904.3107]


The bottom-line

Remarkably, the best fluids that have been observed are the *coldest* and the *hottest* fluid ever created in the laboratory, cold atomic gases (10^{-6}K) and the quark gluon plasma (10^{12}K) at RHIC.

Both of these fluids come close to a bound on the shear viscosity that was first proposed based on calculations in string theory, involving non-equilibrium evolution of black holes in 5 (and more) dimensions.


Measures of Perfection

Viscosity determines shear stress (“friction”) in fluid flow



$$F = A \eta \frac{\partial v_x}{\partial y}$$

Dimensionless measure of shear stress: Reynolds number



$$Re = \underbrace{\frac{n}{\eta}}_{\text{fluid property}} \times \underbrace{mvr}_{\text{flow property}}$$

- $[\eta/n] = \hbar$

- Relativistic systems $Re = \frac{s}{\eta} \times \tau T$


Other sources of dissipation (thermal conductivity, bulk viscosity, ...) vanish for certain fluids, but shear viscosity is always non-zero.

There are reasons to believe that η is bounded from below by a constant times $\hbar s/k_B$. In a large class of theories $\eta/s \geq \hbar/(4\pi k_B)$.


A fluid that saturates the bound is a “perfect fluid”.

Fluids: Gases, Liquids, Plasmas, ...

Hydrodynamics: Long-wavelength, low-frequency dynamics of conserved or spontaneously broken symmetry variables.




$$\tau \sim \tau_{micro}$$



$$\tau \sim \lambda^{-1}$$

Historically: Water
 $(\rho, \epsilon, \vec{\pi})$



Example: Simple Fluid

Conservation laws: mass, energy, momentum

$$\frac{\partial \rho}{\partial t} + \vec{\nabla}(\rho \vec{v}) = 0$$

$$\frac{\partial \epsilon}{\partial t} + \vec{\nabla} \cdot \vec{j}^\epsilon = 0$$

$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0$$

[Euler/Navier-Stokes equation]



Constitutive relations: Energy momentum tensor

$$\Pi_{ij} = P\delta_{ij} + \rho v_i v_j + \eta \left(\partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right) + O(\partial^2)$$

reactive

dissipative


2nd order

Kinetic Theory

Kinetic theory: conserved quantities carried by quasi-particles

$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p + \vec{F} \cdot \vec{\nabla}_p f_p = C[f_p]$$

$$\eta \sim \frac{1}{3} n \bar{p} l_{mfp}$$



Normalize to density. Uncertainty relation suggests

$$\frac{\eta}{n} \sim \bar{p} l_{mfp} \geq \hbar$$

Also: $s \sim k_B n$ and $\eta/s \geq \hbar/k_B$

Validity of kinetic theory as $\bar{p} l_{mfp} \sim \hbar$?

Effective Theories for Fluids (Here: Weak Coupling QCD)



$$\mathcal{L} = \bar{q}_f (i \not{D} - m_f) q_f - \frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a$$



$$\frac{\partial f_p}{\partial t} + \vec{v} \cdot \vec{\nabla}_x f_p = C[f_p] \quad (\omega < T)$$



$$\frac{\partial}{\partial t} (\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad (\omega < g^4 T)$$


Holographic Duals: Transport Properties

Thermal (conformal) field theory \equiv AdS_5 black hole

CFT entropy \Leftrightarrow Hawking-Bekenstein entropy
 \sim area of event horizon

shear viscosity \Leftrightarrow Graviton absorption cross section
 \sim area of event horizon

$$T_{\mu\nu} = \frac{\delta S}{\delta g_{\mu\nu}} \quad g_{\mu\nu} = g_{\mu\nu}^0 + \gamma_{\mu\nu}$$



Holographic Duals: Transport Properties

Thermal (conformal) field theory \equiv AdS_5 black hole


CFT entropy \Leftrightarrow Hawking-Bekenstein entropy
 \sim area of event horizon

shear viscosity \Leftrightarrow Graviton absorption cross section
 \sim area of event horizon

Strong coupling limit

$$\frac{\eta}{s} = \frac{\hbar}{4\pi k_B}$$

Son and Starinets (2001)



Strong coupling limit universal? Provides lower bound for all theories?

Effective Theories (Strong coupling)




$$\mathcal{L} = \bar{\lambda}(i\sigma \cdot D)\lambda - \frac{1}{4}G_{\mu\nu}^a G_{\mu\nu}^a + \dots \Leftrightarrow S = \frac{1}{2\kappa_5^2} \int d^5x \sqrt{-g} \mathcal{R} + \dots$$




$$\frac{\partial}{\partial t}(\rho v_i) + \frac{\partial}{\partial x_j} \Pi_{ij} = 0 \quad (\omega < T)$$

Kinetics vs No-Kinetics

Spectral function $\rho(\omega) = \text{Im}G_R(\omega, 0)$ associated with T_{xy}



weak coupling QCD



strong coupling AdS/CFT

transport peak vs no transport peak

Perfect Fluids: How to be a contender?

Bound is quantum mechanical

need quantum fluids



Bound is incompatible with weak coupling and kinetic theory

strong interactions, no quasi-particles

Model system has conformal invariance (essential?)

(Almost) scale invariant systems

Perfect Fluids: The contenders





QGP ($T=180$ MeV)

Trapped Atoms
($T=0.1$ neV)



Liquid Helium
($T=0.1$ meV)

Perfect Fluids: The contenders



QGP $\eta = 5 \cdot 10^{11} Pa \cdot s$

Trapped Atoms

$\eta = 1.7 \cdot 10^{-15} Pa \cdot s$



Liquid Helium

$\eta = 1.7 \cdot 10^{-6} Pa \cdot s$

Consider ratios

η/s

Kinetic Theory: Quasiparticles



low temperature

high temperature

unitary gas

phonons



atoms



helium

phonons, rotons



atoms






QCD

pions

quarks, gluons



Theory Summary



I. Experiment (Liquid Helium)





FIG. 1. The viscosity of liquid helium II measured by flow through a 10^{-4} cm channel.

Kapitza (1938)

viscosity vanishes below T_c
capillary flow viscometer



Hollis-Hallett (1955)


roton minimum, phonon rise
rotation viscometer

$$\eta/s \simeq 0.8 \hbar/k_B$$

II. Collective Modes (Fermions)


Radial breathing mode

Ideal fluid hydrodynamics ($P \sim n^{5/3}$)



$$\frac{\partial n}{\partial t} + \vec{\nabla} \cdot (n \vec{v}) = 0$$

$$\frac{\partial \vec{v}}{\partial t} + (\vec{v} \cdot \vec{\nabla}) \vec{v} = -\frac{\vec{\nabla} P}{mn} - \frac{\vec{\nabla} V}{m}$$



Hydro frequency at unitarity

$$\omega = \sqrt{\frac{10}{3}} \omega_{\perp}$$

Damping small, depends on T/T_F .

Viscous Hydrodynamics


Energy dissipation (η, ζ, κ : shear, bulk viscosity, heat conductivity)

$$\begin{aligned} \dot{E} = & -\frac{1}{2} \int d^3x \eta(x) \left(\partial_i v_j + \partial_j v_i - \frac{2}{3} \delta_{ij} \partial_k v_k \right)^2 \\ & - \int d^3x \zeta(x) (\partial_i v_i)^2 - \frac{1}{T} \int d^3x \kappa(x) (\partial_i T)^2 \end{aligned}$$







Shear viscosity to entropy ratio
(assuming $\zeta = \kappa = 0$)

$$\frac{\eta}{s} = \frac{3}{4} \xi^{\frac{1}{2}} (3N)^{\frac{1}{3}} \frac{\Gamma}{\omega_{\perp}} \frac{\bar{\omega}}{\omega_{\perp}} \frac{N}{S}$$

Schaefer (2007), see also Bruun, Smith





Dissipation



III. Elliptic Flow (QGP)

Hydrodynamic
expansion converts
coordinate space
anisotropy
to momentum space
anisotropy



source: U. Heinz (2005)


Viscosity and Elliptic Flow

Consistency condition $T_{\mu\nu} \gg \delta T_{\mu\nu}$
(applicability of Navier-Stokes)

$$\frac{\eta + \frac{4}{3}\zeta}{s} \ll \frac{3}{4}(\tau T)$$

Danielewicz, Gyulassy (1985)

Very restrictive for $\tau < 1$ fm



Romatschke (2007), Teaney (2003)

Many questions: Dependence on initial conditions, freeze out, etc.

Outlook

Too early to declare a winner.

$$\eta/s \simeq 0.8 \text{ (He)}, \quad \eta/s \leq 0.5 \text{ (CA)}, \quad \eta/s \leq 0.5 \text{ (QGP)}$$

Other experimental constraints, more analysis needed.

Kinetic theory: o.k. in He (all T), o.k. close to T_c in CA, QGP?

New theory tools: AdS/Cold Atom correspondence? Field theory approaches in cross over regime (large N , epsilon expansions, ...)