24–30
08.2013

ВОРКШОП СОЮЗПОЛИГРАФМАШ

КАРТИРОВАНИЕ ПОТЕНЦИАЛА ПРОИЗВОДСТВЕННЫХ ПРОСТРАНСТВ В ГОРОДЕ

Данияр ЮСУПОВ
Дмитрий ВОРОБЬЕВ

u:lab.spb / СПбГАСУ ДАС
v-cafe / ЦНСИ

makeitcenter.com | +7 921 5563537 | make@makeitcenter.com

bing | NOKIA
Maps by

Maps by
bing | NOKIA

100 feet
25 m
© 2013 Microsoft Corporation
Pictometry Bird's Eye © 2012 Pictometry International Corp
Image courtesy of Simmon

99
Bahnanlage
ROW G 208
LOS TX 244
B PD 6914

Soundcraft

посетители ботанического сада
жители окрестных микрорайонов
прохожие
посетители концертов и мероприятий
студенты

population

- 0
- 1
- 2
- 3

Function

- art
- education
- office
- service
- retail
- restaurant
- entertainment
- production
- empty

























metronaps
Google

Terrace
Please close door!

D Dubois

D Dubois

CONTAMINANT		MAXIMUM LEVELS OF CONTAMINANT FOR: Multi Family Housing.. ..Recreation..Park	Single Family Houses..Gardening.. Playground	Farming Animals.. ..Growing Food		TYPICAL PLANTS
As	Arsenic	16ppm	16ppm	13ppm	Often found in lead-acid batteries, light-emitting diodes, paints, dyes, metals, pharmaceuticals, pesticides, herbicides, soaps, and semiconductors.	Chinese Brake Fern *Pteris vittata L.*; Sunflower *Helianthus annuus*; Highland Bent Grass *Agrostis castellana*
Cr	Chromium	180ppm	36ppm	30ppm	PHYTO EXTRACTION	Alpine Pennycress *Thlaspi caerulescens*; Giant Duckweed *Spirdela polyrhiza*
Pb	Lead	400ppm	400ppm	63ppm	PHYTO STABILIZATION	Blue Sheep Fescue *Festuca ovina*; Indian mustard *Brassica juncea L.*; Common Wheat *Triticum estivum*; Common Ragweed Brassica olercea
Hg	Mercury	0.81ppm	0.81ppm	0.18ppm	PHYTO STABILIZATION	Seapink Thrift *Armeria maritima*; Rapeseed Plant Brassica napus
PCB	Polychlorinated biphenyls	1ppm	1ppm	0.1ppm	Colorless to light yellow oily liquids or waxy solids. Accumulate in fish and marine mammals at much higher levels than in sediments and water	Paul's Scarlet Rose *Rosa*; Zuchini *Curcurbita pepo*
TCE	Trichloroethlene	21ppm	10ppm	0.47ppm	PHYTO DEGRADATION	Willow *Salix*; Eastern cottonwoods *Populus deltoides*
MTBE	Methyl tertiary butyl ether	100ppm	62ppm	0.93ppm	Typically used as a fuel additive in gasoline. Common in areas that were exposed to leakage from the gasoline storage and distribution systems.	Pine *Pinus*
DDT	Dichlorophenyltrichloroethane	7.9ppm	1.7ppm	0.0033ppm		Pumpkin *Curcurbita*
PCP	Pentachlorophenol	6.7ppm	2.4ppm	0.8ppm	PHYTO DEGRADATION	White rot fungus *Phanerochaete chrysosporium*; Crested Wheatgrass *Agropyron cristatum*

Bubble Tree




Betahaus
SEAT
SEAT
SEAT
SEAT
SEAT
SEAT
SEAT
SEAT
TABLE

Projects



Bühne 6
57590

©WWW.INDUSTRIE-IKONEN.DE

photopics

Diebels





QUEEN ELIZABETH HALL
PURCELL ROOM

N°0 L'INUSABLE

FORCE
5 TONNES

LOXAM
LA FABRIQUE



24–30
08.2013

ВОРКШОП СОЮЗПОЛИГРАФМАШ

КАРТИРОВАНИЕ ПОТЕНЦИАЛА ПРОИЗВОДСТВЕННЫХ ПРОСТРАНСТВ В ГОРОДЕ

Данияр ЮСУПОВ
Дмитрий ВОРОБЬЕВ

u:lab.spb / СПбГАСУ ДАС
v-cafe / ЦНСИ

makeitcenter.com | +7 921 5563537 | make@makeitcenter.com