

2.25 Solution: (a) The Dirac delta function on the two-dimensional surface with restricted angular range $[0, \beta]$ can be written as

$$\delta(\boldsymbol{\rho} - \boldsymbol{\rho}') = \frac{1}{\rho} \delta(\rho - \rho') \delta(\phi - \phi') = \frac{1}{\rho} \delta(\rho - \rho') \cdot \frac{2}{\beta} \sum_{m=1}^{\infty} \sin(m\pi\phi/\beta) \sin(m\pi\phi'/\beta),$$

and the corresponding Green function must have a similar form,

$$G(\boldsymbol{\rho}, \boldsymbol{\rho}') = \frac{2}{\beta} \sum_{m=1}^{\infty} g_m(\rho, \rho') \sin(m\pi\phi/\beta) \sin(m\pi\phi'/\beta). \quad (1)$$

Applying the Laplace operator, we have

$$\nabla_{\boldsymbol{\rho}'}^2 G(\boldsymbol{\rho}, \boldsymbol{\rho}') = \frac{1}{\rho'} \frac{\partial}{\partial \rho'} \left(\rho' \frac{\partial G}{\partial \rho'} \right) + \frac{1}{\rho'^2} \frac{\partial^2 G}{\partial \phi'^2} = -4\pi \delta(\boldsymbol{\rho} - \boldsymbol{\rho}').$$

With Eq. (1), we have

$$\begin{aligned} \sum_{m=1}^{\infty} \left[\frac{1}{\rho'} \frac{\partial}{\partial \rho'} \left(\rho' \frac{\partial}{\partial \rho'} \right) - \frac{m^2 \pi^2}{\beta^2} \right] g_m(\rho, \rho') \sin(m\pi\phi/\beta) \sin(m\pi\phi'/\beta) \\ = -4\pi \cdot \frac{1}{\rho} \delta(\rho - \rho') \cdot \sum_{m=1}^{\infty} \sin(m\pi\phi/\beta) \sin(m\pi\phi'/\beta). \end{aligned}$$

The above equation must be valid term-wise, which leads to a differential equation for g_m ,

$$\left[\frac{1}{\rho'} \frac{\partial}{\partial \rho'} \left(\rho' \frac{\partial}{\partial \rho'} \right) - \frac{m^2 \pi^2}{\beta^2} \right] g_m(\rho, \rho') = -4\pi \cdot \frac{1}{\rho} \delta(\rho - \rho').$$

Now, we can apply the familiar procedure to determine g_m . First, solutions to g_m must be linear combinations of $\rho'^{m\pi/\beta}$ and $\rho'^{-m\pi/\beta}$. For $\rho' < \rho$, regularity at $\rho' = 0$ indicates that we can only choose the $\rho'^{m\pi/\beta}$ solution, or

$$g_m(\rho, \rho') \propto \rho'^{m\pi/\beta}.$$

Similarly, for $\rho' > \rho$,

$$g_m(\rho, \rho') \propto \rho'^{-m\pi/\beta}.$$

The final solution must be symmetric in ρ and ρ' , which means

$$g_m(\rho, \rho') = C_m \left(\frac{\rho_{<}}{\rho_{>}} \right)^{m\pi/\beta}.$$

where $\rho_{<}$ ($\rho_{>}$) is the smaller (larger) of ρ and ρ' . Integrate the differential equation governing g_m in a small interval around ρ , we can obtain a relation that connects the derivative of g_m on both side of ρ ,

$$\rho' \frac{\partial}{\partial \rho'} g_m(\rho, \rho') \Big|_{\rho-\varepsilon}^{\rho+\varepsilon} = -4\pi.$$

For $\rho' = \rho + \varepsilon$,

$$\rho' \frac{\partial}{\partial \rho'} g_m(\rho, \rho') \Big|_{\rho'=\rho+\varepsilon} = C_m \rho' \frac{\partial}{\partial \rho'} \left(\frac{\rho}{\rho'} \right)^{m\pi/\beta} \Big|_{\rho'=\rho+\varepsilon} = -C_m \frac{m\pi}{\beta},$$

and for $\rho' = \rho - \varepsilon$,

$$\rho' \frac{\partial}{\partial \rho'} g_m(\rho, \rho') \Big|_{\rho'=\rho-\varepsilon} = C_m \rho' \frac{\partial}{\partial \rho'} \left(\frac{\rho'}{\rho} \right)^{m\pi/\beta} \Big|_{\rho'=\rho-\varepsilon} = C_m \frac{m\pi}{\beta},$$

which leads to

$$\rho' \frac{\partial}{\partial \rho'} g_m(\rho, \rho') \Big|_{\rho-\varepsilon}^{\rho+\varepsilon} = -2C_m \frac{m\pi}{\beta} = -4\pi,$$

or,

$$C_m = \frac{2\beta}{m}.$$

Put the solution of g_m back into Eq. (1), the Green function becomes

$$G(\rho, \phi; \rho', \phi') = 4 \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{m\pi/\beta} \sin(m\pi\phi/\beta) \sin(m\pi\phi'/\beta).$$

(b) Using the identity

$$\sin(m\pi\phi/\beta) \sin(m\pi\phi'/\beta) = \frac{1}{2} \left(\cos[m(\phi - \phi')/\beta] - \cos[m(\phi + \phi')/\beta] \right),$$

the Green function can be expressed as

$$\begin{aligned} G(\rho, \phi; \rho', \phi') &= 2 \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{m\pi/\beta} \left(\cos[m(\phi - \phi')/\beta] - \cos[m(\phi + \phi')/\beta] \right) \\ &= 2 \cdot \text{Re} \left\{ \sum_{m=1}^{\infty} \frac{1}{m} \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{m\pi/\beta} \left(e^{im(\phi - \phi')/\beta} - e^{im(\phi + \phi')/\beta} \right) \right\} \\ &= -2 \cdot \text{Re} \left\{ \log \left(1 - \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{\pi/\beta} e^{i(\phi - \phi')/\beta} \right) - \log \left(1 - \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{\pi/\beta} e^{i(\phi + \phi')/\beta} \right) \right\} \\ &= \log \left(1 + \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{2\pi/\beta} - 2 \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{\pi/\beta} \cos(\phi + \phi') \right) \\ &\quad - \log \left(1 + \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{2\pi/\beta} - 2 \left(\frac{\rho_{\leq}}{\rho_{>}} \right)^{\pi/\beta} \cos(\phi - \phi') \right) \\ &= \log \left(\frac{(\rho)^{2\pi/\beta} + (\rho')^{2\pi/\beta} - 2(\rho\rho')^{\pi/\beta} \cos(\phi + \phi')}{(\rho)^{2\pi/\beta} + (\rho')^{2\pi/\beta} - 2(\rho\rho')^{\pi/\beta} \cos(\phi - \phi')} \right), \end{aligned}$$

where we have used the MacLaurin series for $\log(1 - x)$,

$$\log(1 - x) = - \sum_{n=1}^{\infty} \frac{x^n}{n}.$$