

3.8 Solution: (a) Comparing Eq. (3.136) from the book and the alternative form, we need to establish the following identity,

$$\log \left(\frac{2}{\sin \theta} \right) = \sum_{l=0}^{\infty} A_l P_l(\cos \theta),$$

with $A_0 = 1$, $A_{2j-1} = 0$, and

$$A_{2j} = \frac{4j+1}{2j(2j+1)},$$

for $j > 0$.

Multiplying both sides by $\sin \theta P_l(\cos \theta)$ and integrate with respect to θ between $[0, \pi]$, we have

$$A_l = \frac{2l+1}{2} \int_0^\pi \sin \theta \log \left(\frac{2}{\sin \theta} \right) P_l(\cos \theta) d\theta = \frac{2l+1}{2} \int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) P_l(x) dx.$$

For $l = 0$, $P_0(x) \equiv 1$, the coefficient A_0 can be directly determined,

$$\begin{aligned} A_0 &= \frac{1}{2} \int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) dx \\ &= \log 2 - \frac{1}{4} \left((1+x) \log(1+x) - (1-x) \log(1-x) - 2x \right) \Big|_{-1}^1 \\ &= 1. \end{aligned}$$

When $l \neq 0$, using the ordinary Legendre differential equation (3.10)

$$\frac{d}{dx} \left[(1-x^2) \frac{dP_l(x)}{dx} \right] + l(l+1)P_l(x) = 0,$$

and multiplying with $\log(2/\sqrt{1-x^2})$ and integrating from -1 to 1 , we have

$$\int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) \frac{d}{dx} \left[(1-x^2) \frac{dP_l(x)}{dx} \right] dx + l(l+1) \int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) P_l(x) dx = 0.$$

Now, the first term can be manipulated by partial integration,

$$\begin{aligned} & \int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) \frac{d}{dx} \left[(1-x^2) \frac{dP_l(x)}{dx} \right] dx \\ &= \log \left(\frac{2}{\sqrt{1-x^2}} \right) \cdot (1-x^2) \frac{dP_l(x)}{dx} \Big|_{x=-1}^1 - \int_{-1}^1 \frac{d}{dx} \log \left(\frac{2}{\sqrt{1-x^2}} \right) \cdot (1-x^2) \frac{dP_l(x)}{dx} dx \\ &= - \int_{-1}^1 x \frac{dP_l(x)}{dx} dx \\ &= -xP_l(x) \Big|_{x=-1}^1 + \int_{-1}^1 P_l(x) dx \\ &= -P_l(1) - P_l(-1) = -1 - (-1)^l, \end{aligned}$$

which is nonzero only for even l with the value of -2 . Therefore,

$$\int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) P_l(x) dx = -\frac{1}{l(l+1)} \int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) \frac{d}{dx} \left[(1-x^2) \frac{dP_l(x)}{dx} \right] dx$$

$$= \begin{cases} \frac{1}{j(2j+1)}, & l = 2j \\ 0, & l = 2j+1, \end{cases}$$

and

$$A_l = \frac{2l+1}{2} \int_{-1}^1 \log \left(\frac{2}{\sqrt{1-x^2}} \right) P_l(x) dx = \begin{cases} \frac{4j+1}{2j(2j+1)}, & l = 2j \\ 0, & l = 2j+1, \end{cases}$$

for $j > 0$.

Thus, we have established the identity

$$\log \left(\frac{2}{\sin \theta} \right) = 1 + \sum_{j=1}^{\infty} \frac{4j+1}{2j(2j+1)} P_{2j}(\cos \theta),$$

which gives the desired result for the potential.

(b) From the expansion

$$\frac{1}{|\mathbf{x} - \mathbf{x}'|} = \frac{1}{\sqrt{x^2 + x'^2 - 2xx' \cos \theta}} = \sum_{l=0}^{\infty} \frac{x_{<}^l}{x_{>}^{l+1}} P_l(\cos \theta),$$

by setting $x' = 1$ and taking the limit $x \rightarrow 1 - \varepsilon$ (or, from the other side—this does not change the final result), we have

$$\frac{1}{\sqrt{2(1 - \cos \theta)}} = \frac{1}{2 \sin(\theta/2)} = \sum_{l=0}^{\infty} P_l(\cos \theta).$$

Similarly, perform the same calculation on $1/|\mathbf{x} + \mathbf{x}'|$, we have

$$\frac{1}{\sqrt{2(1 + \cos \theta)}} = \frac{1}{2 \cos(\theta/2)} = \sum_{l=0}^{\infty} (-1)^l P_l(\cos \theta).$$

Take the sum,

$$\frac{1}{2} \left(\frac{1}{\sin(\theta/2)} + \frac{1}{\cos(\theta/2)} \right) = \sum_{l=0}^{\infty} (1 + (-1)^l) P_l(\cos \theta) = 2 \sum_{j=0}^{\infty} P_{2j}(\cos \theta).$$

With the above identity, Eq. (3.137) can be written as

$$\begin{aligned} \sigma(\theta) &= -\frac{Q}{4\pi b^2} \left[1 + \sum_{j=1}^{\infty} \frac{4j+1}{2j+1} P_{2j}(\cos \theta) \right] \\ &= -\frac{Q}{4\pi b^2} \left[1 + 2 \sum_{j=1}^{\infty} P_{2j}(\cos \theta) - \sum_{j=1}^{\infty} \frac{1}{2j+1} P_{2j}(\cos \theta) \right] \\ &= -\frac{Q}{4\pi b^2} \left[2 \sum_{j=0}^{\infty} P_{2j}(\cos \theta) - \sum_{j=0}^{\infty} \frac{1}{2j+1} P_{2j}(\cos \theta) \right] \\ &= -\frac{Q}{4\pi b^2} \left[\frac{1}{2} \left(\frac{1}{\sin(\theta/2)} + \frac{1}{\cos(\theta/2)} \right) - \sum_{j=0}^{\infty} \frac{1}{2j+1} P_{2j}(\cos \theta) \right]. \end{aligned}$$