Algebra Basics

Section 2.2
Graphing Linear Functions

Turning an equation into a picture (graph)

Video Lesson Idea: Start with a story, show the relationship table, have it hanging, and then introduce the graph.

Key Idea: Every solution point from a table, equation, or situation, for all to see on the coordinate plane.

Learning Target

Students will teach how functions are shown in the basics.

- Use a table to find solutions
- Use graphs to find values
- Use graphs to see the shape of the function
- Use graphs to see how a change in one variable affects the other
- Use a story problem to create a graph, plot, or table and see its real-life context

What's To Come

Here's a graph that shows the relationship between the number of hours a worker worked and the amount of money earned. Use the graph to answer the questions below, and also write down the full domain and range. Use all of the symbols.

Video Link

The video below shows a bit of the graph. Can you figure out what it means? Explain.

- Is the graph increasing, decreasing, or constant?
- Is the graph linear or nonlinear?
- Is the graph discrete or continuous?
- Is the graph positive or negative?

Intro

Number

Number is basically the math symbol for 1, representing 1. So when we're talking about the number 1 we're talking about the symbol 1. Number is also a word that means the amount of something. For example, I have a number of pencils.

Every function takes a specific input. The set of inputs that the function accepts is called the domain of the function. The domain is the set of all possible values that can be substituted into the function. For example, the function $f(x) = x^2$ has a domain of all real numbers, because x^2 is defined for all real numbers. The function $f(x) = \frac{1}{x}$ has a domain of all real numbers except 0, because $\frac{1}{0}$ is undefined. The function $f(x) = \sqrt{x}$ has a domain of all positive real numbers, because $\sqrt{x}$ is undefined for negative numbers.

Every function also has a specific output. The set of outputs that the function produces is called the range of the function. For example, the function $f(x) = x^2$ has a range of all positive real numbers, because x^2 is always positive. The function $f(x) = \frac{1}{x}$ has a range of all real numbers except 0, because $\frac{1}{x}$ is never equal to 0. The function $f(x) = \sqrt{x}$ has a range of all positive real numbers, because $\sqrt{x}$ is always positive.

Domain and range can be written in interval notation, which is a way of writing the set of all numbers between two endpoints. For example, the interval $[1, 5]$ means all numbers between 1 and 5, including 1 and 5. The interval $(1, 5)$ means all numbers between 1 and 5, but not including 1 and 5. The interval $[1, 5)$ means all numbers between 1 and 5, including 1 but not including 5. The interval $(1, 5]$ means all numbers between 1 and 5, including 5 but not including 1.

Domain and range can be written in set notation, which is a way of writing the set of all numbers that satisfy a certain condition. For example, the set $\{x \mid x > 0\}$ means all numbers x such that x is greater than 0. The set $\{x \mid x \leq 5\}$ means all numbers x such that x is less than or equal to 5. The set $\{x \mid x \neq 0\}$ means all numbers x such that x is not equal to 0.

Domain and range can be written in words, which is a way of describing the set of all numbers that satisfy a certain condition. For example, the domain of all positive real numbers can be written as "all positive real numbers". The range of all real numbers except 0 can be written as "all real numbers except 0".

Domain and range can be written in graphical form, which is a way of showing the set of all numbers that satisfy a certain condition on a graph. For example, the domain of all positive real numbers can be shown on a graph by shading the region to the right of the y-axis. The range of all real numbers except 0 can be shown on a graph by shading the region above and below the x-axis, but not including the x-axis.

Domain and range can be written in table form, which is a way of showing the set of all numbers that satisfy a certain condition in a table. For example, the domain of all positive real numbers can be shown in a table by listing the numbers 1, 2, 3, 4, 5, and so on. The range of all real numbers except 0 can be shown in a table by listing the numbers 1, -1, 2, -2, 3, -3, and so on.

Domain and range can be written in function notation, which is a way of writing the set of all numbers that satisfy a certain condition for a function. For example, the domain of all positive real numbers can be written as $f(x) = x^2, x > 0$. The range of all real numbers except 0 can be written as $f(x) = \frac{1}{x}, x \neq 0$.

Domain and range can be written in piecewise form, which is a way of writing the set of all numbers that satisfy a certain condition for a piecewise function. For example, the domain of all positive real numbers can be written as $f(x) = \begin{cases} x^2, & x > 0 \\ 0, & x \leq 0 \end{cases}$. The range of all real numbers except 0 can be written as $f(x) = \begin{cases} \frac{1}{x}, & x \neq 0 \\ 1, & x = 0 \end{cases}$.

Domain and range can be written in parametric form, which is a way of writing the set of all numbers that satisfy a certain condition for a parametric function. For example, the domain of all positive real numbers can be written as $x = t^2, y = t, t > 0$. The range of all real numbers except 0 can be written as $x = \frac{1}{t}, y = t, t \neq 0$.

Domain and range can be written in polar form, which is a way of writing the set of all numbers that satisfy a certain condition for a polar function. For example, the domain of all positive real numbers can be written as $r = \theta^2, \theta > 0$. The range of all real numbers except 0 can be written as $r = \frac{1}{\theta}, \theta \neq 0$.

Domain and range can be written in vector form, which is a way of writing the set of all numbers that satisfy a certain condition for a vector function. For example, the domain of all positive real numbers can be written as $\mathbf{r}(t) = t^2\mathbf{i} + t\mathbf{j}, t > 0$. The range of all real numbers except 0 can be written as $\mathbf{r}(t) = \frac{1}{t}\mathbf{i} + t\mathbf{j}, t \neq 0$.

Domain and range can be written in matrix form, which is a way of writing the set of all numbers that satisfy a certain condition for a matrix function. For example, the domain of all positive real numbers can be written as $\begin{bmatrix} t^2 \\ t \end{bmatrix}, t > 0$. The range of all real numbers except 0 can be written as $\begin{bmatrix} \frac{1}{t} \\ t \end{bmatrix}, t \neq 0$.

Domain and range can be written in complex form, which is a way of writing the set of all numbers that satisfy a certain condition for a complex function. For example, the domain of all positive real numbers can be written as $f(z) = z^2, z > 0$. The range of all real numbers except 0 can be written as $f(z) = \frac{1}{z}, z \neq 0$.

Domain and range can be written in quaternion form, which is a way of writing the set of all numbers that satisfy a certain condition for a quaternion function. For example, the domain of all positive real numbers can be written as $q(t) = t^2 + t\mathbf{i} + t\mathbf{j} + t\mathbf{k}, t > 0$. The range of all real numbers except 0 can be written as $q(t) = \frac{1}{t} + t\mathbf{i} + t\mathbf{j} + t\mathbf{k}, t \neq 0$.

Domain and range can be written in octonion form, which is a way of writing the set of all numbers that satisfy a certain condition for an octonion function. For example, the domain of all positive real numbers can be written as $o(t) = t^2 + t\mathbf{i} + t\mathbf{j} + t\mathbf{k} + t\mathbf{l} + t\mathbf{m} + t\mathbf{n} + t\mathbf{o}, t > 0$. The range of all real numbers except 0 can be written as $o(t) = \frac{1}{t} + t\mathbf{i} + t\mathbf{j} + t\mathbf{k} + t\mathbf{l} + t\mathbf{m} + t\mathbf{n} + t\mathbf{o}, t \neq 0$.

Domain and range can be written in sedenion form, which is a way of writing the set of all numbers that satisfy a certain condition for a sedenion function. For example, the domain of all positive real numbers can be written as $s(t) = t^2 + t\mathbf{i} + t\mathbf{j} + t\mathbf{k} + t\mathbf{l} + t\mathbf{m} + t\mathbf{n} + t\mathbf{o} + t\mathbf{p} + t\mathbf{q} + t\mathbf{r} + t\mathbf{s} + t\mathbf{t} + t\mathbf{u} + t\mathbf{v} + t\mathbf{w}, t > 0$. The range of all real numbers except 0 can be written as $s(t) = \frac{1}{t} + t\mathbf{i} + t\mathbf{j} + t\mathbf{k} + t\mathbf{l} + t\mathbf{m} + t\mathbf{n} + t\mathbf{o} + t\mathbf{p} + t\mathbf{q} + t\mathbf{r} + t\mathbf{s} + t\mathbf{t} + t\mathbf{u} + t\mathbf{v} + t\mathbf{w}, t \neq 0$.

Domain and range can be written in trigintanion form, which is a way of writing the set of all numbers that satisfy a certain condition for a trigintanion function. For example, the domain of all positive real numbers can be written as $t(t) = t^2 + t\mathbf{i} + t\mathbf{j} + t\mathbf{k} + t\mathbf{l} + t\mathbf{m} + t\mathbf{n} + t\mathbf{o} + t\mathbf{p} + t\mathbf{q} + t\mathbf{r} + t\mathbf{s} + t\mathbf{t} + t\mathbf{u} + t\mathbf{v} + t\mathbf{w} + t\mathbf{x} + t\mathbf{y} + t\mathbf{z} + t\mathbf{A} + t\mathbf{B} + t\mathbf{C} + t\mathbf{D} + t\mathbf{E} + t\mathbf{F} + t\mathbf{G} + t\mathbf{H} + t\mathbf{I} + t\mathbf{J} + t\mathbf{K} + t\mathbf{L} + t\mathbf{M} + t\mathbf{N} + t\mathbf{O} + t\mathbf{P} + t\mathbf{Q} + t\mathbf{R} + t\mathbf{S} + t\mathbf{T} + t\mathbf{U} + t\mathbf{V} + t\mathbf{W} + t\mathbf{X} + t\mathbf{Y} + t\mathbf{Z}, t > 0$. The range of all real numbers except 0 can be written as $t(t) = \frac{1}{t} + t\mathbf{i} + t\mathbf{j} + t\mathbf{k} + t\mathbf{l} + t\mathbf{m} + t\mathbf{n} + t\mathbf{o} + t\mathbf{p} + t\mathbf{q} + t\mathbf{r} + t\mathbf{s} + t\mathbf{t} + t\mathbf{u} + t\mathbf{v} + t\mathbf{w} + t\mathbf{x} + t\mathbf{y} + t\mathbf{z} + t\mathbf{A} + t\mathbf{B} + t\mathbf{C} + t\mathbf{D} + t\mathbf{E} + t\mathbf{F} + t\mathbf{G} + t\mathbf{H} + t\mathbf{I} + t\mathbf{J} + t\mathbf{K} + t\mathbf{L} + t\mathbf{M} + t\mathbf{N} + t\mathbf{O} + t\mathbf{P} + t\mathbf{Q} + t\mathbf{R} + t\mathbf{S} + t\mathbf{T} + t\mathbf{U} + t\mathbf{V} + t\mathbf{W} + t\mathbf{X} + t\mathbf{Y} + t\mathbf{Z}, t \neq 0$.

Examples

Example 1

Here's a table of values for a function. Plot the points on the graph and connect them.

x	y
0	0
1	1
2	4
3	9
4	16
5	25

Notice how the table and graph tell the same story, they are simply two different ways.

Try! Try it :)

Graph the following: $y = -2x + 1$ and $y = 3$ using the table below and then plot the graph on the same axes.

x	y

Need more practice

- Use the graph and table above to work on the graph!
- Find a function online to make the graph!

Slope and Intercept Can Build the Line

Consider the function $y = 2x + 1$. Using the slope and the y-intercept, we can draw a line. To do this, we first find the slope, then the y-intercept, and then we draw the line.

Practice Now: Find the equation of the line that has a slope of 2 and a y-intercept of 1.

Example 1

Graph $y = -\frac{1}{2}x + 3$ using the slope and the y-intercept. Then find the x-intercept and the domain and range.

Try! Try it :)

Graph $y = \frac{3}{4}x - 2$ using the slope and the y-intercept. Then find the x-intercept and the domain and range.

Graphs Show Direction and Key Features

Graphs are a visual way to show how one thing affects another. For example, how the price of a product affects the number of people who buy it. Or how the number of hours you study affects your test score.

Example 2

Here's a graph of a function. Identify where the increasing and where the decreasing. The maximum, the minimum, and the zeros.

Try! Try it :)

Here's a graph of a function. Identify where the increasing and where the decreasing. The maximum, the minimum, and the zeros.

Need more practice

- Use the graph and table above to work on the graph!
- Find a function online to make the graph!

A Graph Should Match the Situation

When a story describes a function, the graph should match the story. Each part of the graph should tell some story. A graph can be thought of as a picture of a story. Each point on the graph tells a piece of the story.

Example 3

Here's a graph that shows the relationship between the number of hours a worker worked and the amount of money earned. Use the graph to answer the questions below, and also write down the full domain and range. Use all of the symbols.

Try! Try it :)

Here's a graph that shows the relationship between the number of hours a worker worked and the amount of money earned. Use the graph to answer the questions below, and also write down the full domain and range. Use all of the symbols.

Need more practice

- Use the graph and table above to work on the graph!
- Find a function online to make the graph!

Summary

- A linear function graphs as a straight line.
- Slope is the rate of change, and it tells you how steep a line is. A positive slope means the line goes up from left to right, a negative slope means the line goes down from left to right, a slope of zero means the line is horizontal, and an undefined slope means the line is vertical.
- A graph's slope tells you how fast the function is increasing or decreasing.
- A graph's y-intercept tells you where the line crosses the y-axis.
- A graph's x-intercept tells you where the line crosses the x-axis.

Common Mistakes

- **Slope and y-intercept** mixed up. Remember: $y = mx + b$, m is slope, b is y-intercept.
- **Using rise/run backwards** Rise is the change in y, run is the change in x. Always $\frac{\text{rise}}{\text{run}}$.
- **Forgetting the negative** when a line goes down from left to right.
- **Starting at the wrong point** when counting rise and run. Always start at the point you're given.
- **Drawing a line** without using at least two points or the slope and y-intercept.

Additional Practice Tools Reflection

Student Reflection: Which practice tool helped you the most and why?

- Graphing a function on desmos to see how the shape of the graph changes.
- Using the graph that's already there.
- Using the animated graph of the function to see what's going on.