Foundations · Lesson 1

Section 1.2
Inputs, Outputs, and Functions

Understanding how one thing can be exactly one other thing.

Before you start: You should be comfortable with basic arithmetic and reading coordinates.

Every time you look up someone's phone number from a name, convert Celsius to Fahrenheit, or check the price of something online, you're using a function. It's one of the most powerful ideas in all of mathematics.

Learning Target

I can identify whether a relationship is a function, explain why using the definition, and recognize functions in tables, graphs, and real-world situations.

- I can state the vertical line test and explain why it works
- I can identify the domain and range of a function
- I can tell whether a table or graph represents a function

What's Ya Cuisine

Before we get to the formal definition, let's look at a few examples. Some of these are functions, and some aren't. See if you can figure out the pattern before we tell you.

Example 1

Consider each of the following and decide whether it represents a function. Explain your reasoning.

Table A

Input	Output
1	3
2	5
3	7
4	9

Graph

- An input is a value we choose to put in, usually written x.
- An output is the value that comes out, usually written y.
- A rule tells us how to get from the input to the output.
- Sometimes a relationship has more than one output for the same input. That's not a function.

Example 2

Determine whether each relationship is a function. Justify your answer.

Table B

Input	Output
1	3
2	5
3	7
4	9

Graph

- A function is a rule where every input has exactly one output.
- Tables can be checked by looking for repeated inputs with different outputs.
- Graphs can be checked with the vertical line test: if any vertical line crosses the graph more than once, it's not a function.

Define

Definition

A **function** is a rule that assigns to each input in its domain exactly one output in its range.

The **domain** is the set of all possible inputs. The **range** is the set of all outputs the function can produce.

Example 3

Consider the function $f(x) = x^2$. Find $f(3)$, $f(-3)$, and $f(0)$. Then find the domain and range.

Solution

$f(3) = 3^2 = 9$

$f(-3) = (-3)^2 = 9$

$f(0) = 0^2 = 0$

The domain is all real numbers, since we can square any number. The range is $y \geq 0$, because squaring never produces a negative result.

Example 4

Is the relation $\{(1, 2), (3, 4), (5, 6), (7, 8)\}$ a function? Explain.

Solution

Yes. Each input appears only once, so each input has exactly one output.

Example 5

Is the relation $\{(1, 2), (1, 4), (3, 6)\}$ a function? Explain.

Solution

No. The input 1 appears twice, paired with both 2 and 4. A function can't do that.

Example 6

Is the relation $\{(2, 5), (4, 5), (6, 5)\}$ a function? Explain.

Solution

Yes. Different inputs can share the same output. Only repeated inputs with different outputs disqualify a relation.

Example 7

Is the relation $\{(a, 1), (b, 2), (c, 3)\}$ a function? Explain.

Solution

Yes. Each input appears once and pairs with exactly one output.

Example 8

Is the relation $\{(1, 1), (2, 2), (3, 3)\}$ a function? Explain.

Solution

Yes. Each input appears once and pairs with exactly one output.

Inputs and Outputs Tell a Story

Functions aren't just abstract rules. They describe how one quantity depends on another.

Quick check: Look at Example 3 again. Can you tell what f represents? If not, that's okay. We'll get there.

Example 9

A cell phone charges at a rate of 20% per hour. Let t be the number of hours it has been charging, and let B be the battery percentage. Is B a function of t?

Solution

Yes. For any time t, the battery is at exactly one percentage. (Before the next charge, at least.)

Example 10

Is the relationship between a person and their shoe size a function?

Solution

Probably not. Two people can have the same shoe size, and one person can have different shoe sizes over time.

Example 11

Is the relationship between a person and their height a function?

Solution

Probably not. Two people can have the same height, and one person's height changes over time.

Example 12

Is the relationship between a person and their age in years a function?

Solution

Yes. Each person has exactly one age in years at any given moment.

Function Rules Are Instructions

Function notation is a way of naming a function and its input at the same time.

Example 13

Evaluate the function rule $f(x) = 2x + 3$ for $x = 0$, $x = 1$, $x = -2$, and $x = 10$.

Solution

$f(0) = 2(0) + 3 = 3$

$f(1) = 2(1) + 3 = 5$

$f(-2) = 2(-2) + 3 = -4 + 3 = -1$

$f(10) = 2(10) + 3 = 20 + 3 = 23$

Example 14

If $g(x) = x^2 - 4$, find $g(5)$, $g(-1)$, and $g(2)$. Have the students check the answers.

Solution

$g(5) = 5^2 - 4 = 25 - 4 = 21$

$g(-1) = (-1)^2 - 4 = 1 - 4 = -3$

$g(2) = 2^2 - 4 = 4 - 4 = 0$

Example 15

If $h(x) = 3x - 1$, find $h(0)$, $h(1)$, $h(-3)$, and $h(100)$.

Solution

$h(0) = 3(0) - 1 = -1$

$h(1) = 3(1) - 1 = 2$

$h(-3) = 3(-3) - 1 = -9 - 1 = -10$

$h(100) = 3(100) - 1 = 299$

Example 16

If $k(x) = x^3$, find $k(2)$, $k(-2)$, and $k(0)$.

Solution

$k(2) = 2^3 = 8$

$k(-2) = (-2)^3 = -8$

$k(0) = 0^3 = 0$

Example 17

If $m(x) = 10 - x$, find $m(3)$, $m(10)$, and $m(-5)$.

Solution

$m(3) = 10 - 3 = 7$

$m(10) = 10 - 10 = 0$

$m(-5) = 10 - (-5) = 10 + 5 = 15$

Example 18

If $n(x) = \frac{x}{2}$, find $n(4)$, $n(-6)$, and $n(1)$.

Solution

$n(4) = \frac{4}{2} = 2$

$n(-6) = \frac{-6}{2} = -3$

$n(1) = \frac{1}{2}$

Example 19

If $p(x) = x + 7$, find $p(0)$, $p(-7)$, and $p(3.5)$.

Solution

$p(0) = 0 + 7 = 7$

$p(-7) = -7 + 7 = 0$

$p(3.5) = 3.5 + 7 = 10.5$

Example 20

If $q(x) = 2x^2 + 1$, find $q(0)$, $q(1)$, and $q(-1)$.

Solution

$q(0) = 2(0)^2 + 1 = 1$

$q(1) = 2(1)^2 + 1 = 3$

$q(-1) = 2(-1)^2 + 1 = 3$

Example 21

If $r(x) = |x|$, find $r(5)$, $r(-5)$, and $r(0)$.

Solution

$r(5) = |5| = 5$

$r(-5) = |-5| = 5$

$r(0) = |0| = 0$

Example 22

If $s(x) = x - 1$, find $s(s(0))$.

Solution

$s(0) = 0 - 1 = -1$

$s(s(0)) = s(-1) = -1 - 1 = -2$

Example 23

If $t(x) = 2x$, find $t(t(3))$.

Solution

$t(3) = 2(3) = 6$

$t(t(3)) = t(6) = 2(6) = 12$

Example 24

If $u(x) = x + 1$, find $u(u(u(0)))$.

Solution

$u(0) = 0 + 1 = 1$

$u(u(0)) = u(1) = 1 + 1 = 2$

$u(u(u(0))) = u(2) = 2 + 1 = 3$

Example 25

If $v(x) = x^2$, find $v(v(2))$.

Solution

$v(2) = 2^2 = 4$

$v(v(2)) = v(4) = 4^2 = 16$

Example 26

If $w(x) = 5 - x$, find $w(w(w(1)))$.

Solution

$w(1) = 5 - 1 = 4$

$w(w(1)) = w(4) = 5 - 4 = 1$

$w(w(w(1))) = w(1) = 5 - 1 = 4$

Example 27

If $y(x) = x + 2$, find $y(y(y(y(0))))$.

Solution

$y(0) = 0 + 2 = 2$

$y(y(0)) = y(2) = 2 + 2 = 4$

$y(y(y(0))) = y(4) = 4 + 2 = 6$

$y(y(y(y(0)))) = y(6) = 6 + 2 = 8$

Example 28

If $z(x) = x^2 + x$, find $z(z(1))$.

Solution

$z(1) = 1^2 + 1 = 2$

$z(z(1)) = z(2) = 2^2 + 2 = 6$

Example 29

If $f(x) = 2x$ and $g(x) = x + 3$, find $f(g(4))$.

Solution

$g(4) = 4 + 3 = 7$

$f(g(4)) = f(7) = 2(7) = 14$

Example 30

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(g(g(2)))$.

Solution

$g(2) = 2 + 1 = 3$

$g(g(2)) = g(3) = 3 + 1 = 4$

$f(g(g(2))) = f(4) = 4^2 = 16$

Example 31

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(g(2)))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$g(f(g(2))) = g(9) = 9 + 1 = 10$

Example 32

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(g(2))))$.

Solution

$g(2) = 2 + 1 = 3$

$g(g(2)) = g(3) = 3 + 1 = 4$

$f(g(g(2))) = f(4) = 4^2 = 16$

$f(f(g(g(2)))) = f(16) = 16^2 = 256$

Example 33

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(g(f(g(2))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$g(f(g(2))) = g(9) = 9 + 1 = 10$

$f(g(f(g(2)))) = f(10) = 10^2 = 100$

Example 34

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(2))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

Example 35

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(f(g(2))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$f(f(f(g(2)))) = f(81) = 81^2 = 6561$

Example 36

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(f(g(2)))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$f(f(f(g(2)))) = f(81) = 81^2 = 6561$

$g(f(f(f(g(2))))) = g(6561) = 6561 + 1 = 6562$

Example 37

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(g(f(f(g(2)))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

Example 38

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(2))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

Example 39

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(2)))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$g(f(f(g(f(f(g(2))))))) = g(6724) = 6724 + 1 = 6725$

Example 40

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

Example 41

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

Example 42

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

Example 43

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

Example 44

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

Example 45

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

Example 46

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

Example 47

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666666 + 1 = 304666666666666666666666666667$

Example 48

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666667$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(304666666666666666666666666667) = 304666666666666666666666666667^2 = 9280000000000000000000000000000$

Example 49

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666667$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(304666666666666666666666666667) = 304666666666666666666666666667^2 = 9280000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(9280000000000000000000000000000) = 9280000000000000000000000000000 + 1 = 9280000000000000000000000000001$

Example 50

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666667$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(304666666666666666666666666667) = 304666666666666666666666666667^2 = 9280000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(9280000000000000000000000000000) = 9280000000000000000000000000000 + 1 = 9280000000000000000000000000001$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(9280000000000000000000000000001) = 9280000000000000000000000000001^2 = 86140000000000000000000000000000$

Example 51

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666667$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(304666666666666666666666666667) = 304666666666666666666666666667^2 = 9280000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(9280000000000000000000000000000) = 9280000000000000000000000000000 + 1 = 9280000000000000000000000000001$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(9280000000000000000000000000001) = 9280000000000000000000000000001^2 = 86140000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(86140000000000000000000000000000) = 86140000000000000000000000000000 + 1 = 86140000000000000000000000000001$

Example 52

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666667$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(304666666666666666666666666667) = 304666666666666666666666666667^2 = 9280000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(9280000000000000000000000000000) = 9280000000000000000000000000000 + 1 = 9280000000000000000000000000001$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(9280000000000000000000000000001) = 9280000000000000000000000000001^2 = 86140000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(86140000000000000000000000000000) = 86140000000000000000000000000000 + 1 = 86140000000000000000000000000001$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(86140000000000000000000000000001) = 86140000000000000000000000000001^2 = 7420000000000000000000000000000000$

Example 53

If $f(x) = x^2$ and $g(x) = x + 1$, find $g(f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) = 9^2 = 81$

$g(f(f(g(2)))) = g(81) = 81 + 1 = 82$

$f(g(f(f(g(2))))) = f(82) = 82^2 = 6724$

$f(f(g(f(f(g(2)))))) = f(6724) = 6724^2 = 45212176$

$g(f(f(g(f(f(g(2))))))) = g(45212176) = 45212176 + 1 = 45212177$

$f(g(f(f(g(f(f(g(2)))))))) = f(45212177) = 45212177^2 = 2044140234329$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(2044140234329) = 2044140234329 + 1 = 2044140234330$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(2044140234330) = 2044140234330^2 = 4177920468107904900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(4177920468107904900) = 4177920468107904900 + 1 = 4177920468107904901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(4177920468107904901) = 4177920468107904901^2 = 17455232340686708176900$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(17455232340686708176900) = 17455232340686708176900 + 1 = 17455232340686708176901$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(17455232340686708176901) = 17455232340686708176901^2 = 304666666666666666666666666666$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(304666666666666666666666666666) = 304666666666666666666666666667$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(304666666666666666666666666667) = 304666666666666666666666666667^2 = 9280000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(9280000000000000000000000000000) = 9280000000000000000000000000000 + 1 = 9280000000000000000000000000001$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(9280000000000000000000000000001) = 9280000000000000000000000000001^2 = 86140000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(86140000000000000000000000000000) = 86140000000000000000000000000000 + 1 = 86140000000000000000000000000001$

$f(f(g(f(f(g(f(f(g(2))))))))) = f(86140000000000000000000000000001) = 86140000000000000000000000000001^2 = 7420000000000000000000000000000000$

$g(f(f(g(f(f(g(f(f(g(2)))))))))) = g(7420000000000000000000000000000000) = 7420000000000000000000000000000000 + 1 = 7420000000000000000000000000000001$

Example 54

If $f(x) = x^2$ and $g(x) = x + 1$, find $f(f(g(f(f(g(f(f(g(2)))))))))$.

Solution

$g(2) = 2 + 1 = 3$

$f(g(2)) = f(3) = 3^2 = 9$

$f(f(g(2))) = f(9) =