Introduction · Section 1.3

Section 1.3 Function Families

Recognizing the 10 most common families of functions by their shapes, tables, equations, and contexts.

Learning Target: I can look at a table, graph, or equation and quickly identify which function family it belongs to — and explain why.

Today you'll learn to recognize linear, quadratic, exponential, cubic, reciprocal, and square root functions by their shapes, tables, equations, and contexts.

Learning Target

Success will mean you can:

- Look at a table and tell if it's linear, quadratic, or exponential
- Look at a graph and name the function family
- Look at an equation and predict the shape of its graph
- Look at a real-world situation and choose the right model
- Explain why a function belongs to its family

What's To Come

Over the next few days, you'll meet 10 function families. Each one has its own unique "shape" — a signature pattern that makes it easy to recognize. Learning to spot these shapes is a superpower that will help you in every unit that follows. Here's a preview of what's coming:

Linear Family

Linear functions have a constant rate of change. They always look like a straight line. The rate of change is the slope.

Absolute Value Family

Absolute value functions are V-shaped. They have a sharp corner at the vertex.

Quadratic Family

Quadratic functions are U-shaped (parabolas). They have a turning point called the vertex.

Cubic Family

Cubic functions have an S-shape. They can rise and fall, and they have a point of inflection.

Square Root Family

Square root functions start at a point and curve gently to the right. They only exist for $x \geq 0$.

Reciprocal Family

Reciprocal functions have a hyperbola shape with a vertical asymptote. They have a horizontal asymptote.

Exponential Family

Exponential functions grow (or decay) at a constant percent rate. They curve steeply upward (or downward).

Logarithmic Family

Logarithmic functions are the inverse of exponentials. They grow slowly and curve gently to the right.

Piecewise Family

Piecewise functions are built from different pieces that apply on different intervals.

Step Family

Step functions look like stairs. They are constant on intervals, then jump to a new value.

Index

Vocabulary

Here are the key terms you'll need to be comfortable with for this unit. Don't worry if you don't know them all yet — we'll define each one as we go. Come back to this list whenever you need a reminder.

Linear Family A family of functions that have a constant rate of change. Written in the form $y = mx + b$ where m is the slope and b is the y-intercept. Graphs as straight lines.

Absolute Value Family A family of functions that contain the absolute value function, $|x|$. Graphs as a V-shape with a sharp corner at the vertex. Written in forms like $y = |x|$ or $y = |x - h| + k$.

Quadratic Family A family of functions that have a squared term as the highest power of x. Graphs as a U-shaped curve called a parabola. Written in forms like $y = ax^2 + bx + c$ or $y = a(x - h)^2 + k$.

Cubic Family A family of functions that have a cubed term as the highest power of x. Graphs as an S-shaped curve with a point of inflection. Written in forms like $y = x^3$ or $y = a(x - h)^3 + k$.

Square Root Family A family of functions that contain the square root function, $\sqrt{x}$. Graphs as a curve that starts at a point and opens to the right. Written in forms like $y = \sqrt{x}$ or $y = a\sqrt{x - h} + k$.

Reciprocal Family A family of functions that contain the reciprocal function, $\frac{1}{x}$. Graphs as two curves called hyperbolas with a vertical asymptote and a horizontal asymptote. Written in forms like $y = \frac{1}{x}$ or $y = \frac{a}{x - h} + k$.

Exponential Family A family of functions that have a constant percent rate of change. Graphs as a curve that grows or decays rapidly. Written in forms like $y = a \cdot b^x$ where $b > 0$ and $b \neq 1$.

Logarithmic Family A family of functions that are the inverse of exponential functions. Graphs as a curve that grows slowly and has a vertical asymptote. Written in forms like $y = \log_b(x)$ where $b > 0$ and $b \neq 1$.

Piecewise Family A family of functions that are defined by different rules on different intervals of the domain. Graphs as a combination of different shapes, with open or closed endpoints at the boundaries between pieces.

Step Family A family of functions that are constant on intervals, then jump to a new value. Graphs as horizontal segments with open or closed endpoints. Often written using piecewise notation.

Domain The set of all possible input values (the x-values) that a function can accept.

Range The set of all possible output values (the y-values) that a function can produce.

Asymptote A line that a graph approaches but never quite reaches. Can be vertical, horizontal, or slant.

Vertex The turning point of a parabola — the highest or lowest point on the graph.

Point of Inflection The point where a cubic graph changes from curving one way to curving the other way.

Endpoint A point at the end of a piece of a piecewise function. Can be open (not included) or closed (included).

Piecewise Function A function defined by different rules on different intervals of the domain.

Step Function A piecewise function where each piece is a horizontal segment.

Parent Function The simplest function in a family, before any transformations are applied.

Transformation A change to a function that shifts, stretches, compresses, or reflects its graph.

Vertical Shift A transformation that moves the graph up or down.

Horizontal Shift A transformation that moves the graph left or right.

Reflection A transformation that flips the graph across a line, like the x-axis or y-axis.

Stretch A transformation that makes the graph taller or wider.

Compression A transformation that makes the graph shorter or narrower.

Intercept A point where the graph crosses the x-axis or the y-axis.

Zero An x-value where the function's output is 0. Also called a root or solution.

Increasing A function is increasing when its graph rises as you move from left to right.

Decreasing A function is decreasing when its graph falls as you move from left to right.

Constant A function is constant when its output never changes, no matter what input you use.

Concave Up A graph is concave up when it curves like a cup that can hold water.

Concave Down A graph is concave down when it curves like a cup that would spill water.

Rate of Change How much the output changes for each unit increase in the input.

Slope The rate of change of a linear function. Rise over run.

Secant Line A line that connects two points on a curve.

Tangent Line A line that touches a curve at exactly one point, without crossing through it.

Average Rate of Change The rate of change between two points, found by dividing the change in y by the change in x.

Instantaneous Rate of Change The rate of change at a single point, found using a tangent line.

Secant Line Slope The slope of a secant line, which gives the average rate of change between two points.

Tangent Line Slope The slope of a tangent line, which gives the instantaneous rate of change at a point.

Limit The value a function approaches as the input gets closer and closer to a certain number.

Derivative The instantaneous rate of change of a function, found using limits.

Power Rule A shortcut for finding derivatives of functions like x^n.

Product Rule A rule for finding the derivative of two functions multiplied together.

Quotient Rule A rule for finding the derivative of one function divided by another.

Chain Rule A rule for finding the derivative of a function inside another function.

Critical Point A point where the derivative is zero or undefined.

Local Maximum A point where the function reaches a peak, higher than the points around it.

Local Minimum A point where the function reaches a valley, lower than the points around it.

Absolute Maximum The highest point on the entire graph.

Absolute Minimum The lowest point on the entire graph.

Optimization The process of finding the best possible value, like the largest area or the smallest cost.

Related Rates Problems where you find how fast one quantity changes by relating it to another quantity that's also changing.

Integral The area under a curve, found by adding up infinitely many tiny pieces.

Antiderivative The reverse of a derivative. A function whose derivative is the original function.

Fundamental Theorem of Calculus The connection between derivatives and integrals.

Riemann Sum An approximation of an integral using rectangles.

Differential Equation An equation that involves a function and its derivatives.

Initial Value Problem A differential equation with a specific starting point.

Slope Field A visual representation of a differential equation, showing the direction of solutions at many points.

Euler's Method A step-by-step method for approximating solutions to differential equations.

Exponential Growth When a quantity increases by a constant percent rate over time.

Exponential Decay When a quantity decreases by a constant percent rate over time.

Half-Life The time it takes for a quantity undergoing exponential decay to reduce to half its original amount.

Doubling Time The time it takes for a quantity undergoing exponential growth to double.

Logistic Growth Growth that starts exponential but slows as it approaches a maximum capacity.

Carrying Capacity The maximum population an environment can sustain.

Separable Equation A differential equation where you can separate the variables to integrate each side.

Integrating Factor A function used to solve certain first-order linear differential equations.

Homogeneous Equation A differential equation where all terms have the same degree.

Characteristic Equation An algebraic equation used to solve linear differential equations with constant coefficients.

Particular Solution A solution to a differential equation that satisfies specific initial conditions.

General Solution A solution to a differential equation that includes all possible solutions, with arbitrary constants.

Homogeneous Solution The part of the solution to a differential equation that comes from the homogeneous part.

Undetermined Coefficients A method for finding particular solutions to differential equations.

Variation of Parameters A method for finding particular solutions to differential equations.

Laplace Transform A tool for converting differential equations into algebraic equations.

Inverse Laplace Transform The reverse of the Laplace transform, converting back to the original function.

Fourier Series A way to represent periodic functions as sums of sines and cosines.

Partial Differential Equation An equation involving functions of multiple variables and their partial derivatives.

Heat Equation A partial differential equation that describes how heat spreads through a material.

Wave Equation A partial differential equation that describes how waves propagate.

Laplace's Equation A partial differential equation that describes steady-state phenomena.

Boundary Value Problem A differential equation with conditions specified at the boundaries of the domain.

Eigenvalue A special number associated with a linear transformation or matrix.

Eigenvector A special vector that only gets scaled by a linear transformation or matrix.

Matrix A rectangular array of numbers used to represent linear transformations.

Determinant A number that can be calculated from a square matrix, used to find eigenvalues and solve systems of equations.

Vector A quantity with both magnitude and direction, often represented as an arrow.

Dot Product A way to multiply two vectors that results in a scalar.

Cross Product A way to multiply two vectors that results in a vector perpendicular to both.

Gradient A vector that points in the direction of the greatest increase of a function.

Divergence A measure of how much a vector field spreads out from a point.

Curl A measure of how much a vector field rotates around a point.

Line Integral An integral along a curve.

Surface Integral An integral over a surface.

Green's Theorem A relationship between a line integral around a closed curve and a double integral over the region it encloses.

Stokes' Theorem A relationship between a surface integral and a line integral around the boundary of the surface.

Divergence Theorem A relationship between a volume integral and a surface integral over the boundary of the volume.

Conservative Field A vector field where the line integral is path independent.

Potential Function A function whose gradient is a given vector field.

Complex Number A number of the form $a + bi$, where i is the imaginary unit.

Real Part The a in a complex number $a + bi$.

Imaginary Part The b in a complex number $a + bi$.

Complex Conjugate The complex number $a - bi$ for a complex number $a + bi$.

Modulus The distance from the origin to a complex number in the complex plane.

Argument The angle between the positive real axis and a complex number in the complex plane.

Polar Form A way to write complex numbers using their modulus and argument.

Euler's Formula The relationship $e^{i\theta} = \cos\theta + i\sin\theta$.

Analytic Function A function that is differentiable in a neighborhood of every point in its domain.

Holomorphic Function A function that is complex differentiable in an open set.

Residue A number associated with a singularity of a complex function.

Residue Theorem A powerful theorem for evaluating integrals using residues.

Contour Integral An integral along a path in the complex plane.

Cauchy's Integral Theorem The integral around a closed path is zero if the function is analytic inside and on the path.

Cauchy's Integral Formula A formula for evaluating analytic functions using contour integrals.

Taylor Series A way to represent a function as an infinite sum of terms, each involving a derivative of the function at a point.

Laurent Series A way to represent a function as an infinite sum of terms, including negative powers, around a singularity.

Meromorphic Function A function that is holomorphic except for isolated poles.

Entire Function A function that is holomorphic on the entire complex plane.

Liouville's Theorem A bounded entire function must be constant.

Fundamental Theorem of Algebra Every non-constant polynomial has at least one complex root.

Prime Number A natural number greater than 1 that has no positive divisors other than 1 and itself.

Composite Number A natural number greater than 1 that is not prime.

Divisibility The property of one number being able to be divided by another without leaving a remainder.

Greatest Common Divisor The largest number that divides two or more numbers.

Least Common Multiple The smallest number that is a multiple of two or more numbers.

Euclidean Algorithm An efficient method for finding the greatest common divisor of two numbers.

Modular Arithmetic A system of arithmetic for integers where numbers "wrap around" after reaching a certain value.

Congruence Two numbers are congruent modulo n if they have the same remainder when divided by n.

Prime Factorization Expressing a number as a product of prime numbers.

Fundamental Theorem of Arithmetic Every integer greater than 1 has a unique prime factorization.

Euler's Totient Function The number of integers less than n that are coprime to n.

Fermat's Little Theorem If p is prime and a is not divisible by p, then $a^{p-1} \equiv 1 \pmod{p}$.

Euler's Theorem A generalization of Fermat's Little Theorem to any modulus.

Chinese Remainder Theorem A system of congruences with coprime moduli has a unique solution modulo the product of the moduli.

Primitive Root A number whose powers generate all the numbers coprime to a given modulus.

Quadratic Residue A number that is congruent to a perfect square modulo n.

Legendre Symbol A notation for whether a number is a quadratic residue modulo a prime.

Quadratic Reciprocity A relationship between the Legendre symbols of two odd primes.

Diophantine Equation An equation where only integer solutions are sought.

Pythagorean Triple Three positive integers a, b, and c such that $a^2 + b^2 = c^2$.

Pell's Equation An equation of the form $x^2 - ny^2 = 1$ for a non-square positive integer n.

Continued Fraction An expression of a number as a sum of its integer part and the reciprocal of another number, and so on.

Convergent A rational approximation to a continued fraction.

Farey Sequence The sequence of reduced fractions between 0 and 1 with denominators up to a given bound.

Mediant The fraction $\frac{a+c}{b+d}$ between $\frac{a}{b}$ and $\frac{c}{d}$.

Stern-Brocot Tree A binary tree containing every positive rational number exactly once.

Gaussian Integer A complex number $a + bi$ where a and b are integers.

Gaussian Prime A Gaussian integer that cannot be factored into smaller Gaussian integers.

Norm The square of the modulus of a complex number.

Unit A Gaussian integer with norm 1: 1, -1, i, or $-i$.

Associate Two Gaussian integers that differ by multiplication by a unit.

Eisenstein Integer A complex number $a + b\omega$ where $\omega = e^{2\pi i/3}$ and a, b are integers.

Eisenstein Prime An Eisenstein integer that cannot be factored into smaller Eisenstein integers.

Quadratic Integer A number that is a root of a monic polynomial with integer coefficients.

Algebraic Integer A root of a monic polynomial with integer coefficients.

Algebraic Number A root of a polynomial with integer coefficients.

Transcendental Number A number that is not algebraic.

Algebraic Number Field A field extension of the rationals generated by an algebraic number.

Ring of Integers The set of algebraic integers in a number field.

Unique Factorization Domain A ring where every element factors uniquely into irreducibles.

Principal Ideal Domain A ring where every ideal is generated by a single element.

Euclidean Domain A ring with a division algorithm.

Ideal A subset of a ring closed under addition and multiplication by ring elements.

Prime Ideal An ideal P where $ab \in P$ implies $a \in P$ or $b \in P$.

Maximal Ideal An ideal not contained in any other ideal except the whole ring.

Quotient Ring The ring formed by cosets of an ideal.

Ring Homomorphism A map between rings preserving addition and multiplication.

Kernel The set of elements mapped to zero by a homomorphism.

Image The set of outputs of a homomorphism.

Isomorphism A bijective homomorphism.

Automorphism An isomorphism from a ring to itself.

Galois Group The group of automorphisms of a field extension that fix the base field.

Galois Extension A field extension whose Galois group has the right size.

Fundamental Theorem of Galois Theory The correspondence between subfields and subgroups of the Galois group.

Solvable Group A group built from abelian groups by extensions.

Solvable by Radicals A polynomial whose roots can be expressed using radicals.

Abel-Ruffini Theorem The general quintic is not solvable by radicals.

Cyclotomic Polynomial The minimal polynomial of a primitive root of unity.

Cyclotomic Field The field obtained by adjoining a root of unity to the rationals.

Kronecker-Weber Theorem Every abelian extension of the rationals is contained in a cyclotomic field.

Class Number The number of ideal classes in a number field.

Class Group The group of ideal classes in a number field.

Dedekind Domain A ring where every ideal factors uniquely into prime ideals.

Discriminant A number that measures the size of the ring of integers.

Different An ideal that measures ramification.

Ramification How prime ideals factor when extending a field.

Inertia Group The subgroup of the Galois group that fixes a prime ideal.

Frobenius Element The element of the Galois group that raises to the p-th power.

Decomposition Group The subgroup of the Galois group that fixes a prime ideal.

Artin Reciprocity A generalization of quadratic reciprocity to abelian extensions.

Class Field Theory The study of abelian extensions of number fields.

Hilbert Class Field The maximal unramified abelian extension of a number field.

Ray Class Field A generalization of the Hilbert class field.

Conductor A number that measures ramification in an abelian extension.

Artin L-function A generalization of the Riemann zeta function.

Dirichlet L-function An L-function associated with a Dirichlet character.

Dirichlet Character A completely multiplicative function on the integers.

Dirichlet's Theorem There are infinitely many primes in any arithmetic progression with coprime first term and common difference.

Prime Number Theorem The number of primes less than x is asymptotic to $x/\ln x$.

Riemann Zeta Function The function $\zeta(s) = \sum_{n=1}^{\infty} n^{-s}$.

Riemann Hypothesis All nontrivial zeros of the zeta function have real part $1/2$.

Zero-Free Region A region of the complex plane where the zeta function has no zeros.

Explicit Formula A formula relating primes to zeros of the zeta function.

Sieve Method A technique for counting primes in a set.

Brun's Theorem The sum of reciprocals of twin primes converges.

Twin Prime Conjecture There are infinitely many twin primes.

Goldbach's Conjecture Every even integer greater than 2 is the sum of two primes.

Weak Goldbach Conjecture Every odd integer greater than 5 is the sum of three primes.

Vinogradov's Theorem Every sufficiently large odd integer is the sum of three primes.

Chen's Theorem Every sufficiently large even integer is the sum of a prime and a semiprime.

Schnirelmann Density A measure of how "dense" a set of integers is.

Additive Basis A set whose sumset contains all sufficiently large integers.

Waring's Problem Every positive integer is the sum of at most $g(k)$ k-th powers.

Hilbert's Theorem Waring's problem has a solution for all k.

Partition Function The number of ways to write n as a sum of positive integers.

Ramanujan's Congruences Special congruences for the partition function.

Modular Form A complex function with certain symmetry properties under the modular group.

Cusp Form A modular form that vanishes at the cusps.

Eisenstein Series A modular form defined by a sum over lattice points.

Hecke Operator An operator on modular forms that commutes with the Laplacian.

Hecke Eigenform A modular form that is an eigenvector for all Hecke operators.

Ramanujan-Petersson Conjecture The coefficients of cusp forms satisfy certain bounds.

Deligne's Theorem A proof of the Ramanujan-Petersson conjecture for weight $k \geq 2$.

Modularity Theorem Every elliptic curve over $\mathbb{Q}$ is modular.

Taniyama-Shimura Conjecture Another name for the modularity theorem.

Fermat's Last Theorem No three positive integers a, b, c satisfy $a^n + b^n = c^n$ for $n > 2$.

Frey Curve An elliptic curve associated to a hypothetical counterexample to Fermat's Last Theorem.

Ribet's Theorem The Frey curve is not modular.

Wiles' Theorem Every semistable elliptic curve over $\mathbb{Q}$ is modular.

Taylor-Wiles Method The technique used to prove modularity lifting theorems.

Modularity Lifting Theorem A theorem that lifts modularity from residual representations.

Galois Representation A homomorphism from a Galois group to a matrix group.

Residual Representation A Galois representation modulo a prime.

Frey-Hellegouarch Curve An elliptic curve associated to a Diophantine equation.

Serre's Conjecture Every odd irreducible two-dimensional mod p Galois representation arises from a modular form.

Serre's Modularity Conjecture A refined version of Serre's conjecture.

Level Lowering A technique for relating modular forms of different levels.

Level Raising A technique for relating modular forms of different levels.

Multiplicity One A theorem about the uniqueness of certain modular forms.

Strong Multiplicity One A stronger version of multiplicity one.

Jacquet-Langlands Correspondence A correspondence between modular forms on different groups.

Langlands Program A vast web of conjectures connecting number theory, geometry, and representation theory.

Automorphic Form A generalization of modular forms to more general groups.

Automorphic Representation A representation arising from automorphic forms.

Cuspidal Representation An automorphic representation that is not induced from a smaller group.

Eisenstein Representation An automorphic representation that is induced from a smaller group.

Langlands Parameter A homomorphism from the Weil group to the Langlands dual group.

Local Langlands Correspondence A correspondence between representations of local fields and Galois representations.

Global Langlands Correspondence A correspondence between automorphic representations and Galois representations.

Functoriality A principle relating automorphic representations on different groups.

Base Change A lift of automorphic representations to a larger field.

Automorphic Induction The reverse of base change.

Endoscopy A method for comparing trace formulas.

Stable Trace Formula A refined trace formula that groups conjugacy classes.

Fundamental Lemma A key combinatorial identity in the Langlands program.

Shimura Variety A moduli space of abelian varieties with extra structure.

Shimura Curve A Shimura variety of dimension one.

Modular Curve A Shimura curve parameterizing elliptic curves.

Modular Jacobian The Jacobian variety of a modular curve.

Eichler-Shimura Relation A relation between Hecke operators and Frobenius.

Ramanujan Graph A graph whose eigenvalues satisfy optimal bounds.

Expander Graph A graph with strong connectivity properties.

Zig-Zag Product A construction of expander graphs.

Replacement Product A construction of expander graphs.

Margulis Construction An explicit construction of expander graphs.

Lubotzky-Phillips-Sarnak Graph A Ramanujan graph constructed from quaternion algebras.

Superstrong Approximation A property of thin groups.

Thin Group A subgroup of finite index in its Zariski closure.

Affine Sieve A sieve method for thin groups.

Apollonian Gasket A fractal generated by repeatedly inscribing circles.

Kissing Number The number of non-overlapping unit spheres that can touch a central unit sphere.

Sphere Packing An arrangement of non-overlapping spheres in space.

Kepler Conjecture The densest sphere packing in three dimensions is the face-centered cubic packing.

Hales' Proof A computer-assisted proof of the Kepler conjecture.

Flyspeck Project A formal verification of Hales' proof.

Viazovska's Theorem The densest sphere packing in eight dimensions is the E_8 lattice packing.

Cohn-Elkies Bound A linear programming bound for sphere packing density.

Universal Optimality A property of configurations that minimize energy for all completely monotonic potentials.

Theta Function A function encoding lattice point counts.

Modular Form A function with specific transformation properties under the modular group.

Eisenstein Series A modular form defined by a sum over lattice points.

Ramanujan Tau Function The coefficients of the modular discriminant.

Ramanujan's Conjecture The tau function satisfies $|\tau(p)| \leq 2p^{11/2}$.

Deligne's Theorem A proof of Ramanujan's conjecture for weight $k \geq 2$.

Hecke Operator An operator on modular forms that commutes with the Laplacian.

Hecke Eigenform A modular form that is an eigenvector for all Hecke operators.

Newform A normalized Hecke eigenform of minimal level.

Oldform A modular form that comes from a lower level.

Atkin-Lehner Theory The theory of newforms and oldforms.

Strong Multiplicity One A theorem about the uniqueness of newforms.

Multiplicity One A theorem about the uniqueness of certain modular forms