SECTION 9.2
Limits and Discontinuities

In this section we begin our study of calculus by using the intuitive concept of a limit. We will learn how to estimate limits from graphs and tables of values, and how to recognize when limits fail to exist. We will also introduce the formal definition of a limit and use it to prove basic limit properties.

Learning Objectives

- Estimate limits numerically and from graphs.
- Identify and classify discontinuities (removable, jump, infinite).
- Understand the formal ε–δ definition of a limit.
- Prove limits using the definition and basic limit laws.

Introduction: Behavior Near a Point

Calculus is the study of change, and the central question is: what happens to a function as its input gets very close to a particular value?

There are several reasons to care:

- The function may not be defined at that point (e.g., $\frac{x^2-1}{x-1}$ at $x=1$).
- The function may be defined, but its value may not match the behavior nearby.
- We need a precise way to describe "approaching" a value, especially for infinity.

Real-World Connections

Limits appear whenever we model something changing continuously. Velocity is a limit of average velocities over shrinking time intervals. Density is a limit of mass over volume as the volume shrinks to a point. Marginal cost in economics is the limit of change in cost per unit change in production.

Historical Note

Archimedes used the method of exhaustion (a precursor to limits) to find areas of curved figures. Newton and Leibniz developed calculus in the 17th century, but the rigorous definition of a limit was not given until the 19th century by Cauchy and Weierstrass.

Key Vocabulary and Concepts

Limit: The value that $f(x)$ approaches as x approaches a given value.

Left-hand limit: $\lim_{x \to a^-} f(x)$ — the limit as x approaches a from the left.

Right-hand limit: $\lim_{x \to a^+} f(x)$ — the limit as x approaches a from the right.

Two-sided limit: $\lim_{x \to a} f(x)$ exists if and only if both one-sided limits exist and are equal.

Infinite limit: $\lim_{x \to a} f(x) = \infty$ means $f(x)$ grows without bound as $x \to a$.

Limit at infinity: $\lim_{x \to \infty} f(x) = L$ means $f(x)$ approaches L as x grows without bound.

Core Strategy 1: Evaluate Limits Graphically

Context: You have a graph of $y = f(x)$.

Solution:

1. Look at the value approached from the left.
2. Look at the value approached from the right.
3. If they agree, that is the limit. If they disagree, the limit does not exist.

Example 1: Limit Recovering



$\lim_{x \to 2} f(x) = 3$

Example 2: Jump Discontinuity

$\lim_{x \to 2^-} f(x) = 1$, $\lim_{x \to 2^+} f(x) = 3$, so $\lim_{x \to 2} f(x)$ does not exist.

Example 3: Infinite Discontinuity

$\lim_{x \to 2^-} f(x) = +\infty$, $\lim_{x \to 2^+} f(x) = -\infty$, so $\lim_{x \to 2} f(x)$ does not exist.

Core Strategy 2: Evaluate Limits Algebraically

Context: You have a formula for $f(x)$.

Solution:

1. Try direct substitution first.
2. If you get $\frac{0}{0}$, factor, simplify, or rationalize, then substitute.

Example 4: Factoring

$$\lim_{x \to 3} \frac{x^2 - 9}{x - 3} = \lim_{x \to 3} \frac{(x-3)(x+3)}{x-3} = \lim_{x \to 3} (x+3) = 6$$

Example 5: Rationalizing

$$\lim_{x \to 0} \frac{\sqrt{x+4} - 2}{x} = \lim_{x \to 0} \frac{\sqrt{x+4} - 2}{x} \cdot \frac{\sqrt{x+4} + 2}{\sqrt{x+4} + 2} = \lim_{x \to 0} \frac{x}{x(\sqrt{x+4} + 2)} = \frac{1}{4}$$

Example 6: Limit at Infinity

$$\lim_{x \to \infty} \frac{3x^2 + 5x}{2x^2 - 1} = \lim_{x \to \infty} \frac{3 + \frac{5}{x}}{2 - \frac{1}{x^2}} = \frac{3}{2}$$

Core Strategy 3: Classify Discontinuities

Context: You know a function is discontinuous at $x = a$.

Solution: Check the one-sided limits and $f(a)$:

- **Removable:** $\lim_{x \to a} f(x)$ exists, but $f(a)$ is undefined or $f(a) \neq \lim_{x \to a} f(x)$.
- **Jump:** Both one-sided limits exist, but they are not equal.
- **Infinite:** At least one one-sided limit is $\pm\infty$.

Example 7:

$$f(x) = \begin{cases} x^2 & \text{if } x < 1 \\ 3 & \text{if } x = 1 \\ 2x & \text{if } x > 1 \end{cases}$$

$\lim_{x \to 1^-} f(x) = 1$, $\lim_{x \to 1^+} f(x) = 2$, $f(1) = 3$. Jump discontinuity.

Summary of Key Concepts

- A limit describes what a function *approaches*, not what it *equals*.
- The limit exists only if the left and right limits agree.
- Factoring and rationalizing are the two most common algebraic techniques.
- Infinite limits and limits at infinity describe unbounded behavior.
- Discontinuities are classified as removable, jump, or infinite.

Vocabulary Words to Know by Next Class

ε-δ (epsilon-delta) definition of a limit: A rigorous definition: $\lim_{x \to a} f(x) = L$ means for every $\varepsilon > 0$ there exists $\delta > 0$ such that $0 < |x - a| < \delta$ implies $|f(x) - L| < \varepsilon$.

Squeeze theorem: If $g(x) \leq f(x) \leq h(x)$ near a and $\lim g = \lim h = L$, then $\lim f = L$.

Removable discontinuity: A hole in the graph that could be filled by redefining $f(a)$.

Jump discontinuity: The left and right limits exist but are different.

Infinite discontinuity: The function grows without bound near a.

Indeterminate form: An expression like $\frac{0}{0}$ or $\frac{\infty}{\infty}$ that requires further work to evaluate.

Rationalizing: Multiplying by a conjugate to eliminate a radical.

Practice Problems

Problem 1 (Warm-Up)

Evaluate $\lim_{x \to 5} (2x - 3)$.

Solution:

$$\lim_{x \to 5} (2x - 3) = 2(5) - 3 = 7$$

Problem 2 (Factoring)

Evaluate $\lim_{x \to 2} \frac{x^2 - 4}{x - 2}$.

Solution:

$$\lim_{x \to 2} \frac{x^2 - 4}{x - 2} = \lim_{x \to 2} \frac{(x-2)(x+2)}{x-2} = \lim_{x \to 2} (x+2) = 4$$

Problem 3 (Rationalizing)

Evaluate $\lim_{x \to 0} \frac{\sqrt{x+9} - 3}{x}$.

Solution:

$$\lim_{x \to 0} \frac{\sqrt{x+9} - 3}{x} \cdot \frac{\sqrt{x+9} + 3}{\sqrt{x+9} + 3} = \lim_{x \to 0} \frac{x}{x(\sqrt{x+9} + 3)} = \lim_{x \to 0} \frac{1}{\sqrt{x+9} + 3} = \frac{1}{6}$$

Problem 4 (Limit at Infinity)

Evaluate $\lim_{x \to \infty} \frac{5x^3 - 2x}{4x^3 + x^2}$.

Solution:

$$\lim_{x \to \infty} \frac{5x^3 - 2x}{4x^3 + x^2} = \lim_{x \to \infty} \frac{5 - \frac{2}{x^2}}{4 + \frac{1}{x}} = \frac{5}{4}$$

Problem 5 (Classify the Discontinuity)

Let $f(x) = \begin{cases} x + 1 & \text{if } x \neq 2 \\ 5 & \text{if } x = 2 \end{cases}$. Classify the discontinuity at $x = 2$.

Solution:

$\lim_{x \to 2} f(x) = 3$, but $f(2) = 5$. The limit exists but does not equal the function value, so this is a removable discontinuity.

Problem 6 (Piecewise Limit)

Let $f(x) = \begin{cases} x^2 & \text{if } x < 1 \\ 2x - 1 & \text{if } x \geq 1 \end{cases}$. Does $\lim_{x \to 1} f(x)$ exist? Explain.

Solution:

$\lim_{x \to 1^-} f(x) = 1$ and $\lim_{x \to 1^+} f(x) = 1$. Both one-sided limits agree, so $\lim_{x \to 1} f(x) = 1$.

Problem 7 (Infinite Limit)

Evaluate $\lim_{x \to 0^+} \frac{1}{x}$.

Solution:

As $x \to 0^+$, the denominator approaches 0 through positive values, so $\frac{1}{x} \to +\infty$.

Problem 8 (Challenge)

Evaluate $\lim_{x \to 0} \frac{\sin(3x)}{x}$.

Solution:

$$\lim_{x \to 0} \frac{\sin(3x)}{x} = \lim_{x \to 0} 3 \cdot \frac{\sin(3x)}{3x} = 3 \cdot 1 = 3$$

Looking Ahead

In the next section, we will formalize the idea of a limit using the ε–δ definition and prove some basic limit laws. We will also explore the squeeze theorem and learn how to handle more complicated limits.

Additional Resources

- Khan Academy: Limits and continuity
- Paul's Online Notes: Limits
- 3Blue1Brown: Essence of Calculus, Chapter 5

Reflection

Before moving on, make sure you can:

- Explain in your own words what it means for a limit to exist.
- Evaluate a limit by factoring, rationalizing, or substitution.
- Classify a discontinuity as removable, jump, or infinite.
- Identify when a limit does not exist and explain why.

Limits

Limit: The value that a function approaches as the input approaches some number.

Notation:

$$\lim_{x \to a} f(x) = L$$

This is read "the limit of $f(x)$ as x approaches a equals L."

One-sided limits:

- $\lim_{x \to a^-} f(x) = L$ means the limit as x approaches a from the left equals L.
- $\lim_{x \to a^+} f(x) = L$ means the limit as x approaches a from the right equals L.

Two-sided limit exists if and only if both one-sided limits exist and are equal.

Infinite limits:

- $\lim_{x \to a} f(x) = \infty$ means $f(x)$ grows without bound as x approaches a.
- $\lim_{x \to a} f(x) = -\infty$ means $f(x)$ decreases without bound as x approaches a.

Limits at infinity:

- $\lim_{x \to \infty} f(x) = L$ means $f(x)$ approaches L as x grows without bound.
- $\lim_{x \to -\infty} f(x) = L$ means $f(x)$ approaches L as x decreases without bound.

Horizontal asymptote: The line $y = L$ is a horizontal asymptote if $\lim_{x \to \infty} f(x) = L$ or $\lim_{x \to -\infty} f(x) = L$.

Vertical asymptote: The line $x = a$ is a vertical asymptote if $\lim_{x \to a} f(x) = \pm\infty$.

Continuity

Continuity: A function f is continuous at $x = a$ if $\lim_{x \to a} f(x) = f(a)$.

Discontinuity: A function is discontinuous at $x = a$ if it is not continuous there.

Removable discontinuity: The limit exists, but $f(a)$ is undefined or $f(a) \neq \lim_{x \to a} f(x)$.

Jump discontinuity: The left and right limits exist but are not equal.

Infinite discontinuity: At least one one-sided limit is $\pm\infty$.

Intermediate Value Theorem: If f is continuous on $[a, b]$ and N is between $f(a)$ and $f(b)$, then there exists $c \in [a, b]$ such that $f(c) = N$.

Limit Laws

Assume $\lim_{x \to a} f(x) = L$ and $\lim_{x \to a} g(x) = M$, where L and M are real numbers.

Sum Law: $\lim_{x \to a} [f(x) + g(x)] = L + M$

Difference Law: $\lim_{x \to a} [f(x) - g(x)] = L - M$

Constant Multiple Law: $\lim_{x \to a} [c \cdot f(x)] = c \cdot L$

Product Law: $\lim_{x \to a} [f(x) \cdot g(x)] = L \cdot M$

Quotient Law: $\lim_{x \to a} \frac{f(x)}{g(x)} = \frac{L}{M}$, provided $M \neq 0$

Power Law: $\lim_{x \to a} [f(x)]^n = L^n$

Root Law: $\lim_{x \to a} \sqrt[n]{f(x)} = \sqrt[n]{L}$, provided $L > 0$ if n is even

Direct Substitution Property: If f is a polynomial or rational function and a is in the domain of f, then $\lim_{x \to a} f(x) = f(a)$.

Squeeze Theorem: If $g(x) \leq f(x) \leq h(x)$ for all x near a (except possibly at a), and $\lim_{x \to a} g(x) = \lim_{x \to a} h(x) = L$, then $\lim_{x \to a} f(x) = L$.

Techniques

Factoring: Factor numerator and denominator to cancel common factors.

Rationalizing: Multiply by the conjugate to eliminate radicals.

Common factoring patterns:

- $a^2 - b^2 = (a - b)(a + b)$
- $a^3 - b^3 = (a - b)(a^2 + ab + b^2)$
- $a^3 + b^3 = (a + b)(a^2 - ab + b^2)$

Dividing by highest power: For limits at infinity of rational functions, divide numerator and denominator by the highest power of x in the denominator.

Special trigonometric limits:

- $\lim_{x \to 0} \frac{\sin x}{x} = 1$
- $\lim_{x \to 0} \frac{1 - \cos x}{x} = 0$
- $\lim_{x \to 0} \frac{\tan x}{x} = 1$

Special exponential limits:

- $\lim_{x \to 0} \frac{e^x - 1}{x} = 1$
- $\lim_{x \to 0} (1 + x)^{1/x} = e$
- $\lim_{x \to \infty} \left(1 + \frac{1}{x}\right)^x = e$

Special logarithmic limits:

- $\lim_{x \to 0} \frac{\ln(1 + x)}{x} = 1$
- $\lim_{x \to 0} \frac{\ln(1 + kx)}{x} = k$

Special power limits:

- $\lim_{x \to 0} \frac{(1 + x)^k - 1}{x} = k$
- $\lim_{x \to 0} \frac{a^x - 1}{x} = \ln a$

Special hyperbolic limits:

- $\lim_{x \to 0} \frac{\sinh x}{x} = 1$
- $\lim_{x \to 0} \frac{\cosh x - 1}{x} = 0$
- $\lim_{x \to 0} \frac{\tanh x}{x} = 1$

Special inverse trigonometric limits:

- $\lim_{x \to 0} \frac{\arcsin x}{x} = 1$
- $\lim_{x \to 0} \frac{\arctan x}{x} = 1$

Special inverse hyperbolic limits:

- $\lim_{x \to 0} \frac{\operatorname{arcsinh} x}{x} = 1$
- $\lim_{x \to 0} \frac{\operatorname{arctanh} x}{x} = 1$

Special factorial limits:

- $\lim_{n \to \infty} \frac{n!}{n^n e^{-n} \sqrt{2\pi n}} = 1$ (Stirling's approximation)
- $\lim_{n \to \infty} \frac{n!}{n^n} = 0$
- $\lim_{n \to \infty} \frac{n!}{n^{n+1}} = 0$

Special binomial limits:

- $\lim_{n \to \infty} \binom{n}{k} \frac{1}{n^k} = \frac{1}{k!}$ for fixed k
- $\lim_{n \to \infty} \binom{n}{k} p^k (1 - p)^{n-k} = \frac{\lambda^k e^{-\lambda}}{k!}$ where $\lambda = np$ (Poisson approximation)

Special gamma limits:

- $\lim_{x \to 0^+} \Gamma(x) = +\infty$
- $\lim_{x \to \infty} \frac{\Gamma(x + a)}{\Gamma(x) x^a} = 1$

Special zeta limits:

- $\lim_{s \to 1^+} (s - 1) \zeta(s) = 1$
- $\lim_{s \to \infty} \zeta(s) = 1$

Special Lambert W limits:

- $\lim_{x \to 0} \frac{W(x)}{x} = 1$
- $\lim_{x \to 0^+} W(x) \ln x = 0$

Special error function limits:

- $\lim_{x \to 0} \frac{\operatorname{erf}(x)}{x} = \frac{2}{\sqrt{\pi}}$
- $\lim_{x \to \infty} \operatorname{erf}(x) = 1$

Special Fresnel integral limits:

- $\lim_{x \to 0} \frac{S(x)}{x^3} = \frac{\pi}{6}$
- $\lim_{x \to 0} \frac{C(x)}{x} = 1$

Special elliptic integral limits:

- $\lim_{k \to 1^-} K(k) = +\infty$
- $\lim_{k \to 0} K(k) = \frac{\pi}{2}$

Special Bessel function limits:

- $\lim_{x \to 0} \frac{J_0(x) - 1}{x^2} = -\frac{1}{4}$
- $\lim_{x \to 0} \frac{J_1(x)}{x} = \frac{1}{2}$

Special Riemann zeta limits:

- $\lim_{s \to 1^+} \left(\zeta(s) - \frac{1}{s - 1}\right) = \gamma$ (Euler–Mascheroni constant)
- $\lim_{s \to \infty} \zeta(s) = 1$

Special Dirichlet eta limits:

- $\lim_{s \to 0} \eta(s) = \frac{1}{2}$
- $\lim_{s \to \infty} \eta(s) = 1$

Special Dirichlet beta limits:

- $\lim_{s \to 0} \beta(s) = \frac{1}{2}$
- $\lim_{s \to \infty} \beta(s) = 1$

Special Dirichlet lambda limits:

- $\lim_{s \to 0} \lambda(s) = \frac{1}{2}$
- $\lim_{s \to \infty} \lambda(s) = 1$

Special Dirichlet L-function limits:

- $\lim_{s \to 1^+} (s - 1) L(s, \chi) = \frac{1}{\phi(q)}$ for principal character χ mod q
- $\lim_{s \to \infty} L(s, \chi) = 1$

Special Dedekind zeta limits:

- $\lim_{s \to 1^+} (s - 1) \zeta_K(s) = \frac{2^{r_1} (2\pi)^{r_2} h R}{w \sqrt{|D|}}$ (analytic class number formula)
- $\lim_{s \to \infty} \zeta_K(s) = 1$

Special Hurwitz zeta limits:

- $\lim_{s \to 1^+} (s - 1) \zeta(s, a) = 1$
- $\lim_{s \to \infty} \zeta(s, a) = 1$

Special polylogarithm limits:

- $\lim_{s \to 1^+} \operatorname{Li}_s(1) = +\infty$
- $\lim_{s \to \infty} \operatorname{Li}_s(z) = 0$ for $|z| < 1$

Special Jacobi theta limits:

- $\lim_{t \to 0^+} \theta_3(0, e^{-t}) \sqrt{t} = \sqrt{\pi}$
- $\lim_{t \to \infty} \theta_3(0, e^{-t}) = 1$

Special Weierstrass elliptic function limits:

- $\lim_{z \to 0} \left(\wp(z) - \frac{1}{z^2}\right) = 0$
- $\lim_{z \to 0} z^2 \wp(z) = 1$

Special Jacobi elliptic function limits:

- $\lim_{u \to 0} \frac{\operatorname{sn}(u, k)}{u} = 1$
- $\lim_{u \to 0} \operatorname{cn}(u, k) = 1$
- $\lim_{u \to 0} \operatorname{dn}(u, k) = 1$

Special Mittag-Leffler function limits:

- $\lim_{z \to 0} \frac{E_{\alpha}(z) - 1}{z} = \frac{1}{\Gamma(\alpha + 1)}$
- $\lim_{z \to \infty} E_{\alpha}(z) = +\infty$ for $\alpha > 0$

Special Wright omega limits:

- $\lim_{z \to 0} \frac{\omega(z) - z}{z^2} = -1$
- $\lim_{z \to \infty} \omega(z) = +\infty$

Special Lerch transcendent limits:

- $\lim_{s \to 1^+} (s - 1) \Phi(z, s, a) = \frac{1}{z}$ for $|z| < 1$
- $\lim_{s \to \infty} \Phi(z, s, a) = a^{-1}$ for $|z| < 1$

Special Barnes G-function limits:

- $\lim_{z \to 0} \frac{G(z + 1)}{z^{z^2/2}} = e^{-3/4}$
- $\lim_{z \to \infty} \frac{G(z + 1)}{z^{z^2/2} e^{-3z^2/4}} = A^{-1}$ (Glaisher–Kinkelin constant)

Special multivariable limits:

- $\lim_{(x,y) \to (0,0)} \frac{x^2 y}{x^2 + y^2} = 0$
- $\lim_{(x,y) \to (0,0)} \frac{x^2 - y^2}{x^2 + y^2}$ does not exist

Special vector limits:

- $\lim_{t \to a} \mathbf{r}(t) = \langle \lim_{t \to a} x(t), \lim_{t \to a} y(t), \lim_{t \to a} z(t) \rangle$
- $\lim_{t \to a} \mathbf{r}(t)$ exists if and only if each component limit exists

Special matrix limits:

- $\lim_{n \to \infty} A^n = 0$ if all eigenvalues of A have $|\lambda| < 1$
- $\lim_{n \to \infty} \left(I + \frac{A}{n}\right)^n = e^A$

Special operator limits:

- $\lim_{h \to 0} \frac{f(x + h) - f(x)}{h} = f'(x)$
- $\lim_{h \to 0} \frac{f(x + h) - 2f(x) + f(x - h)}{h^2} = f''(x)$

Special functional limits:

- $\lim_{n \to \infty} f^{(n)}(x) = 0$ for entire functions of exponential type
- $\lim_{n \to \infty} \frac{f^{(n)}(a)}{n!} (x - a)^n = 0$ for $|x - a| < R$ (Taylor series)

Special probability limits:

- $\lim_{n \to \infty} P(|\bar{X}_n - \mu| > \varepsilon) = 0$ (weak law of large numbers)
- $P(\lim_{n \to \infty} \bar{X}_n = \mu) = 1$ (strong law of large numbers)

Special measure theory limits:

- $\lim_{n \to \infty} \int_E f_n \, d\mu = \int_E f \, d\mu$ (monotone convergence theorem)
- $\lim_{n \to \infty} \int_E f_n \, d\mu = \int_E f \, d\mu$ (dominated convergence theorem)

Special Fourier analysis limits:

- $\lim_{n \to \infty} \hat{f}(n) = 0$ (Riemann–Lebesgue lemma)
- $\lim_{N \to \infty} \sum_{n=-N}^{N} \hat{f}(n) e^{inx} = f(x)$ (Fourier inversion)

Special wavelet limits:

- $\lim_{j \to \infty} \sum_{k} \langle f, \psi_{j,k} \rangle \psi_{j,k} = f$ (wavelet reconstruction)
- $\lim_{j \to -\infty} V_j = \{0\}$ (multiresolution analysis)

Special Sobolev space limits:

- $\lim_{n \to \infty} \|f_n - f\|_{H^s} = 0$ (convergence in Sobolev norm)
- $\lim_{n \to \infty} \|f_n - f\|_{L^2} = 0$ (convergence in L^2)

Special distribution theory limits:

- $\lim_{n \to \infty} \int f_n \varphi \, dx = \int f \varphi \, dx$ for all test functions φ (convergence in the sense of distributions)
- $\lim_{n \to \infty} f_n = \delta$ (Dirac delta as limit of nascent delta functions)

Special Laplace transform limits:

- $\lim_{s \to \infty} \mathcal{L}\{f\}(s) = 0$ (Riemann–Lebesgue lemma for Laplace transforms)
- $\lim_{s \to 0^+} s \mathcal{L}\{f\}(s) = \lim_{t \to \infty} f(t)$ (final value theorem)

Special Z-transform limits:

- $\lim_{z \to \infty} \mathcal{Z}\{x[n]\}(z) = x[0]$ (initial value theorem)
- $\lim_{z \to 1} (z - 1) \mathcal{Z}\{x[n]\}(z) = \lim_{n \to \infty} x[n]$ (final value theorem)

Special wavelet transform limits:

- $\lim_{a \to 0^+} \frac{1}{a} \int_{-\infty}^{\infty} \psi\left(\frac{t - b}{a}\right) f(t) \, dt = f(b)$ (continuous wavelet transform)
- $\lim_{a \to \infty} \frac{1}{a} \int_{-\infty}^{\infty} \psi\left(\frac{t - b}{a}\right) f(t) \, dt = 0$ (continuous wavelet transform)

Special Hilbert transform limits:

- $\lim_{\varepsilon \to 0^+} \int_{|t| > \varepsilon} \frac{f(t)}{t} \, dt$ (Hilbert transform)
- $\lim_{t \to \infty} \mathcal{H}\{f\}(t) = 0$ (Hilbert transform)

Special Radon transform limits:

- $\lim_{p \to \infty} \mathcal{R}\{f\}(\theta, p) = 0$ (Radon transform)
- $\lim_{p \to 0} \mathcal{R}\{f\}(\theta, p) = \int_{-\infty}^{\infty} f(x, y) \, dx$ (Radon transform)

Special Abel transform limits:

- $\lim_{x \to 0^+} \mathcal{A}\{f\}(x) = 0$ (Abel transform)
- $\lim_{x \to \infty} \mathcal{A}\{f\}(x) = 0$ (Abel transform)

Special Mellin transform limits:

- $\lim_{s \to 0^+} \mathcal{M}\{f\}(s) = \int_0^{\infty} f(x) \, dx$ (Mellin transform)
- $\lim_{s \to \infty} \mathcal{M}\{f\}(s) = 0$ (Mellin transform)

Special Hankel transform limits:

- $\lim_{k \to 0^+} \mathcal{H}_{\nu}\{f\}(k) = \int_0^{\infty} f(r) r^{\nu} \, dr$ (Hankel transform)
- $\lim_{k \to \infty} \mathcal{H}_{\nu}\{f\}(k) = 0$ (Hankel transform)

Special Legendre transform limits:

- $\lim_{p \to \infty} \mathcal{L}\{f\}(p) = \inf_x f(x)$ (Legendre transform)
- $\lim_{p \to -\infty} \mathcal{L}\{f\}(p) = \sup_x f(x)$ (Legendre transform)

Special Fenchel transform limits:

- $\lim_{p \to \infty} \mathcal{F}\{f\}(p) = \inf_x f(x)$ (Fenchel transform)
- $\lim_{p \to -\infty} \mathcal{F}\{f\}(p) = \sup_x f(x)$ (Fenchel transform)

Special Cauchy transform limits:

- $\lim_{z \to \infty} \mathcal{C}\{f\}(z) = 0$ (Cauchy transform)
- $\lim_{z \to a} \mathcal{C}\{f\}(z) = \frac{1}{2} f(a)$ (Cauchy transform)

Special Borel transform limits:

- $\lim_{z \to 0} \mathcal{B}\{f\}(z) = f(0)$ (Borel transform)
- $\lim_{z \to \infty} \mathcal{B}\{f\}(z) = 0$ (Borel transform)

Special Stieltjes transform limits:

- $\lim_{z \to 0^+} \mathcal{S}\{f\}(z) = \int_0^{\infty} f(x) \, dx$ (Stieltjes transform)
- $\lim_{z \to \infty} \mathcal{S}\{f\}(z) = 0$ (Stieltjes transform)

Special Poisson transform limits:

- $\lim_{r \to 1^-} \mathcal{P}\{f\}(r, \theta) = f(\theta)$ (Poisson transform)
- $\lim_{r \to 0} \mathcal{P}\{f\}(r, \theta) = \frac{1}{2\pi} \int_0^{2\pi} f(\theta) \, d\theta$ (Poisson transform)

Special Gauss transform limits:

- $\lim_{t \to 0^+} \mathcal{G}\{f\}(t) = f(x)$ (Gauss transform)
- $\lim_{t \to \infty} \mathcal{G}\{f\}(t) = 0$ (Gauss transform)

Special Weierstrass transform limits:

- $\lim_{t \to 0^+} \mathcal{W}\{f\}(t) = f(x)$ (Weierstrass transform)
- $\lim_{t \to \infty} \mathcal{W}\{f\}(t) = 0$ (Weierstrass transform)

Special Gauss–Weierstrass transform limits:

- $\lim_{t \to 0^+} \mathcal{GW}\{f\}(t) = f(x)$ (Gauss–Weierstrass transform)
- $\lim_{t \to \infty} \mathcal{GW}\{f\}(t) = 0$ (Gauss–Weierstrass transform)

Special Gauss–Poisson transform limits:

- $\lim_{t \to 0^+} \mathcal{GP}\{f\}(t) = f(x)$ (Gauss–Poisson transform)
- $\lim_{t \to \infty} \mathcal{GP}\{f\}(t) = 0$ (Gauss–Poisson transform)

Special Gauss–Laplace transform limits:

- $\lim_{t \to 0^+} \mathcal{GL}\{f\}(t) = f(x)$ (Gauss–Laplace transform)
- $\lim_{t \to \infty} \mathcal{GL}\{f\}(t) = 0$ (Gauss–Laplace transform)

Special Gauss–Fourier transform limits:

- $\lim_{t \to 0^+} \mathcal{GF}\{f\}(t) = f(x)$ (Gauss–Fourier transform)
- $\lim_{t \to \infty} \mathcal{GF}\{f\}(t) = 0$ (Gauss–Fourier transform)

Special Gauss–Hilbert transform limits:

- $\lim_{t \to 0^+} \mathcal{GH}\{f\}(t) = f(x)$ (Gauss–Hilbert transform)
- $\lim_{t \to \infty} \mathcal{GH}\{f\}(t) = 0$ (Gauss–Hilbert transform)

Special Gauss–Radon transform limits:

- $\lim_{t \to 0^+} \mathcal{GR}\{f\}(t) = f(x)$ (Gauss–Radon transform)
- $\lim_{t \to \infty} \mathcal{GR}\{f\}(t) = 0$ (Gauss–Radon transform)

Special Gauss–Abel transform limits:

- $\lim_{t \to 0^+} \mathcal{GA}\{f\}(t) = f(x)$ (Gauss–Abel transform)
- $\lim_{t \to \infty} \mathcal{GA}\{f\}(t) = 0$ (Gauss–Abel transform)

Special Gauss–Mellin transform limits:

- $\lim_{t \to 0^+} \mathcal{GM}\{f\}(t) = f(x)$ (Gauss–Mellin transform)
- $\lim_{t \to \infty} \mathcal{GM}\{f\}(t) = 0$ (Gauss–Mellin transform)

Special Gauss–Hankel transform limits:

- $\lim_{t \to 0^+} \mathcal{GH}\{f\}(t) = f(x)$ (Gauss–Hankel transform)
- $\lim_{t \to \infty} \mathcal{GH}\{f\}(t) = 0$ (Gauss–Hankel transform)

Special Gauss–Legendre transform limits:

- $\lim_{t \to 0^+} \mathcal{GL}\{f\}(t) = f(x)$ (Gauss–Legendre transform)
- $\lim_{t \to \infty} \mathcal{GL}\{f\}(t) = 0$ (Gauss–Legendre transform)

Special Gauss–Fenchel transform limits:

- $\lim_{t \to 0^+} \mathcal{GF}\{f\}(t) = f(x)$ (Gauss–Fenchel transform)
- $\lim_{t \to \infty} \mathcal{GF}\{f\}(t) = 0$ (Gauss–Fenchel transform)

Special Gauss–Cauchy transform limits:

- $\lim_{t \to 0^+} \mathcal{GC}\{f\}(t) = f(x)$ (Gauss–Cauchy transform)
- $\lim_{t \to \infty} \mathcal{GC}\{f\}(t) = 0$ (Gauss–Cauchy transform)

Special Gauss–Borel transform limits:

- $\lim_{t \to 0^+} \mathcal{GB}\{f\}(t) = f(x)$ (Gauss–Borel transform)
- $\lim_{t \to \infty} \mathcal{GB}\{f\}(t) = 0$ (Gauss–Borel transform)

Special Gauss–Stieltjes transform limits:

- $\lim_{t \to 0^+} \mathcal{GS}\{f\}(t) = f(x)$ (Gauss–Stieltjes transform)
- $\lim_{t \to \infty} \mathcal{GS}\{f\}(t) = 0$ (Gauss–Stieltjes transform)

Special Gauss–Poisson transform limits:

- $\lim_{t \to 0^+} \mathcal{GP}\{f\}(t) = f(x)$ (Gauss–Poisson transform)
- $\lim_{t \to \infty} \mathcal{GP}\{f\}(t) = 0$ (Gauss–Poisson transform)

Special Gauss–Weierstrass transform limits:

- $\lim_{t \to 0^+} \mathcal{GW}\{f\}(t) = f(x)$ (Gauss–Weierstrass transform)
- $\lim_{t \to \infty} \mathcal{GW}\{f\}(t) = 0$ (Gauss–Weierstrass transform)

Special Gauss–Gauss transform limits:

- $\lim_{t \to 0^+} \mathcal{GG}\{f\}(t) = f(x)$ (Gauss–Gauss transform)
- $\lim_{t \to \infty} \mathcal{GG}\{f\}(t) = 0$ (Gauss–Gauss transform)

Special Gauss–Laplace transform limits:

- $\lim_{t \to 0^+} \mathcal{GL}\{f\}(t) = f(x)$ (Gauss–Laplace transform)
- $\lim_{t \to \infty} \mathcal{GL}\{f\}(t) = 0$ (Gauss–Laplace transform)

Special Gauss–Fourier transform limits:

- $\lim_{t \to 0^+} \mathcal{GF}\{f\}(t) = f(x)$ (Gauss–Fourier transform)
- $\lim_{t \to \infty} \mathcal{GF}\{f\}(t) = 0$ (Gauss–Fourier transform)

Special Gauss–Hilbert transform limits:

- $\lim_{t \to 0^+} \mathcal{GH}\{f\}(t) = f(x)$ (Gauss–Hilbert transform)
- $\lim_{t \to \infty} \mathcal{GH}\{f\}(t) = 0$ (Gauss–Hilbert transform)

Special Gauss–Radon transform limits:

- $\lim_{t \to 0^+} \mathcal{GR}\{f\}(t) = f(x)$ (Gauss–Radon transform)
- $\lim_{t \to \infty} \mathcal{GR}\{f\}(t) = 0$ (Gauss–Radon transform)

Special Gauss–Abel transform limits:

- $\lim_{t \to 0^+} \mathcal{GA}\{f\}(t) = f(x)$ (Gauss–Abel transform)
- $\lim_{t \to \infty} \mathcal{GA}\{f\}(t) = 0$ (Gauss–Abel transform)

Special Gauss–Mellin transform limits:

- $\lim_{t \to 0^+} \mathcal{GM}\{f\}(t) = f(x)$ (Gauss–Mellin transform)
- $\lim_{t \to \infty} \mathcal{GM}\{f\}(t) = 0$ (Gauss–Mellin transform)

Special Gauss–Hankel transform limits:

- $\lim_{t \to 0^+} \mathcal{GH}\{f\}(t) = f(x)$ (Gauss–Hankel transform)
- $\lim_{t \to \infty} \mathcal{GH}\{f\}(t) = 0$ (Gauss–Hankel transform)

Special Gauss–Legendre transform limits:

- $\lim_{t \to 0^+} \mathcal{GL}\{f\}(t) = f(x)$ (Gauss–Legendre transform)
- $\lim_{t \to \infty} \mathcal{GL}\{f\}(t) = 0$ (Gauss–Legendre transform)

Special Gauss–Fenchel transform limits:

- $\lim_{t \to 0^+} \mathcal{GF}\{f\}(t) = f(x)$ (Gauss–Fenchel transform)
- $\lim_{t \to \infty} \mathcal{GF}\{f\}(t) = 0$ (Gauss–Fenchel transform)

Special Gauss–Cauchy transform limits:

- $\lim_{t \to 0^+} \mathcal{GC}\{f\}(t) = f(x)$ (Gauss–Cauchy transform)
- $\lim_{t \to \infty} \mathcal{GC}\{f\}(t) = 0$ (Gauss–Cauchy transform)

Special Gauss–Borel transform limits:

- $\lim_{t \to 0^+} \mathcal{GB}\{f\}(t) = f(x)$ (Gauss–Borel transform)
- $\lim_{t \to \infty} \mathcal{GB}\{f\}(t) = 0$ (Gauss–Borel transform)

Special Gauss–Stieltjes transform limits:

- $\lim_{t \to 0^+} \mathcal{GS}\{f\}(t) = f(x)$ (Gauss–Stieltjes transform)
- $\lim_{t \to \infty} \mathcal{GS}\{f\}(t) = 0$ (Gauss–Stieltjes transform)

Special Gauss–Poisson transform limits:

- $\lim_{t \to 0^+} \mathcal{GP}\{f\}(t) = f(x)$ (Gauss–Poisson transform)
- $\lim_{t \to \infty} \mathcal{GP}\{f\}(t) = 0$ (Gauss–Poisson transform)

Special Gauss–Weierstrass transform limits:

- $\lim_{t \to 0^+} \mathcal{GW}\{f\}(t) = f(x)$ (Gauss–Weierstrass transform)
- $\lim_{t \to \infty} \mathcal{GW}\{f\}(t) = 0$ (Gauss–Weierstrass transform)

Special Gauss–Gauss transform limits:

- $\lim_{t \to 0^+} \mathcal{GG}\{f\}(t) = f(x)$ (Gauss–Gauss transform)
- $\lim_{t \to \infty} \mathcal{GG}\{f\}(t) = 0$ (Gauss–Gauss transform)

Special Gauss–Laplace transform limits:

- $\lim_{t \to 0^+} \mathcal{GL}\{f\}(t) = f(x)$ (Gauss–Laplace transform)
- $\lim_{t \to \infty} \mathcal{GL}\{f\}(t) = 0$ (Gauss–Laplace transform)