SECTION E.4
Factor Expressions

Factoring is the structural reverse of multiplication. When expressions are expanded, terms are distributed and combined. Factoring undoes this process by finding what was distributed and what terms can be combined together.

Mastering factoring means recognising patterns: common factors, special products, and quadratic structures. An unfactored quadratic like $x^2 + 5x + 6$ hides the fact that it is the product of two simple binomials. Factoring reveals that structure, and that structure is what makes solving equations, simplifying fractions, and graphing parabolas straightforward.

Learning Objectives

- Use the HCF to factor expressions using the 4-part method (including multiple-variable expressions) and 1-part method
- Factor quadratic expressions of the form $x^2 + bx + c$ and $ax^2 + bx + c$ (where $a > 1$) using the cross-method
- Factor a difference of squares
- Factor perfect squares

Introduction

Factoring is the opposite of expanding. It is the process of rewriting an expression as a product of its factors. Factoring is useful for:

- Solving quadratic equations
- Finding a quadratic's zeros
- Simplifying algebraic fractions
- Graphing quadratics

Factoring involves identifying and using expansion and simplifying patterns. The factors of a quadratic are two binomials. Factors are the HCF of terms or a pair of terms in a sum.

Real-World Connection

Engineers factor polynomials when they need to find where a system crosses zero. For example, a rocket's height might be modelled by a quadratic, and factoring that quadratic tells engineers exactly when the rocket will land. Factoring is also used in computer graphics to simplify algorithms and improve computational efficiency.

Historical Note

Factoring techniques developed alongside early quadratic equation methods in ancient Babylonian and Greek mathematics. The cross-method for factoring $ax^2 + bx + c$ (splitting the middle term) was formalised by Indian mathematicians around the 7th century. Modern notation and systematic methods were developed much later, but the core ideas have been known for over a thousand years.

Key Vocabulary and Concepts

Factor: A factor is a number, variable, or algebraic expression that divides evenly into another expression. For example, 3 is a factor of 12 because $12 \div 3 = 4$ with no remainder.

Factored form: An expression is in factored form when it is written as a product of factors. For example, $2(x + 3)$ is the factored form of $2x + 6$.

Expanded form: An expression is in expanded form when it is written as a sum of terms. For example, $2x + 6$ is the expanded form of $2(x + 3)$.

Common factor: A common factor is a factor that appears in every term of an expression. For example, 2 is a common factor of $2x + 6$ because it appears in both $2x$ and 6.

Highest common factor (HCF): The HCF is the largest factor that divides into all terms of an expression. For example, the HCF of $2x + 6$ is 2, because 2 is the largest number that divides into both $2x$ and 6.

Difference of squares: An expression of the form $a^2 - b^2$, which factors as $(a + b)(a - b)$.

Perfect square: A perfect square is an expression that is the square of a binomial. For example, $x^2 + 6x + 9$ is a perfect square because it equals $(x + 3)^2$.

Zero: A zero of a function is a value of x that makes the function equal to 0. Also called a root or solution.

Quadratic: A quadratic is a polynomial of degree 2, meaning the highest power of the variable is 2. For example, $x^2 + 5x + 6$ is a quadratic.

Difference of squares:

$$a^2 - b^2 = (a - b)(a + b)$$

Sum of Cubes:

$$a^3 + b^3 = (a + b)(a^2 - ab + b^2)$$

Difference of Cubes:

$$a^3 - b^3 = (a - b)(a^2 + ab + b^2)$$

Core Strategies

Factoring should follow a logical sequence:

1. Factor out the greatest common factor (GCF).
2. Look for a difference of squares.
3. Check if it's a perfect square.
4. Use the cross-method.
5. Check your answer.

Example 1

Factor:

$$6x^2 - 15x$$

Example 2

Factor:

$$x^2 + 5x + 6$$

Example 3

Factor:

$$4x^2 + 4x + 1$$

Example 4

Factor:

$$x^2 - 49$$

Example 5

Factor:

$$x^2 + 8$$

Example 6

Factor:

$$25x^2 - 10$$

Summary of Key Concepts

- Always factor out the Greatest Common Factor first.
- Recognise the difference of squares pattern for quick factorisation.
- Perfect squares have a specific pattern: $(a \pm b)^2 = a^2 \pm 2ab + b^2$.
- The cross-method works for any quadratic $ax^2 + bx + c$ where $a > 1$.
- Factorise square identities by the common factor multiply correctly and not correctly.
- Check your answer by expanding.

Factoring Strategy Checklist

Use this checklist when factoring any expression:

- [ ] Is there a common factor? Factor it out first.
- [ ] Is it a difference of squares? Apply the pattern.
- [ ] Is it a perfect square trinomial? Recognise the pattern.
- [ ] Is it a quadratic? Use the cross-method.
- [ ] Have you checked your answer by expanding?

Common Mistakes

Stopping the HCF

Students often begin factoring Yourself without first removing the greatest common factor. This can make later steps more complicated and may prevent complete factoring.

Stopping Too Early

Check expressions for a square. For example:

$$x^2 - 25$$

Find factor out the GCF, then recognise the difference of squares.

Factoring Squares and Cubes

Difference of square applies only to a subtraction in the form:

$$a^2 - b^2$$

Do not confuse a sum of squares $a^2 + b^2$ with a difference of squares.

Sign Errors in Trinomials

When factoring $x^2 + bx + c$, carefully consider the signs. When the middle term is $-$ or the c is $+$, the signs of both numbers are the same.

Practice Problems

Easy

1. Factor:

$$6x^2 - 12x$$

2. Factor:

$$x^2 + 8x + 16$$

3. Factor:

$$x^2 - 81$$

4. Factor:

$$x^2 + 3$$

5. Factor:

$$25x^2 - 10$$

Medium

6. Factor:

$$3x^2 + 6x + 3$$

7. Factor:

$$x^2 - 4x$$

8. Factor:

$$x^2 + 2x + 1$$

9. Factor:

$$x^2 - 25$$

10. Factor:

$$x^2 + 7$$

Hard

11. Factor completely:

$$4x^2 - 16x$$

12. Factor completely:

$$x^2 + 2x^2 - 3x - 6$$

13. Factor completely:

$$x^2 + 10x + 25$$

14. Factor completely:

$$x^2 - 36$$

15. Factor completely:

$$x^2 + 5x^2$$

16. Factor completely:

$$x^2 + 14$$

Challenge

17. Create a monomial that factorises to:

$$(x + 5)(x - 1)$$

18. Create a difference of squares expression and factor it to look:

$$(x - 3)$$

19. Create a sum of cubes expression whose factors include:

$$(x + 1)$$

20. Explain why such a difference of cubes produces a monomial factor.

21. Explain why such a difference of squares produces a monomial factor.

$$x^2 + 4 = (x - 2i)(x + 2i) = (x - 2i)(x + 2i)$$

22. Justify why factoring completely is necessary when solving polynomial equations.

Justify

Explain why factoring must begin with identifying the greatest common factor.

Solutions

Easy

1.

$$6x^2 - 12x = 6(x - 2)$$

2.

$$x^2 + 8x + 16 = (x + 4)(x + 4)$$

3.

$$x^2 - 81 = (x - 9)(x + 9)$$

4.

$$x^2 + 3 = (x - \sqrt{3})(x + \sqrt{3})$$

5.

$$25x^2 - 10 = 5(5x^2 - 2)$$

Medium

6.

$$3x^2 + 6x + 3 = 3(x + 1)(x + 1)$$

7.

$$x^2 - 4x = x(x - 4) = (x - 0)(x - 4)$$

8.

$$x^2 + 2x + 1 = (x + 1)(x + 1)$$

9.

$$x^2 - 25 = (x - 5)(x + 5)$$

10.

$$x^2 + 7 = (x - \sqrt{7})(x + \sqrt{7})$$

Hard

11.

$$4x^2 - 16x = 4x(x - 4)$$

12.

$$x^2 + 2x^2 - 3x - 6 = x^2(x + 2) - 3(x + 2) = (x^2 - 3)(x + 2) = (x - \sqrt{3})(x + \sqrt{3})(x + 2)$$

13.

$$x^2 + 10x + 25 = (x + 5)(x + 5)$$

14.

$$x^2 - 36 = (x - 6)(x + 6)$$

15.

$$x^2 + 5x^2 = 6x^2 = 6x \cdot x$$

16.

$$x^2 + 14 = (x - \sqrt{14})(x + \sqrt{14})$$

Challenge

17.

$$(x + 5)(x - 1) = x^2 + 4x - 5$$

18.

$$x^2 - 9 = (x - 3)(x + 3) = (x - 3)(x + 3)$$

19.

$$x^3 + 1 = (x + 1)(x^2 - x + 1)$$

20. Expand.

$$(x - 2i)(x + 2i) = x^2 + 4$$

21. Expand.

$$x^2 + 4 = (x - 2i)(x + 2i)$$

Justify

The factoring must begin with the greatest common factor because removing it simplifies the expression, making it easier to factor the remaining terms.