9.A Graph Transformations of Rational Functions

Building on your understanding of parent functions, you'll now explore how to apply transformations—shifts, stretches, reflections—to rational functions. Rather than starting at zero for each problem, you'll learn to recognize how changes to the equation affect the graph's asymptotes, intercepts, and overall shape.

This skill is essential for analyzing real-world models, such as how a medication's concentration changes over time or how the cost per unit decreases as production increases.

Prerequisite concept structure — 2 class set review ⓘ

Learning Targets

- I can identify the parent function of a transformed rational function and describe the sequence of transformations.
- I can write a function that performs a specified transformation.

Introduction

Recurring student misstep ⓘ

The function is undefined at x = 0, and as $x \to \pm\infty$, $f(x) \to 0$. Its domain is $(-\infty, 0) \cup (0, \infty)$, and its range is $(0, \infty)$. The graph has two branches, one in Quadrant I and one in Quadrant II.

The function $f(x) = \frac{1}{x}$ is the simplest rational function. Its graph has a distinctive shape with two separate branches. Understanding this parent function is crucial because all other rational functions we'll study are transformations of it—shifted, stretched, reflected, or combinations of these. Once you know how $\frac{1}{x}$ behaves, you can predict how any transformed version will behave.

Key Idea: Transformations of rational functions follow the same rules you learned for other function families. Horizontal and vertical shifts move the graph and its asymptotes. Stretches and reflections change the shape and orientation. The key is tracking how each transformation affects the asymptotes and intercepts.

Connections

Prior Knowledge
- Function notation, domain, and transformation behavior
- Horizontal and vertical asymptotes as boundary behavior
- Graphing reciprocal functions and identifying key features

Cross-Disciplinary Connections
- **Physics:** Inverse relationships between variables, where one quantity decreases as the other increases
- **Economics:** Cost per unit models and average cost functions

Real-World Applications
- **Medicine:** Drug concentration in the bloodstream over time
- **Business:** Average cost per item as production volume changes
- **Engineering:** Electrical resistance in parallel circuits

Key Concepts

Vertical Shift of a Transformed Rational Function

$f(x) = \frac{1}{x} + k$ shifts the graph of $f(x) = \frac{1}{x}$ vertically k units.

- If $k > 0$, the graph shifts up k units.
- If $k < 0$, the graph shifts down $|k|$ units.
- The horizontal asymptote moves from $y = 0$ to $y = k$.

Horizontal Shift of a Transformed Rational Function

$f(x) = \frac{1}{x - h}$ shifts the graph of $f(x) = \frac{1}{x}$ horizontally h units.

- If $h > 0$, the graph shifts right h units.
- If $h < 0$, the graph shifts left $|h|$ units.
- The vertical asymptote moves from $x = 0$ to $x = h$.

Reciprocal Reference

The function $f(x) = \frac{1}{x}$ has a vertical asymptote at $x = 0$ and a horizontal asymptote at $y = 0$. Its graph has two branches, one in Quadrant I and one in Quadrant II. Each branch approaches the asymptotes but never touches them.

Examples

Example 1 — Horizontal Shift

Identify the vertical asymptote and x-intercept of $f(x) = \frac{1}{x - 3}$.

Example 2 — Reflection Across the x-axis

Identify the vertical asymptote and x-intercept of $f(x) = -\frac{1}{x}$.



Example 3 — Vertical Stretch

Identify the vertical asymptote and x-intercept of $f(x) = \frac{2}{x}$.

Example 4 — Combined Transformations

Identify the vertical asymptote and x-intercept of $f(x) = \frac{1}{x + 2} - 1$.



Example 5 — Reflection and Translation

Identify the vertical asymptote and x-intercept of $f(x) = -\frac{1}{x - 1}$.

Common Mistakes

- **Ignoring asymptotes:** Forgetting to move the asymptote when applying a shift. Always track how transformations affect the asymptotes.
- **Confusing horizontal and vertical shifts:** $f(x) = \frac{1}{x - 2}$ shifts right 2, not up 2. $f(x) = \frac{1}{x} + 2$ shifts up 2, not right 2.
- **Forgetting reflections:** A negative in front of the function reflects the graph across the x-axis.
- **Assuming the graph touches the asymptote:** Rational functions approach but never reach their asymptotes.

Summary

- Transformations of rational functions follow the same rules as other function families.
- Horizontal shifts move the function and its asymptote left or right.
- Vertical shifts move the function and its asymptote up or down.
- Reflections flip the graph across the x-axis or y-axis.
- Stretches and compressions change the steepness of the graph.
- Always identify how transformations affect the asymptotes and intercepts, not just the graph.

Practice Set

Level 1 — Warm

Problem 1

Problem 2

Problem 3

Problem 4

Problem 5

Level 2 — Practice

Problem 6

Problem 7

Problem 8

Problem 9

Problem 10

Level 3 — Stuck

Problem 11

Problem 12

Problem 13

Problem 14

Problem 15

Level 4 — Challenge

Problem 16

Problem 17

Problem 18

Problem 19

Problem 20

Level 5 — Justification / Conceptual

Problem 21 Explain why the graph of $f(x) = \frac{1}{x}$ never touches the x-axis or the y-axis.

Problem 22 Describe the sequence of transformations that takes the graph of $f(x) = \frac{1}{x}$ to the graph of $f(x) = \frac{1}{x + 3} - 2$.

Problem 23 Explain why the graph of $f(x) = \frac{1}{x - 4}$ has a vertical asymptote at $x = 4$ but no horizontal asymptote.

Problem 24 Compare the graphs of $f(x) = \frac{1}{x}$ and $g(x) = \frac{3}{x}$. Describe how they are similar and how they are different.

Problem 25 Explain why the graph of $f(x) = \frac{1}{x}$ has no x-intercept, using your own words.

Answer Key

Level 1 — Warm

Problem 1
$f(x) = \frac{1}{x} + 3$
$f(x) = \frac{1}{x}$ shifted up 3 units.

Problem 2
$f(x) = \frac{1}{x - 5}$
$f(x) = \frac{1}{x}$ shifted right 5 units.

Problem 3
$f(x) = -\frac{1}{x}$
$f(x) = \frac{1}{x}$ reflected across the x-axis.

Problem 4
$f(x) = \frac{4}{x}$
$f(x) = \frac{1}{x}$ stretched vertically by a factor of 4.

Problem 5
$f(x) = \frac{1}{x + 2} - 1$
$f(x) = \frac{1}{x}$ shifted left 2 units and down 1 unit.

Level 2 — Practice

Problem 6
$f(x) = \frac{1}{x - 1} + 4$
$f(x) = \frac{1}{x}$ shifted right 1 unit and up 4 units.

Problem 7
$f(x) = -\frac{1}{x + 3}$
$f(x) = \frac{1}{x}$ shifted left 3 units and reflected across the x-axis.

Problem 8
$f(x) = \frac{1}{x} - 2$
$f(x) = \frac{1}{x}$ shifted down 2 units.

Problem 9
$f(x) = \frac{1}{x - 6}$
$f(x) = \frac{1}{x}$ shifted right 6 units.

Problem 10
$f(x) = \frac{1}{x + 4} + 1$
$f(x) = \frac{1}{x}$ shifted left 4 units and up 1 unit.

Level 3 — Stuck

Problem 11
$f(x) = \frac{1}{x + 2} - 3$
$f(x) = \frac{1}{x}$ shifted left 2 units and down 3 units.

Problem 12
$f(x) = -\frac{1}{x - 1} + 2$
$f(x) = \frac{1}{x}$ shifted right 1 unit, reflected across the x-axis, and shifted up 2 units.

Problem 13
$f(x) = \frac{1}{x - 3}$
$f(x) = \frac{1}{x}$ shifted right 3 units.

Problem 14
$f(x) = \frac{1}{x} + 5$
$f(x) = \frac{1}{x}$ shifted up 5 units.

Problem 15
$f(x) = \frac{1}{x + 1} - 2$
$f(x) = \frac{1}{x}$ shifted left 1 unit and down 2 units.

Level 4 — Challenge

Problem 16
$f(x) = \frac{1}{x - 2} + 3$
$f(x) = \frac{1}{x}$ shifted right 2 units and up 3 units.

Problem 17
$f(x) = -\frac{1}{x + 4} - 1$
$f(x) = \frac{1}{x}$ shifted left 4 units, reflected across the x-axis, and shifted down 1 unit.

Problem 18
$f(x) = \frac{2}{x} + 1$
$f(x) = \frac{1}{x}$ stretched vertically by a factor of 2 and shifted up 1 unit.

Problem 19
$f(x) = \frac{1}{x - 5} - 4$
$f(x) = \frac{1}{x}$ shifted right 5 units and down 4 units.

Problem 20
$f(x) = \frac{1}{x + 3} + 2$
$f(x) = \frac{1}{x}$ shifted left 3 units and up 2 units.

Level 5 — Justification / Conceptual

Problem 21 The graph of $f(x) = \frac{1}{x}$ never touches the x-axis because there is no value of x that makes $\frac{1}{x} = 0$. Similarly, the graph never touches the y-axis because $x = 0$ makes the function undefined.

Problem 22 Shift left 3 units, then shift down 2 units.

Problem 23 The vertical asymptote is at $x = 4$ because that is where the denominator equals zero. There is no horizontal asymptote because the degree of the numerator is greater than the degree of the denominator.

Problem 24 Both graphs have the same shape and asymptotes. The graph of $g(x) = \frac{3}{x}$ is stretched vertically by a factor of 3 compared to $f(x) = \frac{1}{x}$, so it is steeper.

Problem 25 The function has no x-intercept because there is no value of x that makes the function equal zero. The numerator is always 1, which can never be zero.

Detailed Example Solutions

Problem 1 — Horizontal Shift

Given: $f(x) = \frac{1}{x - 3}$

Step 1: Identify transformation
The -3 in the denominator means the graph is shifted right 3 units.

Step 2: Identify the asymptote
The vertical asymptote is at $x = 3$.

Step 3: Find the x-intercept
The function never equals zero, so there is no x-intercept.

Final Answer: Vertical asymptote: $x = 3$; no x-intercept

Problem 2 — Vertical Translation

Given: $f(x) = \frac{1}{x} + 4$

Step 1: Identify transformation
The $+4$ means the graph is shifted up 4 units.

Step 2: Identify the asymptote
The horizontal asymptote is at $y = 4$.

Step 3: Find the x-intercept
Set $\frac{1}{x} + 4 = 0$
$\frac{1}{x} = -4$
$x = -\frac{1}{4}$

Final Answer: Horizontal asymptote: $y = 4$; x-intercept: $\left(-\frac{1}{4}, 0\right)$

Problem 3 — Reflection and Translation

Given: $f(x) = -\frac{1}{x + 2}$

Step 1: Identify transformations
Adding 2 shifts left 2 units.
The negative reflects across the x-axis.

Step 2: Identify the asymptote
The vertical asymptote is at $x = -2$.

Step 3: Find the x-intercept
Set $-\frac{1}{x + 2} = 0$
No solution, so no x-intercept.

Final Answer: Vertical asymptote: $x = -2$; no x-intercept

Problem 4 — Vertical Stretch

Given: $f(x) = \frac{3}{x}$

Step 1: Identify transformation
The 3 in the numerator means the graph is stretched vertically by a factor of 3.

Step 2: Identify the asymptote
The vertical asymptote is at $x = 0$.

Step 3: Find the x-intercept
Set $\frac{3}{x} = 0$
No solution, so no x-intercept.

Final Answer: Vertical asymptote: $x = 0$; no x-intercept

Problem 5 — Combined Transformations

Given: $f(x) = \frac{1}{x + 1} - 2$

Step 1: Identify transformations
Adding 1 shifts left 1 unit.
Subtracting 2 shifts down 2 units.

Step 2: Identify the asymptote
The vertical asymptote is at $x = -1$.

Step 3: Find the x-intercept
Set $\frac{1}{x + 1} - 2 = 0$
$\frac{1}{x + 1} = 2$
$x + 1 = \frac{1}{2}$
$x = -\frac{1}{2}$

Final Answer: Vertical asymptote: $x = -1$; x-intercept: $\left(-\frac{1}{2}, 0\right)$