

Problem 1.

Let X follows the exponential distribution with the rate parameter λ , then its (uncentered) MGF is

$$\mathbb{E}[\exp(tX)] = \frac{\lambda}{\lambda - t}, \quad \forall t < \lambda$$

and its mean is

$$\mathbb{E}[X] = \frac{1}{\lambda}$$

With the same r.v. X , we have

$$\begin{aligned} & \mathbb{E}[\exp(t(X - \mathbb{E}[X]))] \\ &= \frac{1}{\exp(t\mathbb{E}[X])} \mathbb{E}[\exp(tX)] \\ &= \frac{1}{\exp(\frac{t}{\lambda})} \mathbb{E}[\exp(tX)] \\ &= \frac{1}{\exp(\frac{t}{\lambda})} \frac{\lambda}{\lambda - t} \\ &= \frac{1}{\exp(\frac{t}{\lambda})} \left(1 + \frac{t}{\lambda - t}\right) \\ &\leq \frac{1}{\exp(\frac{t}{\lambda})} \exp\left(\frac{t}{\lambda - t}\right) \quad (\text{bc } 1 + x \leq \exp(x), \forall x) \\ &= \exp\left(\frac{t}{\lambda - t} - \frac{t}{\lambda}\right) \\ &= \exp\left(\frac{t^2}{\lambda(\lambda - t)}\right) \\ &= \exp\left(\frac{t^2 \frac{2}{\lambda^2}}{2(1 - \frac{1}{\lambda}t)}\right) \\ &\leq \exp\left(\frac{t^2 \frac{2}{\lambda^2}}{2(1 - \frac{1}{\lambda}|t|)}\right), \quad \forall |t| \leq \lambda \quad (\text{from the assumption of exponential distribution } t \leq \lambda) \end{aligned}$$

Therefore, the smallest parameters are $c = \frac{1}{\lambda}$ and $\nu^2 = \frac{2}{\lambda^2}$

Problem 3.**Problem 3.a.**

By definition, we have

$$Q = (1 - 2\varepsilon) \begin{pmatrix} 1 & -\tau \\ -\tau & \tau^2 \end{pmatrix} + \varepsilon \begin{pmatrix} 1 & \tau \\ \tau & \tau^2 \end{pmatrix} + \varepsilon \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}$$

We set $\varepsilon = \tau^2$, then

$$\begin{aligned} Q &= (1 - 2\tau^2) \begin{pmatrix} 1 & -\tau \\ -\tau & \tau^2 \end{pmatrix} + \tau^2 \begin{pmatrix} 1 & \tau \\ \tau & \tau^2 \end{pmatrix} + \tau^2 \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \\ &= \begin{pmatrix} 1 - \tau^2 & 3\tau^3 - r \\ 3\tau^3 - r & -\tau^4 + 2\tau^2 \end{pmatrix} \end{aligned}$$

Use wolfram alpha to find the Q^{-1} , we have

$$Q^{-1} = \frac{1}{8r^4 - 3r^2 - 1} \begin{pmatrix} \tau^2(\tau^2 - 2) & r(3\tau^3 - 1) \\ r(3\tau^3 - 1) & \tau^2 - 1 \end{pmatrix}$$

Problem 3.b.

From Q^{-1} , we have $(Q^{-1})_{11} = \frac{\tau^2(\tau^2-2)}{8r^4-3r^2-1}$