

# HW 1

## 1.1

We roll a fair die twice. Describe a sample space  $\Omega$  and a probability measure  $P$  to model this experiment. Let  $A$  be the event that the second roll is larger than the first. Find the probability  $P(A)$  that the event  $A$  occurs.

✓ Answer ✓

Let  $\alpha = \{1, 2, 3, 4, 5, 6\}$

Let  $\Omega = (i, j) : i, j \in \alpha$

Since  $\forall i \in \alpha : P(\{i\}) = \frac{1}{6}$  for a fair die,

Let  $P(\{(i, j)\}) = \frac{1}{36} : (i, j) \in \Omega$

First Roll	$P(A)$
1	5/6
2	4/6
3	3/6
4	2/6
5	1/6
6	0/6

With each of the states of the first roll having a  $\frac{1}{6}$  chance of occurring, the probability of the second roll being larger must be  $\frac{15}{36} = \frac{5}{12}$ .

---

$$P(A) = \frac{5}{12}$$

□

## 1.5

In one type of state lottery 5 distinct numbers are picked from  $1, 2, 3, \dots, 40$  uniformly at random.

**a**

Describe a sample space  $\Omega$  and a probability measure  $P$  to model this experiment.

✓ Answer

Assuming replacement of the number, and independence in drawings

Let  $\alpha = \{1, 2, 3, \dots, 40\}$

Let  $\Omega = \alpha^5$

Let  $A$  be when exactly 3 of the 5 numbers are even

Where  $\forall \omega \in \Omega : P(\{\omega\}) = \frac{1}{40^5}$

What is the probability that out of the 5 picked numbers exactly three will be even?

✓ Answer

$\forall a \in \alpha : P(a) = \frac{1}{40}$

Let  $\alpha' = \{a \text{ is even}, a \text{ is odd}\}$

$P(a \text{ is even}) = P(a \text{ is odd}) = \frac{1}{2}$  in  $\alpha'$  since  $\frac{20}{40}$  possibilities are even.

Let  $\Omega' = \alpha'^5$

Since the elements of  $\alpha'$  are equally probable, an independent combination of samples from  $\alpha'$  must also be equally probable.

$P(\{\omega\}) = \frac{1}{2^5} : \omega \in \Omega'$

There are  $5C3 = 10$  possible samples  $\omega \in \Omega'$  where 3 are even.

$P(A) = \frac{10}{32} = \frac{5}{16}$

---

$P(A) = \frac{5}{16}$

□

## 1.7

We have an urn with 3 green and 4 yellow balls. We draw 3 balls one by one without replacement.

**a**

Find the probability that the colors we see in order are green, yellow, green.

✓ Answer

$\frac{3}{7} \cdot \frac{4}{6} \cdot \frac{2}{5} = \frac{24}{210} = \frac{4}{35}$

□

**b**

Find the probability that our sample of 3 balls contains 2 green balls and one yellow ball.

✓ **Answer**

$$\text{GYG: } \frac{3}{7} \cdot \frac{4}{6} \cdot \frac{2}{5} = \frac{24}{210} = \frac{4}{35}$$

$$\text{YGG: } \frac{4}{7} \cdot \frac{3}{6} \cdot \frac{2}{5} = \frac{24}{210} = \frac{4}{35}$$

$$\text{GGY: } \frac{3}{7} \cdot \frac{2}{6} \cdot \frac{4}{5} = \frac{24}{210} = \frac{4}{35}$$

The order does not matter for picking the colors of 3 samples.

The probability of pulling any of those three combinations is  $\frac{12}{35}$

---

$$\frac{12}{35}$$

## 1.10

We roll a fair die repeatedly until we see the number four appear and then we stop. The outcome of the experiment is the number of rolls.

**a**

Following Example 1.16 describe a sample space  $\Omega$  and a probability measure  $P$  to model this situation.

✓ **Answer**

Let  $\Omega = \{\infty, 1, 2, \dots\}$

$$\forall \omega \in \Omega : P(\{\omega\}) = \left(\frac{5}{6}\right)^{\omega-1} \cdot \frac{1}{6}$$

**b**

Calculate the probability that the number four never appears.

✓ **Answer**

$$\lim_{\omega \rightarrow \infty} P(\{\omega\}) = \lim_{\omega \rightarrow \infty} \left(\frac{5}{6}\right)^{\omega-1} \cdot \frac{1}{6} = 0$$

---

$$P(\{\infty\}) = 0$$

## 1.12

We roll a fair die repeatedly until we see the number four appear and then we stop.

**a**

What is the probability that we need at most 3 rolls?

✓ **Answer**

The probability of needing 1, 2, or 3 rolls is equivalent to not needing more than rolls.

$$P(\omega > 3) = \left(\frac{5}{6}\right)^3$$

$$P(\omega \leq 3) = 1 - \left(\frac{5}{6}\right)^3 = \frac{91}{216}$$

---

$$P(\omega \leq 3) = \frac{91}{216}$$

□

**b**

What is the probability that we needed an even number of die rolls?

✓ **Answer**

Since the number of rolls are disjoint, we may add the probabilities directly.

$$P(\omega \text{ is even}) = P(\{2\}) + P(\{4\}) + P(\{6\}) + \dots$$

$$= \frac{5}{36} \left(1 + \frac{25}{36} + \left(\frac{26}{36}\right)^2 + \dots\right)$$

$$= \frac{5}{11} \text{ By the geometric sum rule}$$

---

$$P(\omega \text{ is even}) = \frac{5}{11}$$

## 1.20

We roll a fair die four times.

**a**

Describe the sample space  $\Omega$  and the probability measure  $P$  that model this experiment. To describe  $P$ , give the value  $P(\{\omega\})$  for each outcome  $\omega \in \Omega$ .

✓ **Answer**

Let  $\alpha = \{1, 2, \dots, 6\}$

Let  $\Omega = \alpha^4$

$$\forall \omega \in \Omega : P(\{\omega\}) = \frac{1}{6^4}$$

**b**

Let  $A$  be the event that there are at least two fives among the four rolls. Let  $B$  be the event that there is at most one five among the four rolls. Find the probabilities  $P(A)$  and  $P(B)$  by finding the ratio of the number of favorable outcomes to the total, as in Fact 1.8.

✓ **Answer**

Number of 5s	Number of Combinations
0	$1 \cdot 5^4$
1	$4 \cdot 5^3$
2	$6 \cdot 5^2$
3	$4 \cdot 5^1$
4	1

$$P(A) = \frac{171}{1296}$$

$$P(B) = \frac{1125}{1296}$$

□

**c**

What is the set  $A \cup B$ ? What equality should  $P(A)$  and  $P(B)$  satisfy? Check that your answers to part (b) satisfy this equality.

✓ **Answer**

Since  $A$  and  $B$  are complementary,  $P(A) + P(B) = 1$

$P(A) + P(B) = 1$  as checked.

□

## 1.22

We pick a card uniformly at random from a standard deck of 52 cards. (If you are unfamiliar with the deck of 52 cards, see the description above Example C.19 in Appendix C.)

**a**

Describe the sample space  $\Omega$  and the probability measure  $P$  that model this experiment.

✓ Answer

$$\Omega = \{\text{Diamonds, Clubs, Hearts, Spades}\} \times \{A, 2, 3, 4, 5, 6, 7, 8, 9, 10, J, Q, K\}$$

$$\forall \omega \in \Omega : P(\{\omega\}) = \frac{1}{52}$$

**b**

Give an example of an event in this probability space with probability  $\frac{3}{52}$

✓ Answer

$$P(\omega \in \text{Royal Spades}) : \omega \in \Omega$$

**c**

Show that there is no event in this probability space with probability  $\frac{1}{5}$ .

✓ Answer

Since there are 52 equal probability cards, any subset  $A \subset \Omega$  must contain an integer amount of elements less than or equal to 52.

We may sum over the cards in  $A$  as pulling any card is disjoint from pulling any other card.

$$P(A) = \sum_{i=1}^n P(\{A_i\}) \implies P(A) = \frac{|A|}{52}$$

$\frac{1}{5}$  is not a multiple of  $\frac{1}{52}$  and thus there cannot exist a set  $A$  such that  $\frac{|A|}{52} = \frac{1}{5}$ .

## 1.34

Pick a uniformly chosen random point inside a unit square (a square of sidelength 1) and draw a circle of radius  $1/3$  around the point. Find the probability that the circle lies entirely inside the square.

✓ Answer

We may divide the square into 9 equal squares, dividing by three vertically and horizontally.

Any circle centered in the center of the 9 squares will be entirely contained by the square, as the distance to its closest edge is at least  $\frac{1}{3}$ .

Any circle not centered on the center of the 9 squares will not be entirely contained, as the distance to the closest edge is less than  $\frac{1}{3}$ .

Since the area of each of the sub-squares are the same, they have the same probability of being sampled:

$$\Omega = \llbracket 1, 9 \rrbracket$$

$$\forall \omega \in \Omega : P(\{\omega\}) = \frac{1}{9}$$

$$P(\{5\}) = \frac{1}{9}$$