

Given $0 \leq x < \infty$; $\alpha, \beta > 0$

$$f(x) = \frac{x^{\alpha-1}e^{-x/\beta}}{\Gamma(\alpha)\beta^\alpha}$$
$$\mu = \alpha\beta$$
$$\sigma^2 = \alpha\beta^2$$
$$M(t) = (\frac{1}{1-\beta t})^\alpha$$

Logistic (μ, β)

Given $-\infty < x < \infty$; $-\infty < \mu < \infty$; $\beta > 0$

$$f(x) = \frac{e^{-(x-\mu)/\beta}}{\beta(1+e^{-(x-\mu)/\beta})^2}$$
$$\mu = \mu$$
$$\sigma^2 = \frac{\pi^2}{3}$$
$$M(t) = e^{\mu t}\Gamma(1+\beta t)$$

Lognormal (μ, σ^2)

Distribution where the log of a variable is normally distributed. Similar in appearance to the gamma distribution.

Given $0 \leq x < \infty$; $-\infty < \mu < \infty$; $\sigma > 0$

$$f(x) = \frac{1}{\sqrt{2\pi}\sigma} \frac{e^{-(\ln x - \mu)^2/(2\sigma^2)}}{x}$$
$$\mu = e^{\mu + (\sigma^2/2)}$$
$$\sigma^2 = e^{2(\mu + \sigma^2)} - e^{2\mu + \sigma^2}$$
$$EX^n = e^{n\mu + n^2\sigma^2/2}$$

Normal (μ, σ^2)

Can be used to approximate many different kinds of distributions as their population sizes increase.

Given $-\infty < x < \infty$; $-\infty < \mu < \infty$; $\sigma > 0$

$$f(x) = \frac{e^{-(x-\mu)^2/(2\sigma^2)}}{\sqrt{2\pi}\sigma}$$
$$\mu = \mu$$
$$\sigma^2 = \sigma^2$$
$$M(t) = e^{\mu t + \sigma^2 t^2/2}$$

Pareto (α, β)

Given $a < x < \infty$; $\alpha, \beta > 0$

$$f(x) = \frac{\beta\alpha^\beta}{x^{\beta+1}}$$
$$\mu = \frac{\beta\alpha}{\beta-1} \quad ; \beta > 1$$
$$\sigma^2 = \frac{\beta\alpha^2}{(\beta-1)^2(\beta-2)} \quad ; \beta > 2$$

T (v)

Given $-\infty < x < \infty$; $v = 1, 2, 3, \dots$

$$f(x) = \frac{\Gamma(\frac{v+1}{2})}{\Gamma(\frac{v}{2})} \frac{1}{\sqrt{v\pi}} \frac{1}{(1+(\frac{x^2}{v}))^{(v+1)/2}}$$
$$\mu = 0 \quad ; v > 1$$
$$\sigma^2 = \frac{v}{v-2} \quad ; v > 2$$
$$MX^n = \begin{cases} \frac{\Gamma(\frac{v+1}{2})\Gamma(\frac{v-n}{2})}{\sqrt{\pi}\Gamma(v/2)}v^{n/2} & n < v; n \text{ is even} \\ 0 & n < v; n \text{ is odd} \end{cases}$$

$$T(v) = \frac{N(0,1)}{\sqrt{\frac{1}{1v}}}$$

Uniform (a, b)

All values between a and b are evenly distributed and x has equal chance of landing anywhere on that range.

Given $a \leq x \leq b$

- a is the lower bound of the distribution
- b is the upper bound
- All values between a and b are equally distributed

$$f(x) = \frac{1}{b-a}$$
$$\mu = \frac{b+a}{2}$$
$$\sigma^2 = \frac{(b-a)^2}{12}$$
$$M(t) = \frac{e^{bt}-e^{at}}{t(b-a)}$$

Weibull (γ, β)

Given $0 \leq x < \infty$; $\gamma, \beta > 0$

$$f(x) = \frac{\gamma}{\beta}x^{\gamma-1}e^{-x^\gamma/\beta}$$
$$\mu = \beta^{1/\gamma}\Gamma(1+\frac{1}{\gamma})$$
$$\sigma^2 = \beta^{2/\gamma}(\Gamma(1+\frac{2}{\gamma})-\Gamma^2(1+\frac{1}{\gamma}))$$
$$EX^n = \beta^{n/\gamma}\Gamma(1+\frac{n}{\gamma})$$

Multivariable Distributions

Covariance and Correlation

$$\text{Cov}(X,Y) = E((X-\mu_X)(Y-\mu_Y))$$
$$\rho_{XY} = \frac{\text{Cov}(X,Y)}{\sigma_X\sigma_Y}$$

Multinomial Distribution

Given $\vec{\mu} \in \Re^k$; $\Sigma \in \Re^{k^2}$; $k \in \mathbb{N}$

$$f(\vec{x}) = (2\pi)^{-k/2} \det(\Sigma)^{-1/2} \exp\left(-\frac{1}{2}(\vec{x}-\vec{\mu})^T \Sigma^{-1}(\vec{x}-\vec{\mu})\right)$$
$$\vec{\mu} = \vec{\mu}$$
$$\sigma^2 = \Sigma$$
$$M(\vec{t}) = \exp\left(\vec{\mu}^T \vec{t} + \frac{1}{2} \vec{t}^T \Sigma \vec{t}\right)$$

Bivariate case ($\mu_X, \mu_Y, \sigma_X^2, \sigma_Y^2, \rho$)

$$f(x,y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp\left(-\frac{1}{2(1-\rho^2)}\left(\left(\frac{\bar{x}-\bar{\mu}_x}{\sigma_X}\right)^2 - 2\rho\left(\frac{\bar{x}-\bar{\mu}_x}{\sigma_X}\right)\left(\frac{\bar{y}-\bar{\mu}_y}{\sigma_Y}\right) + \left(\frac{\bar{y}-\bar{\mu}_y}{\sigma_Y}\right)^2\right)\right)$$

$$\vec{\mu} = \langle \mu_X, \mu_Y \rangle$$
$$\Sigma = \begin{bmatrix} \sigma_X^2 & \rho\sigma_X\sigma_Y \\ \rho\sigma_X\sigma_Y & \sigma_Y^2 \end{bmatrix}$$

Families of Distributions

Exponential Family

Exponential Family

Any statistical distribution or family of distributions that can fit into the form:

$$f(x|\theta) = h(x)c(\theta) \exp\left(\sum_{i=1}^k w_i(\theta)t_i(x)\right)$$

Binomial

$$\binom{n}{x}(1-p)^n \exp\left(\log\left(\frac{p}{1-p}\right)x\right)$$

Normal

$$\frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{\mu^2}{2\sigma^2}\right) \exp\left(-\frac{x^2}{2\sigma^2} + \frac{\mu x}{\sigma^2}\right)$$

Location Scale Family

Location Scale Family

Any statistical distribution or family of distributions that can fit into the form:

$$g(x|\mu,\sigma) = \frac{1}{\sigma} f\left(\frac{x-\mu}{\sigma}\right)$$

Probability Inequalities

Chebychev's Inequality

Let X be a random variable and let $g(x)$ be a non-negative function. Then, for any $r > 0$

$$P(g(X) \geq r) \leq \frac{Eg(X)}{r}$$

Normal Probability Inequality

With Z as a normal distribution,

Very similar to a binomial, except there is more than one possible outcome per trial — as compared to success or failure in the binomial.

- Taking the marginal of any of the possible outcomes results in a regular binomial

Multivariable Normal ($\vec{\mu}, \Sigma$)

$$P(|Z| \geq t) \leq \sqrt{\frac{e}{\pi}} \frac{e^{-t^2/2}}{t} \text{ for all } t > 0$$

Random Samples

Properties of the sample

$$\text{Mean: } \bar{X} = \frac{X_1 + \dots + X_n}{n} = \frac{1}{n} \sum_{i=1}^n X_i$$

$$\text{Variance: } S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

$$E(\sum_{i=1}^n g(X_i)) = nEg(X_1)$$

$$\text{Var}(\sum_{i=1}^n g(X_i)) = n\text{Varg}(X_1)$$

Properties of properties of the sample of random variables

$$E(\bar{X}) = \mu$$

$$\text{Var}(\bar{X}) = \frac{\sigma^2}{n}$$

$$E(S^2) = \sigma^2$$

Sample distributions of common distributions

Normal (μ, σ^2)

$$\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$$

S^2 is independent from \bar{X}

Gamma (α, β)

$$\bar{X} \sim \text{Gamma}\left(n\alpha, \frac{\beta}{n}\right)$$

Cauchy

$$\text{Cauchy}(0, \sigma_1) + \dots + \text{Cauchy}(0, \sigma_n) = \text{Cauchy}(0, \sum_{i=1}^n \sigma_i)$$

Chi-squared

$$\text{Chi}(a_1) + \dots + \text{Chi}(a_n) = \text{Chi}(\sum_{i=1}^n a_i)$$